Published in final edited form as:

J Ethn Subst Abuse. 2019 ; 18(2): 237–256. doi:10.1080/15332640.2017.1336956.

Psychometric evaluation of an abbreviated version of the Intragroup Marginalization Inventory

Jasmin Llamas, Ph.D^a, Louisa M. Holmes, Ph.D^b, Nadra E. Lisha, Ph.D^c, and Pamela M. Ling, M.D., MPH^c

^aSanta Clara University, School of Education and Counseling Psychology, CA, USA

^bBinghamton University, Department of Geography, Binghamton, NY, USA

^cCenter for Tobacco Control Research and Education and Division of General Internal Medicine, Department of Medicine. University of California San Francisco, CA, USA

Abstract

Tobacco and marijuana use among U.S. young adults is a top public health concern and racial/ethnic minorities may be at particular risk. Past research examining cultural variables has focused on the individual in relation to the mainstream U.S. culture, however an individual can also experience within-group stress, or intragroup marginalization. We used the 2014 San Francisco Bay Area Young Adult Health Survey to validate an abbreviated measure of intragroup marginalization and identify associations between intragroup marginalization and tobacco and marijuana use among ethnic minority young adults (N=1058). Exploratory Factor Analysis was conducted to identify factors within the abbreviated scale and logistic regressions were conducted to examine relationships between intragroup marginalization and tobacco and marijuana use. Two factors emerged from the abbreviated scale. The first factor encompassed items related to belonging and membership, capturing if individuals experienced marginalization due to not fitting in because of physical appearance or behavior. The second factor encompassed whether individuals shared similar hopes and dreams to their friends and family members. Factor 1 (membership) was associated with increased odds of marijuana use (OR = 1.34, $p < .05$) and lower odds of using cigars (OR = 0.79, $p < .05$), controlling for sociodemographic factors. Results suggest that young adults may use marijuana as a means to build connection and belonging to cope with feeling marginalized. Health education programs focused on ethnic minority young adults are needed to help them effectively cope with intragroup marginalization without resorting to marijuana use.

Keywords

ethnic minority; young adults; intragroup marginalization; tobacco; marijuana

Tobacco and marijuana use among U.S. young adults is a top public health concern (Chen & Jacobson, 2012); young adults use both substances at higher rates than any other age group

(Cohn, Villanti, Richardson, Rath, Williams, Stanton, & Mermelstein, 2015; Rath, Villanti, Abrams, & Vallone, 2012; Doll, Peto, Boreham, & Sutherland, 2004; Ling, Neilands, & Glantz, 2009). The transition to young adulthood increases the susceptibility to engaging in health-risk behaviors including tobacco and marijuana use (Pearso, Richardson, Niaura, Vallone, & Abrams, 2012; Rath et al. 2012). Non-cigarette tobacco product use (e.g., e-cigarettes, cigarillos, and hookah) has increased among young adults in the past several years (SAMHSA 2013; McMillen, Maduka, & Winickoff 2011; Soneji, Sargent, & Tanski, 2016; U.S. Department of Health and Human Services, 2014). Similarly, trends demonstrate an increase in marijuana use over the past decade (Hasin et al. 2015; U.S. Department of Health and Human Services 2014). In 2013, 37% of all young adults in the U.S. ages 18–25 reported currently using some form of tobacco (SAMHSA, 2013). Additionally, 19.1% of young adults reported using marijuana in the past 30 days in 2013 (SAMHSA, 2013). Tobacco use has been associated with cancer, heart disease, lung disease, and reproductive effects (U.S. Department of Health and Human Services, 2014), and marijuana use has been associated with cognitive impairment, cardiovascular disease, impaired respiratory functioning, and adverse psychosocial development and mental health (Hall & Degenhardt, 2009; NIDA 2016). Given the prevalence of these behaviors among young adults and their considerable health and social consequences, it is important to better understand factors associated with use, including whether shared cultural values, or feelings of marginalization with respect to such values may help explain high rates of use (Chen & Jacobson, 2012).

Among young adults, members of racial/ethnic minority groups are at especially high risk for certain types of tobacco use, such as cigarette and marijuana use, as well as higher rates of co-use (Holmes, Popova, & Ling, 2016; Ramo, Liu, and Prochaska, 2012; SAMHSA, 2013; U.S. Department of Health and Human Services, 2014; U.S. Department of Health and Human Services 1998; Wu, Swartz, Brady, Hoyle, & Workgroup, 2015). Additionally, racial/ethnic minorities suffer disproportionately from tobacco-related illness and death, despite having higher rates of light and intermittent smoking compared to Non-Hispanic Whites (Jemal et al. 2008; Trinidad, Perez-Stable, Emery, White, Grana, & Messer, 2009; U.S. Department of Health and Human Services, 1998). Racial and ethnic differences also exist in tobacco and marijuana use across groups. African American young adults have higher rates of current little cigar use compared to other racial/ethnic groups, while Latin young adult males report higher rates of current and lifetime e-cigarette use (Lariscy, Hummer, Rath, Villanti, Hayward, & Vallone, 2013). Current use of any tobacco product and marijuana use is highest for Mixed Race individuals (Holmes, Popova, and Ling, 2016; SAMHSA, 2013; Wu, Swartz, Brady, Hoyle, & Workgroup, 2015).

Race/ethnicity related stressors -- stressors that are a function of the cultural background and the context of the individual that are unique to being a member of a racial/ethnic minority group -- can make racial/ethnic minority young adults susceptible to tobacco and marijuana use (Kam, Cleveland, & Hecht, 2010; Williams, Neighbors, and Jackson 2003). For example, perceived racial/ethnic discrimination is a type of racial/ethnic stress that has been linked to increased smoking (Williams, Neighbors, and Jackson 2003) and higher odds of lifetime marijuana use (Borrell, Jacobs, Williams, Pletcher, Houston, Kiefe, & 2007). The National Conference on Tobacco and Health Disparities highlighted a need for researchers to examine the social and cultural context of tobacco use among racial/ethnic groups (Fagan,

King, Lawrence, Petrucci, Robinson, Banks, Marable, & Grana, 2004). Past research has also stressed the problematic perspective of viewing tobacco or marijuana use as an isolated problem, rather than being viewed as a part of a larger, more complicated picture that includes social and cultural components (Duff, 2003; Lunnay, Ward, & Borlagdan, 2011; Spooner, Hall, & Lynskey 2001). Additionally, health promotion researchers note that culturally specific interventions are important in addressing smoking-related health disparities. Culturally specific interventions refer to the degree to which ethnicity, attitudinal and behavioral norms, shared beliefs, history, and environment are integrated into the intervention (Resnicow, Baranowski, Ahluwalia, & Braithwaite, 1999). For example, the *Pathways to Freedom* is a smoking cessation guide developed for African Americans that incorporates known smoking patterns of African Americans, religious quotes, pictures of African Americans, and emphasizes family and community (Robinson, Orleans, James & Sutton, 1992).

Definitions of culture vary, but for the context of this paper, we focus on race/ethnicity, and the shared characteristics within these groups, which comprise religion, language, and nationality. The historical experiences of different racial/ethnic groups create unique physiological and social characteristics that can include lifestyle and value systems (Hays & Erford 2014; Napier et al., 2014). Past research examining cultural variables has primarily focused on racial/ethnic minority individuals in relation to the dominant culture, or mainstream U.S. culture, (i.e. discrimination, racism, acculturative stress), however an individual can also experience stress emanating from tensions within their own racial/ethnic group. This phenomenon, known as intragroup marginalization, refers to the perceived interpersonal distancing by members of one's racial/ethnic group when the individual diverges from racial/ethnic norms (Castillo, Conoley, Brossart, & Quiros, 2007). Deviating from racial/ethnic norms can create a backlash whereby group members reject or distance themselves from the individual. The interpersonal distancing occurring from intragroup marginalization can be viewed as a social sanction placed on the individual and can take the form of teasing and criticism. Intragroup marginalization is based on social identity theory (Tajfel & Turner 1986) suggesting that group members marginalize in-group members when they do not conform to group standards in order to maintain the uniqueness and stability of the group (Abrams, Marques, Bown, & Henson 2000). Group members displaying behaviors or attitudes that conflict with group norms can be perceived as threatening the distinctiveness of the group and can then be marginalized in order to preserve the group's distinctiveness.

Intragroup marginalization may be experienced by any racial/ethnic group. Additionally, family, friends, and other racial/ethnic members in the community can all impose group norms and engage in the process of intragroup marginalization. Limited research suggests intragroup marginalization may lead to higher levels of acculturative stress, or stress associated with adapting to a new culture, and increased alcohol use among young adults (Castillo, Cano, Chen, Blucker, & Olds, 2008; Castillo, Zahn, & Cano 2012; Llamas & Ramos-Sanchez 2013; Llamas & Morgan Consoli 2012). Past research, while not directly investigating intragroup marginalization, has made potential links between familial and peer stress with tobacco and marijuana use (e.g., Wills, Knight, Pagano, & Sargent 2015; Zapata Roblyer, Grzywacz, Cervantes, & Merten, 2016; Vitaro, Wanner, Brendgen, Gosselin, &

Gendreau, 2004). Foster and Spencer (2013) suggest that marijuana and other drug use may underlie a deeper need for connection in the absence of close familial connections for marginalized young adults, or young adults that have been rejected by their families. These young adults may be seeking opportunities to connect and create a sense of belonging, and marijuana use can play a common and significant social role in building supportive and caring relationships (Foster & Spencer 2013). Researchers further contend that investigation is needed to better understand how culture impacts these young adults' drug use (Foster & Spencer 2013).

Currently, intragroup marginalization is measured using the Intragroup Marginalization Inventory (Castillo et al. 2007), which is comprised of three separate scales measuring perceived intragroup marginalization from the heritage culture family (12-items), friends (17-items), and other members of the individual's ethnic group (13-items). The inventory is comprised of 42-items rated on a 7-point Likert scale (*never/does not apply [1] to extremely often [7]*). The scale items were developed so that the scale could be tailored to any ethnic group (e.g., 'Chinese friends tell me that I am not really Chinese because I don't act Chinese').

While the scale is comprehensive, the length of the survey can make it difficult for researchers to distribute the entire inventory, with many opting to use only one scale in their research (e.g., Castillo et al., 2008; Castillo, Zahn, & Cano, 2012; Llamas & Morgan Consoli, 2012; Llamas & Ramos-Sanchez, 2013). In practice this has limited studies of intragroup marginalization to focus either on family members or friends, rather than examining both. Due to the length of the survey, the feasibility of using the measure in large-scale studies or with large sample sizes has been limited. Most studies using the inventory have limited sample sizes focused on one racial/ethnic group (under 400 participants; e.g., Castillo et al., 2007; Llamas & Morgan Consoli, 2012; Llamas & Ramos-Sanchez, 2013). Greater sample sizes allow for segmentation of the data across demographic characteristics (i.e. race/ethnicity, gender, etc.), reduce the margin of error, and provide the statistical power to conduct more advanced analyses. In addition, some items may have less applicability for certain groups, such as items related to linguistic expectations (e.g., 'Family members criticize me because I don't speak my ethnic group's language.'). Lastly, the inventory was developed and validated with a college population and has not been validated with non-college populations (Castillo et al., 2007). Tobacco and marijuana use are problematic for all young adults and intragroup marginalization may be an important factor in understanding tobacco and marijuana disparities in this population as a whole. Yet, without an efficient means to assess intragroup marginalization, this important construct will continue to remain absent within health disparities research.

Current Study

Limited research addresses whether shared cultural values or feelings of marginalization may help explain high rates of tobacco and marijuana use among young adults (Chen & Jacobson 2012; Foster & Spencer 2013). The purpose of this study is to provide a psychometrically sound abbreviated measure of intragroup marginalization. Such a measure would have great utility when survey length is of concern and the survey needs to be

distributed across diverse racial/ethnic groups. This study tests and validates an abbreviated measure of the Intragroup Marginalization Inventory, which we refer to as the IMI-6. The IMI-6 consists of six items that measure perceived intragroup marginalization from the heritage culture family and friends. The items of the IMI-6 are hypothesized to have content validity, as items were taken directly from the existing scale, which has already been found to have content validity and were selected in consultation with the survey developer and by the primary author whose research focuses on racial/ethnic minority issues and intragroup marginalization in specific. We hypothesize that the IMI-6 also has construct validity, which we establish in this study through exploratory factor analyses. In addition to testing the feasibility of using this abbreviated measure, a primary aim of this study was to apply the IMI-6 and examine relationships between intragroup marginalization and tobacco and marijuana use. We hypothesize that participants reporting more experiences of intragroup marginalization would be more likely to use cigarettes, e-cigarettes, cigars, blunts, hookahs, and marijuana.

Method

Item selection

The original Intragroup Marginalization Inventory consists of three scales: Family, Friend, and Ethnic Group. The scales have a common factor structure, and while there are slight differences in items and factor names, they fall into five general factors: Homeostatic Pressure (pressure to not change), Linguistic Expectations (expectations that one speak the native language), and Accusations of Assimilation (accusations of adopting values and beliefs of White American culture), Accusations of Differentiation (accusations of looking or acting different), and Discrepant Values (values are too different from the group). The IMI-6 consists of six items that measure perceived intragroup marginalization from the heritage culture family and friends. The original scale developer provided consultation during item selection, ultimately reviewing and approving the final six items. Items were selected based on the researchers' and developer's experience with the survey as well as those items that had the greatest applicability to a diverse pool of respondents and were broad enough to remain appropriate for different racial/ethnic groups. Items from the Accusations of Assimilation and Linguistic Expectations factors were not included as they contained items that were tailored to specific racial/ethnic groups (i.e. an item from the Accusations of Assimilation was relevant only to Latina/os, "Friends from my ethnic group tell me that I am brown on the outside, but white on the inside"). Items from the Homeostatic Pressure were similar to items from the Accusations of Differentiation factor, however items from the Homeostatic Pressure focused solely on the individual's behavior, while items from the Accusations of Differentiation included items assessing both behavior *and* appearance. The selected items were taken from the Discrepant Values factor and the Accusation of Differentiation factor of the full inventory (see Table 1). Two items were taken from the Discrepant Values factor assessing whether family and friends have the same hopes and dreams as the respondent. Four items were taken from the Accusation Differentiation factor assessing whether family and friends accuse the respondent of not really being a member of one's ethnic group because s/he does not *look* like and *act* like members of the group. Responses were rated on a 7-point Likert scale, ranging from 'never/

does not apply' (1) to 'extremely often (7).' Items 3 and 6 were reverse coded, so that higher numbers represent greater experiences of intragroup marginalization. Items were piloted with 45 young adults (ages 18–26) from the San Francisco Bay Area. Participants were recruited from local bars on a Thursday, Friday and Saturday evening to be interviewed that same weekend and received a \$75 incentive if they participated in a one-hour focus group, completed the pilot questionnaire, and engaged in an interview with project staff to share feedback about the questionnaire. Individuals reviewed the item clarity and representation of their experiences. No items were altered and participant feedback suggested that the selected items accurately captured participant experiences.

Participants and procedure

Sample—This study used data we collected in 2014 as part of the San Francisco Bay Area Young Adult Health Survey, a probabilistic multi-mode household survey of 18–26 year old young adults, stratified by race/ethnicity (Holmes, Popova, & Ling 2016). The study was conducted in Alameda and San Francisco Counties in California. We identified potential respondent households using address lists from Marketing Systems Group (MSG; sample 1) in which there was an approximately 30–40% chance that an eligible young adult resided at a selected address ($n=15,000$ addresses). We used 2009–2013 American Community Survey and 2010 decennial census data in a multistage sampling design to identify Census Block Groups and then Census Blocks in which at least 15% of residents were Latino or non-Hispanic Black adults in the eligible age range. Ultimately, we randomly selected 61 blocks, then households within these blocks ($n=1,636$ housing units) then young adults within eligible households (sample 2). We oversampled these blocks because young nonwhite urban adults are among the most difficult populations to survey (Tourangeau, Edwards, Johnson, Wolter, & Bates, 2014), and we wished to ensure appropriate population representation.

We surveyed in three stages and utilized four modes of contact (mail, web, telephone, face-to-face). In the first stage we conducted a series of three mailings with sample 1 households; respondents returned paper questionnaires or completed surveys online using Qualtrics. In the second stage we telephoned those who did not respond to mail, and lastly we performed face-to-face interviews with a random selection of the remaining nonresponders ($n \approx 1,250$) from sample 1 as well as all of the households identified in sample 2. Potential sample 2 respondents did not participate in the mail or telephone phases of the survey; each of these households was visited in person. The final sample consisted of 1,363 young adult participants, for a response rate of approximately 30%, with race, sex and age distributions closely reflecting those of the young adult population overall in the two counties surveyed. Ethnicity/race was measured using items from the Census Bureau's American Community Survey instrument, with participants first asked to identify if there were Hispanic, Latino, or Spanish origin and then to select their race from 14 categories. Race/ethnicity was then collapsed into mutually exclusive categories including Hispanic, non-Hispanic White, non-Hispanic black, non-Hispanic Asian/Pacific Islander and Mixed Race. Those who selected more than one race/ethnic category (e.g. Black and Latino; Japanese and White, etc.) were categorized as Mixed Race. We constructed individual sample and post-stratification adjustment weights during data reduction (Holmes, Popova, and Ling 2016).

Measures

Outcomes—We evaluated associations between intragroup marginalization and current use of cigarettes, cigars, blunts (hollowed out cigars filled with marijuana), hookah, e-cigarettes and marijuana. Each outcome measure was dichotomized and set equal to ‘1’ if a respondent reported using the product in question at least once in the past 30 days.

Main Explanatory Variables

Covariates: Age in years since birth was measured continuously (18–26), sex was measured dichotomously with male set equal to ‘1’ and female to ‘0’, and maternal education was set equal to ‘1’ if the respondent’s mother had completed at least a bachelor’s degree and ‘0’ otherwise. Race/ethnicity was measured as an indicator variable with mutually exclusive categories including Hispanic, non-Hispanic black, non-Hispanic Asian/Pacific Islander and Mixed Race (those who identified as two or more races). We restricted our analysis to young adults in these categories, excluding non-Hispanic white as the intragroup marginalization inventory has only been used and validated among nonwhite populations previously and endorsement of intragroup marginalization was not expected among this population (Castillo et al., 2007). The resulting number of observations was 1058, or 78% of the total sample.

Statistical Analysis

To examine the items in the abbreviated measure we conducted an exploratory factor analysis (EFA). Due to the exploratory nature of our analysis we chose to conduct an EFA rather than confirmatory factor analysis (CFA). CFA is useful to extract latent factors from a set of items based on an a priori theory. This requires a strong empirical or conceptual foundation and a pre-specification of the number of factors pattern of factor loadings. As we are using these items in a relatively innovative fashion, we wanted to determine, without specifying a structure, how the items were related. We conducted an EFA using an oblique geomin rotation (Fabrigar et al. 1999) in *Mplus*. EFA methods typically follow “rules of thumb,” with factor loading cutoff criteria ranging from .30 to .55, to establish a solid factor loading coefficient (Swisher, Beckstead, & Bebeau, 2004); we used a cutoff value of .55 in this study. The number of factors retained was based on eigenvalues > 1. Internal consistency was examined by computing Cronbach’s α for the entire measure and each subscale. Second, we fit multinomial logistic regression models using SAS SURVEYLOGISTIC (SAS, 2008) to account for the complex survey design. This was repeated with six dichotomous outcomes (cigarette use, e-cigarette use, blunt/wraps, hookah and marijuana use,) in two steps: 1) unadjusted analysis (factors were the sole predictors), and 2) controlling for race/ethnicity, age, sex and mother’s highest education.

Results

Sample information

Weighted percentages (or means) and standard error of percent (or standard error of the mean) are presented in Table 2. Approximately one-third of the sample retained for analysis was Latino, 40% was non-Hispanic Asian/Pacific Islander, 15% was non-Hispanic Black and the remaining 10% reported being of two or more races. Close to half of all participants

endorsed feeling marginalized by friends because they did not look (43%) or act (49%) like members of their racial/ethnic group. Approximately a quarter of participants endorsed feeling marginalized by family members because they did not look (23%) or act (27%) like members of their racial/ethnic group. Most participants reported having similar hopes and dreams as their friends (95%) and family (84%).

Exploratory Factor Analysis and Internal Consistency

The EFA indicated two factors (eigenvalue factor 1 = 2.970, eigenvalue factor 2 = 1.591, and eigenvalue factor 3 = 0.688). As shown in Table 3, every item loaded above 0.60 on at least one factor. Factor 1 might be described as looking or acting like your ethnic group and was composed of items 'Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't look like my ethnic group,' 'Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't act like my ethnic group,' 'Family members tell me I am not really a member of my ethnic group because I don't look like my ethnic group,' and 'Family members tell me I am not really a member of my ethnic group because I don't act like my ethnic group.' Factor 2 appears to represent hopes and dreams and was composed of the two items 'Friends and peers in my ethnic group have the same hopes and dreams as me,' and 'My family has the same hopes and dreams as me.' The Cronbach's α for the entire IMI-6 was 0.66. Cronbach's α s were computed for each subscale and found to be: Factor 1, .81; and Factor 2, 0.71.

Mean IMI factor scores by race/ethnicity are presented in Table 4. A regression analysis was conducted to determine mean differences in IMI factor scores by race/ethnicity (Table 5). A significant difference in means scores was found by race/ethnicity for Factor 1, $F(8, 1050) = 20.02$, $p < .001$, $R^2 = 0.04$. Latinos had greater mean scores for Factor 1 than Non-Hispanic Blacks [$t(8) = 4.43$, $p < .01$] and Non-Hispanic Asian/Pacific Islander [$t(8) = 3.81$, $p < .01$]. Mixed Race individuals had greater mean scores for Factor 1 than Non-Hispanic Blacks [$t(8) = -5.00$, $p < .01$] and Non-Hispanic Asian/Pacific Islander [$t(8) = -2.99$, $p < .05$]. No significant difference in means scores was found by race/ethnicity for Factor 2, $F(8, 1050) = 1.81$, $p = 0.22$, $R^2 = 0.005$.

IMI-Discriminant Validity

Non-Hispanic whites were not expected to report intragroup marginalization and not included in the factor and regression analyses as the full inventory has only been used and validated among nonwhite populations. To demonstrate the discriminant validity of the measure two t-tests were conducted. Non-Hispanic Whites were compared to the rest of the sample on the two factors. Non-Hispanic Whites experienced significantly lower scores on discrimination for both factor 1 (1.26 vs. 1.92, $p < .0001$) and factor 2 (3.26 vs. 3.71, $p < .0001$).

IMI-6 in Unadjusted Logistic Regressions

Results varied by outcome such that no significant relationship was found between the two factors and cigarette use, e-cigarette use or blunt use. However, Factor 1 was related to hookah, marijuana and cigar use. Higher scores on Factor 1 were related to higher odds of hookah use (OR = 1.26, 95%CI = 1.07, 1.48) and marijuana use (OR = 1.37, 95%CI = 1.05,

1.79), but lower odds of cigar use (OR = 0.81, 95%CI = 0.70, 0.93). Factor 2 was related to lower odds of hookah use (OR = .85, 95%CI = 0.72, 0.99).

IMI-6 in Multinomial Logistic Regressions

When controlling for race/ethnicity, age, sex and mother's education the results were consistent with the unadjusted models, except that Factor 1 and 2 were no longer associated with hookah use. No significant relationships were found for cigarettes, e-cigarettes, or blunts. When adjusting for covariates, Factor 1 and 2 were no longer associated with hookah use. However, the associations with marijuana and cigar use were robust; Factor 1 was associated with increased odds of marijuana use (OR = 1.34, 95%CI = 1.02, 1.76), and lower odds of using cigars (OR = 0.79, 95%CI = 0.71, 0.87).

Discussion

Results support the use of an abbreviated measure of intragroup marginalization. The IMI-6 was found to be psychometrically sound and representative of the full construct of intragroup marginalization as theorized by Castillo and colleagues (2007). Two factors emerged from the abbreviated scale. The first factor encompassed items related to belonging and membership, capturing whether individuals felt marginalized due to deviations in their physical appearance or behaviors (i.e., hobbies, interests). The second factor encompassed whether the individual shared similar hopes and dreams as their families and friends. These factors reflected similarly identified factors from the validation study of the full inventory scales (Castillo et al., 2007), suggesting good agreement between the original measure and the abbreviated version.

Examining racial/ethnic differences in mean scores across factors demonstrated significant differences in Factor 1. Latinos and Mixed Race young adults experienced greater intragroup marginalization related to not looking or acting like members of their racial/ethnic group compared to non-Hispanic Blacks and Asian Americans/ Pacific Islanders. The full Intragroup Marginalization Inventory (Castillo et al., 2007) was developed with a diverse sample (Asian American, Black/ African American, Latino, Native American and Biracial) and past research has explored intragroup marginalization with African Americans (e.g. Thompson et al. 2010), Asian Americans (e.g., Castillo et al., 2012) and Latinos (e.g., Castillo et al., 2008); however, specific racial/ethnic differences have not been examined.

Latinos may be particularly susceptible to intragroup marginalization given the heterogeneity among Latinos in terms of national origin, physical appearance, political ideology, immigration status, and class status (Fry, 2002; Johnson, Farrell, & Guinn, 1997). In particular, Latinos can encompass different racial groups (i.e. Afro-Latino, Asian Latino, etc.), which can contribute to differences in appearance one of the concepts captured in Factor 1. Physical appearance can limit the extent to which people are accepted as belonging to a certain racial/ethnic group, which is also especially relevant for multiracial individuals, whose physical appearance may not align with any specific ethnic/racial group (Ahn Allen, Suyemoto, & Carter, 2006). Additionally, multiracial individuals describe feeling marginalized from peers rooted in having different appearance, culture, and/or beliefs than

their peers (Jackson, 2010), explaining the higher rates of intragroup marginalization observed in this study.

Research examining young adult tobacco and marijuana use often relies on college samples, thereby neglecting individuals in this age group that may be at greater risk of substance use (e.g., Moran, Wechsler, & Rigotti, 2004; Morrell, Cohen, Bacchi, & West, 2005; National Cancer Institute, 2008; Rigotti, Lee, & Wechsler, 2000). The Intragroup Marginalization Inventory, which may have particular utility with young adults who are negotiating the stresses of transitioning to adulthood, was also developed and tested with a college-only sample (Ferenczi & Marshall, 2014). This study validates an abbreviated version of the IMI, the IMI-6, which was developed to capture tensions experienced within racial/ethnic groups. We tested the IMI-6 in a large representative household sample of racially/ethnically diverse young adults in the San Francisco Bay Area in order to better understand the impact of cultural stressors on tobacco and marijuana use among young adults in general.

When controlling for demographic characteristics, Factor 1 (membership) was associated with greater marijuana use. Participants who felt that they did not look or act like members of their racial/ethnic group demonstrated increased odds of marijuana use. Young adults who feel marginalized by family members or friends may seek to find a way to belong and connect with other young adults and marijuana use may be a way to find belonging within a group. This parallels research, which suggests that the decision to engage in marijuana use comes from an internal need for emotional connection and friendship (Pilkington, 2007) and as an opportunity to connect and create a sense of belonging (Foster & Spencer, 2013). Other research has identified marijuana as a more acceptable substance viewed as superior and safer than other substances (Foster & Spencer, 2013). Marijuana may be the substance of choice to build connection with others and combat feelings of intragroup marginalization.

If marijuana use is perceived as a means for social connection, it may help to explain the findings between Factor 1 (membership) and cigar use. When controlling for demographic characteristics, participants who felt as though they did not look or act like members of their racial/ethnic group had decreased odds of cigar use. Cigars were the least frequently used product within the sample retained for analysis. National averages parallel this trend with current cigar use (10%) having lower prevalence than cigarettes (31%) and marijuana (19%) for young adults (SAMHSA, 2013). If marginalized young adults seek to connect with others via substance use, cigar use may not be the best mechanism by which to connect with others and therefore they may be less likely to use cigars. The combination of low rates of use and potential lack of opportunity to build social connection may help explain the decreased odds of cigar use. This finding is unexpected and further research is needed to better understand the relationship between intragroup marginalization and cigar use.

Similarly, cigarette, e-cigarette, blunt use, and hookah use had lower rates compared to marijuana use. While unexpected, cigarette, e-cigarette, and blunt use were not associated with experiences of intragroup marginalization. This may be due in part to the lower rates of use. It is worth noting that blunt use was examined independently, although it is often associated with marijuana use and in this sample most blunt users also reported concurrent marijuana use (104 of 109 blunt users). Additionally, the use of these substances may be less

Author Manuscript

tied to social use and therefore their use may not be linked to developing ways of belonging. Past research has differentiated between ‘social smoking’ and smoking alone (Moran, Wechsler, & Rigotti, 2004). Studies have suggested that young adults not in college may be less likely to be social smokers (Johnston, O’Malley, & Bachman, 2002) and social smoking may not be prevalent across racial/ethnic groups (Moran, Wechsler, & Rigotti, 2004). This study did not differentiate between social smoking and smoking alone and may be another important factor to better understand the role of intragroup marginalization and tobacco use.

Author Manuscript

Intragroup marginalization was associated with higher hookah use; however, when controlling for race/ethnicity this association was no longer significant due to racial/ethnic differences. While hookah use has been noted as a means for socializing (Braun, Glassman, Wohlwend, Whewell, & Reindl, 2012) and is often smoked in a group setting (Ward, Eissenberg, Gray, Srinivas, Wilson, & Maziak, 2007), this may be population specific. Hookah use is common in Middle Eastern countries and has strong cultural underpinnings (Jamil, Elsouhag, Hiller, Arnetz, & Arnetz, 2010). Middle Eastern young adults experiencing intragroup marginalization may use hookah as a means to connect and fit in within their cultural group. Furthermore, African Americans have been found to have lower rates of use compared to other ethnic/racial groups (Barnett, Smith, He, Soule, Curbow, Tomar, & McCarty, 2013; Primack, Fertman, Rice, Adachi-Mejia, & Fine, 2010). Additional research may be needed to further investigate differential impacts of intragroup marginalization on hookah use ethnic/racial group.

Author Manuscript

Factor 1 (membership) captures the challenges young adults face when they feel they do not fit in with members of their ethnic/racial group. Young adults experiencing this may desire to find ways to gain membership and connection with others. While Factor 1 focuses on difficulties in belonging and membership, Factor 2 centered on shared values and dreams. Feeling marginalized due to a lack of similar hopes and dreams was not associated with tobacco or marijuana use. This finding supports the theory that young adults use these substances as a means of building belonging and connection (Foster & Spencer 2013). While having dissimilar hopes and dreams may be stressful, it may not necessarily indicate one does have any connection to others. Given these findings, the scale may be able to be further abbreviated by dropping Factor 2, particular when examining tobacco and marijuana use. Future research may be needed to further investigate impacts of Factor 2 on other health outcomes.

Author Manuscript

Despite the strengths of this research, there are important limitations to note. This study focused on young adults in the San Francisco Bay Area, and findings may be not be generalizable to all young adults. However by using population-based sampling, we were able to obtain a representative sample, which past research has noted the difficulty in reaching urban young adults (Holmes et al., 2016). This study also utilized a cross-sectional design, preventing any potential inference concerning causality. Tobacco and marijuana use were measured using self-report data and use was not biochemically verified. While past research has demonstrated the reliability and validity of self-reported smoking in anonymous surveys with young adults (Ramo, Hall, & Prochaska, 2011) this validation has not extended to non-cigarette tobacco products; this may be a potential area for future research. This study examined intragroup marginalization among Mixed Race young adults; a population often

overlooked in intragroup marginalization studies. Mixed Race participants were not required to identify which group served as the primary source of intragroup marginalization. However, it is possible that different cultural norms around tobacco and marijuana use could influence whether intragroup marginalization impacted behavior. Oyserman and colleagues (2007) have demonstrated the identity-based motivation of health behaviors, with racial/ethnic minorities more likely to identify unhealthy behaviors with their group. Additional research may be needed with Mixed Race individuals to better understand how different groups may impact the relationships between intragroup marginalization and tobacco use. A final limitation is that we did not directly assess reasons or motivations for use. Future qualitative research is needed to explicitly examine motivations for use as a result of experiences of intragroup marginalization.

This study provides the first quantitative examination of intragroup marginalization with tobacco and marijuana use. Results respond to recent calls to better understand motivations for young adult marijuana use (Holmes et al., 2016), with findings demonstrating an association between intragroup marginalization and increased marijuana use. These findings are especially relevant given the changing climate regarding the legalization of marijuana, with California just recently voting to legalize marijuana (NORML, 2016). Results reaffirm existing arguments that drug policy must attend to the social and cultural contexts of use (Duff, Moore, Johnston, & Goren, 2007; Foster & Spencer, 2013). Additionally, findings respond to existing calls in the literature to better understand how culture impacts use (Foster & Spencer 2013). Past intervention research has highlighted the importance of attending to peer smoking behavior and norms, providing further support for the need to attend to social dynamics when addressing young adult tobacco and marijuana use (Kalkhoran, Lisha, Neilands, Jordan, & Ling, 2016). Additional research is needed to further investigate the relationship between intragroup marginalization and marijuana use, which can help in the tailoring and development of targeted health education programs.

References

- Abrams D, Marques JM, Bown N, & Henson M. (2000). Pro-norm and anti-norm deviance within and between groups. *Journal of Personality and Social Psychology*, 78(5), 906. doi: 10.1037/0022-3514.78.5.906 [PubMed: 10821197]
- AhnAllen JM, Suyemoto KL, & Carter AS (2006). Relationship between physical appearance, sense of belonging and exclusion, and racial/ethnic self-identification among multiracial Japanese European Americans. *Cultural Diversity and Ethnic Minority Psychology*, 12(4), 673–686. doi: 10.1037/1099-9809.12.4.673 [PubMed: 17087528]
- Barnett TE, Smith T, He Y, Soule EK, Curbow BA, Tomar SL, & McCarty C (2013). Evidence of emerging hookah use among university students: a cross-sectional comparison between hookah and cigarette use. *BMC public health*, 13(1), 302–309. doi:10.1186/1471-2458-13-302 [PubMed: 23560649]
- Borrell LN, Jacobs DR, Williams DR, Pletcher MJ, Houston TK, & Kiefe CI . (2007). Self-reported racial discrimination and substance use in the Coronary Artery Risk Development in Adults Study. *American Journal of Epidemiology*, 166(9), 1068–1079. [PubMed: 17698506]
- Braun RE, Glassman T, Wohlwend J, Whewell A, & Reindl DM . (2012). Hookah use among college students from a Midwest University. *Journal of community health*, 37(2), 294–298. [PubMed: 21805373]

- Castillo LG, Cano MA, Chen SW, Blucker RT, & Olds TS . (2008). Family conflict and intragroup marginalization as predictors of acculturative stress in Latino college students. *International Journal of Stress Management*, 15(1), 43–52.
- Castillo LG, Conoley CW, Brossart DF, & Quiros AE (2007). Construction and validation of the Intragroup Marginalization Inventory. *Cultural Diversity and Ethnic Minority Psychology*, 13(3), 232–240. doi:10.1037/1099-9809.13.3.232 [PubMed: 17638480]
- Castillo LG, Zahn MP, & Cano MA (2012). Predictors of familial acculturative stress in Asian American college students. *Journal of College Counseling*, 15(1), 52–64.
- Chen P, & Jacobson KC (2012). Developmental trajectories of substance use from early adolescence to young adulthood: gender and racial/ethnic differences. *Journal of Adolescent Health*, 50(2), 154–63. doi: 10.1016/j.jadohealth.2011.05.013. [PubMed: 22265111]
- Cohn A, Villanti A, Richardson A, Rath JM, Williams V, Stanton C, & Mermelstein R (2015). The association between alcohol, marijuana use, and new and emerging tobacco products in a young adult population. *Addictive behaviors*, 48, 79–88. [PubMed: 26042613]
- Doll R, Peto R, Boreham J, & Sutherland I (2004). Mortality in relation to smoking: 50 years' observations on male British doctors. *Bmj*, 328(7455), 1519. [PubMed: 15213107]
- Duff C (2003). Drugs and youth cultures: Is Australia experiencing the “nor- malization” of adolescent drug use? *Journal of Youth Studies*, 6(4), 433–447. doi:10.1080/1367626032000162131
- Fabrigar LR, Wegener DT, MacCallum RC, & Strahan EJ (1999). Evaluating the use of exploratory factor analysis in psychological research. *Psychological methods*, 4(3), 272.
- Fagan P, King G, Lawrence D, Petrucci SA, Robinson RG, Banks D, Marable S, & Grana R (2004). Eliminating tobacco-related health disparities: Directions for future research. *American Journal of Public Health*, 94(2), 211–217. [PubMed: 14759929]
- Ferenczi N, & Marshall TC (2014). Meeting the expectations of your heritage culture Links between attachment orientations, intragroup marginalization and psychological adjustment. *Journal of social and personal relationships*, 0265407514562565.
- Foster K & Spencer D (2013). ‘It’s just a social thing’: Drug use, friendship and borderwork among marginalized young people. *International Journal of Drug Policy*, 24, 223–230. [PubMed: 23352335]
- Fry R (2002). Latinos in Higher Education: Many Enroll, Too Few Graduate. Retrieved from Washington, DC: Pew Hispanic Center.
- Hall W, & Degenhardt L. (2009). Adverse health effects of non-medical cannabis use. *The Lancet*, 374(9698), 1383–1391.
- Hays DG .and Erford BT.2014 Developing Multicultural Counseling Competence: A Systems Approach. Upper Saddle River, NJ: Pearson.
- Hasin DS, Saha TD, Kerridge BT, Goldstein RB, Chou SP, Zhang H, ... & Grant BF . (2015). Prevalence of marijuana use disorders in the United States between 2001–2002 and 2012–2013. *JAMA Psychiatry*, 72(12), 1235–1242. [PubMed: 26502112]
- Hays DGE, (2014). Developing multicultural counseling competence: A systems approach. Upper Saddle River, NJ: Pearson.
- Holmes LM, Popova L, & Ling PM (2016). State of transition: Marijuana use among young adults in the San Francisco Bay Area. *Preventive Medicine*, 90, 11–16. doi:10.1016/j.ypmed.2016.06.025 [PubMed: 27346757]
- Jackson KF (2010). Living the multiracial experience: Shifting racial expressions, resisting race, and seeking community. *Qualitative Social Work*, 1473325010375646.
- Jamil H, Elsouhag D, Hiller S, Arnetz JE, & Arnetz BB . (2010). Sociodemographic risk indicators of hookah smoking among White Americans: A pilot study. *Nicotine & Tobacco Research*(12), 525–529. doi:10.1093/ntq026
- Jemal A, Thun MJ, Ries LA, Howe HL, Weir HK, Center MM, . . . Edwards BK (2008). Annual report to the nation on the status of cancer, 1975–2005, featuring trends in lung cancer, tobacco use, and tobacco control. *J Natl Cancer Inst*, 100(23), 1672–1694. doi:10.1093/jnci/djn389 [PubMed: 19033571]

- Johnson JH Jr, Farrell WC Jr, & Guinn C (1997). Immigration reform and the browning of America: Tensions, conflicts and community instability in metropolitan Los Angeles. *International Migration Review*, 1055–1095.
- Johnston L OMP, Bachman J (2002). *Monitoring the Future: National Survey Results on Drug Use, 1975–2001* (Vol. Volume II: College Students and Adults Ages 19 – 40). Bethesda, MD: National Institute on Drug Abuse.
- Kalkhoran S, Lisha NE, Neilands TB, Jordan JW, & Ling PM . (2016). Evaluation of Bar and Nightclub Intervention to Decrease Young Adult Smoking in New Mexico. *Journal of Adolescent Health*.
- Kam JA CM, Hecht ML . (2010). Applying general strain theory to examine perceived discrimination's indirect relation to Mexican-heritage youth's alcohol, cigarette, and marijuana use. *Prevention Science*, 11(4), 397–410. [PubMed: 20490921]
- Lariscy JT, Hummer RA, Rath JM, Villanti AC, Hayward MD, & Vallone DM (2013). Race/ethnicity, nativity, and tobacco use among US young adults: Results from a nationally representative survey. *Nicotine & Tobacco Research*, nts344.
- Ling PM, Neilands TB, & Glantz SA (2009). Young adult smoking behavior: A national survey. *American journal of preventive medicine*, 36(5), 389–394.e382. doi:10.1016/j.amepre.2009.01.028 [PubMed: 19269128]
- Llamas J, & Ramos-Sánchez L (2013). Role of peer support on intragroup marginalization for latino undergraduates. *Journal of Multicultural Counseling and Development*, 41(3), 158–168.
- Llamas JD, & Morgan Consoli M (2012). The importance of familia for Latina/o college students: examining the role of familial support in intragroup marginalization. *Cultur Divers Ethnic Minor Psychol*, 18(4), 395–403. doi:10.1037/a0029756 [PubMed: 22985244]
- Lunnay B, Ward P, & Borlagdan J . (2011). The practise and practice of Bourdieu: The application of social theory to youth alcohol research. *International Journal of Drug Policy*, 22(6), 428–436. doi: 10.1016/j.drugpo.2011.07.013 [PubMed: 21862305]
- McMillen R, Maduka J, & Winickoff J . (2012). Use of emerging tobacco products in the United States. *Journal of Environmental and Public Health*, 1–8.
- Moran S, Rigotti N. A. WH . (2004). Social smoking among US college students. *Pediatrics*(114), 1028–1034. doi:10.1542/peds.2003-0558-L [PubMed: 15466101]
- Morrell HER, Bacchi D. CLM, West J . (2005). Predictors of smoking and smokeless tobacco use in college students: A preliminary study using web-based survey methodology. *Journal of American College Health*(54), 108–115. doi:10.3200/JACH.54.2.108-115 [PubMed: 16255323]
- Napier AD, Arcano C, Butler B, Calabrese JD, Chater A, Chatterjee H, ... & Wilson KW . (2014). Culture and health. *The Lancet*, 384(9954), 1607–1639.
- National Cancer Institute (NCI). (2008). *The Role of Media in Promoting and Reducing Tobacco Use*. Bethesda (MD): U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute.
- National Institute on Drug Addiction (NIDA). (2016). Marijuana. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/marijuana>
- NORML. (2016). State Marijuana Laws. Retrieved from <http://norml.org/laws>
- Oyserman D, Fryberg SA, & Yoder N. (2007). Identity-based motivation and health. *Journal of Personality and Social Psychology*, 93(6), 1011–27. [PubMed: 18072851]
- Pearson JL, Richardson A, Niaura RS, Vallone DM, & Abrams DB (2012). e-Cigarette awareness, use, and harm perceptions in US adults. *American Journal of Public Health*, 102(9), 1758–1766. [PubMed: 22813087]
- Pilkington H. (2007). In good company: Risk, security and choice in young people's drug decisions. *The Sociological Review*, 55(2), 373–392. doi:10.1111/j.1467-954X.2007.00710.x.
- Primack BA FC, Rice K, Adachi-Mejia A, Fine M. (2010). Waterpipe and cigarette smoking among college athletes in the United States. *J Adolescent Health*, 2010(46), 45–51.
- Ramo DE, Hall SM, & Prochaska JJ . (2011). Reliability and validity of self-reported smoking in an anonymous online survey with young adults. *Health Psychology*, 30(6), 693. [PubMed: 21574709]

- Rath JM, Villanti AC, Abrams DB, & Vallone DM (2012). Patterns of tobacco use and dual use in US young adults: the missing link between youth prevention and adult cessation. *Journal of Environmental and Public Health*, 2012.
- Resnicow K, Baranowski T, Ahluwalia JS, & Braithwaite RL . (1999). Cultural sensitivity in public health: Defined and demystified. *Ethnicity and Disease*(9), 10–21. [PubMed: 10355471]
- Rigotti NA, Wechsler H LJE. . (2000). U.S. college students' use of tobacco products: Results of a national survey. *Journal of the American Medical Association*(284), 699–705. [PubMed: 10927777]
- Robinson RG, Orleans CT, James DA, & Sutton CD . (1992). *Pathways to freedom*. Philadelphia Fox Chase Cancer Center.
- Roblyer Martha I Zapata, Joseph G Grzywacz, Richard C Cervantes, and Michael J Merten. 2016 "Stress and Alcohol, Cigarette, and Marijuana Use Among Latino Adolescents in Families with Undocumented Immigrants." *Journal of Child and Family Studies* 25 (2):475–87. [PubMed: 26900317]
- SAS. (2008). *SAS/STAT® 9.2 User's Guide*.
- Soneji S SJ, Tanski S. (2014). Multiple tobacco product use among US adolescents and young adults. *Tobacco Control*, 25(2), 174–180. doi:10.1136/tobaccocontrol-2014-051638 [PubMed: 25361744]
- Spooner C, Hall W, & Lynskey M. (2001). The structural determinants of youth drug use. Retrieved from University of New South Wales:
- Substance Abuse and Mental Health Services Administration (SAMHSA). (2013). *Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings*. Rockville, MD.
- Swisher L, Beckstead J, & Bebeau M (2004). Factor analysis as a tool for survey analysis using a professional role orientation inventory as an example. *Physical Therapy*, 84(9), 784. [PubMed: 15330692]
- Tajfel H, & Turner JC (1986). The social identity theory of intergroup behaviour *In* Worchel S & Austin WG. *Psychology of Intergroup Relations* (pp. 7–24). Chicago, IL: Nelson-Hall.
- Thompson KV, Lightfoot NL, Castillo LG, & Hurst ML (2010). Influence of family perceptions of acting white on acculturative stress in African American college students. *International Journal for the Advancement of Counselling*, 32(2), 144–152.
- Tourangeau R, Edwards B, Johnson TP, Wolter KM, & Bates N. (Eds.). (2014). *Hard-to-Survey Populations*: Cambridge University Press.
- Trinidad DR, Perez-Stable EJ, Emery SL, White MM, Grana RA, & Messer KS (2009). Intermittent and light daily smoking across racial/ethnic groups in the United States. *Nicotine & Tobacco Research*, 11(2), 203–210. [PubMed: 19246433]
- U.S. Department of Health and Human Services. (1998). *Tobacco Use Among US Racial/Ethnic Minority Groups—African Americans, American Indians and Alaska Natives, Asian Americans and Pacific Islanders, and Hispanics: A Report of the Surgeon General*. Atlanta: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- U.S. Department of Health and Human Services. (2014). *The Health Consequences of Smoking—50 Years of Progress. A Report of the Surgeon General* Atlanta: GA: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health .
- Vitaro F, Wanner B, Brendgen M, Gosselin C, & Gendreau PL. (2004). Differential contribution of parents and friends to smoking trajectories during adolescence. *Addictive behaviors*, 29(4), 831–835. [PubMed: 15135568]
- Ward KD, Eissenberg T, Gray JN, Srinivas V, Wilson N, & Maziak W (2007). Characteristics of US waterpipe users: a preliminary report. *Nicotine & Tobacco Research*, 9(12), 1339–1346. [PubMed: 18058352]
- Williams DR NH, Jackson JS. (2003). Racial/ethnic discrimination and health: findings from community studies. *American Journal of Public Health*, 93(2), 200–208. [PubMed: 12554570]
- Wills TA, Knight R, Williams RJ, Pagano I, & Sargent JD. (2015). Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. *Pediatrics*, 135(1), e43–e51. [PubMed: 25511118]

- Wu LT, Swartz MS, Brady KT, Hoyle RH, & Workgroup NA (2015). Perceived cannabis use norms and cannabis use among adolescents in the United States. *Journal of psychiatric research*(64), 79–87.
- Zapata Roblyer MI, Grzywacz JG, Cervantes RC, & Merten MJ (2016). Stress and Alcohol, Cigarette, and Marijuana Use Among Latino Adolescents in Families with Undocumented Immigrants. *J Child Fam Stud*, 25(2), 475–487. doi:10.1007/s10826-015-0249-9 [PubMed: 26900317]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

Item section for the IMI-6 by factor from the full Intragroup Marginalization Inventory

	Accusations of Differentiation	Discrepant Values
IM1. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	x	
IM2. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	x	
IM3. Friends and peers in my ethnic group have the same hopes and dreams as me.		x
IM4. Family members tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	x	
IM5. Family members tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	x	
IM6. My family has the same hopes and dreams as me.		x

Table 2.

Weighted sample characteristics

	Percent (SE of %)	Mean (SE)
Intragroup Marginalization		
IM1. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	42.72 (2.6)	2.24 (25.43)
IM2. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	48.94 (2.6)	2.43 (26.11)
IM3. Friends and peers in my ethnic group have the same hopes and dreams as me.	95.35 (0.8)	3.05 (24.73)
IM4. Family members tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	22.86 (1.7)	1.48 (14.97)
IM5. Family members tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	27.48 (2.0)	1.71 (19.29)
IM6. My family has the same hopes and dreams as me.	84.33 (2.0)	3.42 (27.05)
Demographics		
Race/Ethnicity		
Latino	35.14 (15.0)	
Non-Hispanic Black	14.94 (8.7)	
Non-Hispanic Asian/Pacific Islander	40.33 (16.8)	
Mixed Race	9.60 (3.2)	
Male	48.78 (3.4)	
Mother's Education		
College graduate or more	66.94 (3.4)	
No college graduation	33.06 (3.4)	
Age		22.60 (0.23)
Outcomes		
Cigarette use	11.69 (2.8)	
E-cigarette use	13.46 (3.6)	
Cigar use	7.86 (2.5)	
Blunt/wrap use	10.79 (2.4)	
Hookah use	11.05 (2.0)	
Marijuana use	24.89 (4.4)	

Note: SE = standard error

Table 3.

Factor loadings from exploratory factor analysis with oblique rotation

	Factor 1	Factor 2
1. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	0.775	−0.407
2. Friends and peers in my ethnic group tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	0.826	−0.311
3. Friends and peers in my ethnic group have the same hopes and dreams as me.	−0.001	0.798
4. Family members tell me I am not really a member of my ethnic group because I don't look like my ethnic group.	0.946	0.011
5. Family members tell me I am not really a member of my ethnic group because I don't act like my ethnic group.	0.857	0.051
6. My family has the same hopes and dreams as me.	0.045	0.676

Note: Factor loadings >.50 are boldfaced

Table 4.

Means and standard errors of IMI factor scores by race/ethnicity

	Factor 1		Factor 2	
	<i>M</i>	<i>SE</i>	<i>M</i>	<i>SE</i>
Latino	2.10	0.04	3.81	0.07
Non-Hispanic Black	1.64	0.10	3.99	0.50
Non-Hispanic Asian/Pacific Islander	1.82	0.05	3.64	0.05
Mixed Race	2.57	0.23	3.82	0.22

Note: SE = standard error

Table 5.

Mean differences of IMI factor scores by race/ethnicity

		Factor 1			Factor 2		
		M _{diff}	SE	t	M _{diff}	SE	t
Latino	Non-Hispanic Black	0.46	0.10	4.43**	-0.18	0.48	-0.39
Latino	Non-Hispanic Asian/Pacific Islander	0.28	0.07	3.81**	0.17	0.08	2.06
Latino	Mixed Race	-0.46	0.24	-1.98	-0.01	0.20	-.04
Non-Hispanic Black	Non-Hispanic Asian/Pacific Islander	-0.18	0.11	-1.67	0.35	0.48	0.73
Non-Hispanic Black	Mixed Race	-0.93	0.19	-5.0**	0.18	0.42	0.43
Non-Hispanic Asian/Pacific Islander	Mixed Race	-0.75	0.25	-2.99*	0.24	0.24	-0.74

Notes: SE = standard error

*
 $p < .05$,**
 $p < .01$

Table 6.

Correlations of IMI factor scores and tobacco use variables

	1	2	3	4	5	6	7	8
Factor 1	—	-.06	.08 [*]	.02	-.01	.02	.09 ^{**}	.07 [*]
Factor 2		—	.02	.04	.03	.04	-.04	.02
Cigarette use			—	.37 ^{**}	.25 ^{**}	.32 ^{**}	.20 ^{**}	.27 ^{**}
E-cigarette use				—	.20 ^{**}	.27 ^{**}	.33 ^{**}	.34 ^{**}
Cigar use					—	.33 ^{**}	.14 ^{**}	.37 ^{**}
Blunt/wrap use						—	.17 ^{**}	.65 ^{**}
Hookah use							—	.24 ^{**}
Marijuana use								—

* Notes: $p < .05$,^{**} $p < .01$