Published in final edited form as:

Chemistry. 2018 July 02; 24(37): 9417–9425. doi:10.1002/chem.201801298.

Reversible Ligand-Centered Reduction in Low-Coordinate Iron Formazanate Complexes

Dr. Daniel L. J. Broere^[a], Dr. Brandon Q. Mercado^[a], James T. Lukens^[b], Avery C. Vilbert^[b], Gourab Banerjee^[a], Hannah M. C. Lant^[a], Shin Hee Lee^[a], Dr. Eckhard Bill^[c], Dr. Stephen Sproules^[d], Prof. Dr. Kyle M. Lancaster^[b], and Prof. Dr. Patrick L. Holland^[a]

^[a]Department of Chemistry, Yale University, New Haven, Connecticut 06520, United States, daniel.broere@yale.edu / patrick.holland@yale.edu ^[b]Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca New York 14853. ^[c]Max Planck Institute for Chemical Energy Conversion, Stiftstrasse 34-36, D-45470 Mülheim an der Ruhr, Germany ^[d]WestCHEM, School of Chemistry, University of Glasgow, Glasgow G12 8QQ, United Kingdom

Abstract

Coordination of redox-active ligands to metals is a compelling strategy for making reduced complexes more accessible. In this work, we explore the use of redox-active formazanate ligands in low-coordinate iron chemistry. Reduction of an iron(II) precursor occurs at milder potentials than analogous non-redox-active β -diketiminate complexes, and the reduced three-coordinate formazanate-iron compound is characterized in detail. Structural, spectroscopic and computational analysis show that the formazanate ligand undergoes reversible ligand-centered reduction to form a formazanate radical dianion in the reduced species. The less negative reduction potential of the reduced low-coordinate iron formazanate complex leads to distinctive reactivity with formation of a new N-I bond that is not seen with the β -diketiminate analogue. Thus, the storage of an electron on the supporting ligand changes the redox potential and enhances certain reactivity.

FULL PAPER

Low-coordinate iron formazanate complexes undergo reversible ligand-centered reduction, which is possible at milder potentials than related iron complexes with non-redox-active β -diketiminate ligands. The ligand-centered radical avoids iron(I), yet retains the capability to abstract halide atoms

Correspondence to: Daniel L. J. Broere; Patrick L. Holland.

Conflict of interest

The authors declare no competing financial interest.

Keywords

Redox-active ligand; low-coordinate; iron; ligand-centered-radical; formazanate

Introduction

β -Diketiminato ligands have found extensive use as auxiliary ligands to stabilize unusual metal complexes throughout the periodic table.¹ In particular, highly reduced low-coordinate base metal complexes featuring β -diketiminato ligands are able to activate a number of challenging chemical bonds.² Reductive processes in β -diketiminato complexes are normally metal-centered due to the very negative potentials required for ligand-centered reduction.³ However, the negative potential required for metal-centered reduction (placing electrons in the LUMO at the left of Figure 1) requires the use of harsh reductants such as Na or KC_8 , which are often incompatible with exogenous substrates for catalytic transformations. Moreover, catalytic turnover is often not feasible because the reduced states are too high in energy.

The ability of redox-active ligands to reversibly accept and donate electrons offers an appealing way to generate more accessible reduced states: as shown on the right of Figure 1, the ligand orbital can be lower in energy, moderating the redox potential. Accessible ligand-centered redox processes can expand upon the intrinsic reactivity of metal centers.⁴ Redox-active ligands have enabled multi-electron processes in systems where the transition metal changes oxidation states by less than two electrons, thereby facilitating elementary reactions such as oxidative addition and reductive elimination.⁵ Alternatively, redox-active ligands can enable single-electron transformations without changing metal oxidation state.⁶ A recent computational investigation by Hu and Chen⁷ demonstrated the utility of multiconfigurational calculations for characterizing the redox-active bisiminopyridine ligands, which enable facile reductive elimination in iron-catalyzed [2+2] cycloadditions developed by Chirik and coworkers.⁸ Given the wide use of β -diketiminates, there is motivation for exploring a redox-active ligand that is structurally similar, but can readily accept electrons, enabling one-electron reduction at less negative potentials.

Recent studies have shown that β -diketiminates can undergo ligand-centered oxidation in homoleptic complexes of zinc,⁹ cobalt,¹⁰ and nickel,^{11,12} but ligand-centered reductions on transition metals have not been reported. Here, we use a formazanate as a redox-active ligand that is structurally similar to β -diketiminates, but which can accept electrons. By placing an unoccupied ligand orbital low in energy, they may make the complex easier to reduce (Figure 1, right). Otten has shown that formazanates undergo facile ligand-centered reduction when bound to boron and to zinc.^{13,14} Although spectroscopically observed,¹⁵ no formazanate complex of any redox-active transition metal that shows ligand-centered reduction has yet been isolated. Moreover, a recently reported homoleptic formazanate iron(II) complex adopted a low-spin iron(I) configuration upon one-electron reduction rather than generating a ligand-centered radical,¹⁶ highlighting the need for further study.

Herein, we describe the synthesis, characterization and reactivity of some new low-coordinate iron formazanate complexes and their one-electron reduction products.

Spectroscopic and computational data are used to characterize the frontier orbitals. These studies reveal that the iron-formazanate complexes undergo ligand-centered reduction at relatively mild potentials, and demonstrate an isolable example of a redox-active metal with a formazanate radical dianion. In order to illustrate the new opportunities coming from redox activity at the formazanate ligand, we demonstrate how the formazanate complexes activate chemical bonds like low-coordinate β -diketimate iron complexes, but in contrast are solely mediated by ligand-centered redox changes. In addition, we show that the less electron donating formazanate ligand affects the stability of higher metal oxidation states through an unusual N-I elimination reaction.

Results and Discussion

Synthesis and characterization.

The dropwise addition of a suspension of the known¹⁷ formazan **1** to a stirred solution of $\text{Fe}[\text{N}(\text{SiMe}_3)_2]_2$ in pentane gives the three-coordinate formazanate complex **2** in near quantitative yield (Scheme 1). The ^1H NMR spectrum of complex **2** in C_6D_6 at 298 K shows the expected number of paramagnetically shifted resonances between +82 and –40 ppm, which are tentatively assigned based on their relative integration and distance from the metal center (Figure S1). The presence of a single equivalent of THF in the synthesis of complex **2** results in the exclusive formation of the four-coordinate complex **3** (Scheme 1). Hence, care must be taken to avoid the presence of THF in the solvent and starting materials.¹⁸ We observed that the resonance for the SiMe_3 protons shifts >40 ppm upfield upon the addition of one equiv of THF to complex **2**. Due to fast exchange on the NMR time scale and an equilibrium constant > 5000 M^{-1} , the chemical shift can be used as an indication of the THF content of the sample (Figure S27).

Cooling a solution of **2** or **3** in pentane yielded crystals suitable for single crystal X-ray diffraction. In **2**, two crystallographically independent molecules were found in the asymmetric unit, which share a π – π interaction between the tolyl groups in the formazanate ligands (Figure S35). Each molecule displays a planar, three-coordinate geometry at iron where the sum of the three angles is $359.8 \pm 0.5^\circ$ (Figure 2, left). A notable feature in the solid state structure of **2** is the small dihedral angle between the formazanate and *N*-phenyl groups ($1.6 - 28.3^\circ$), which is very different from analogous three-coordinate Fe complexes containing a $\text{N}(\text{SiMe}_3)$ and bulky β -diketimate ligands ($65 - 90^\circ$).¹⁹ We attribute this difference to the lack of sterically demanding *ortho*-substituents on the *N*-phenyl groups in complex **2**. The Fe–N bond lengths to the formazanates ($1.95 - 1.97 \text{ \AA}$) are shorter than those in analogous three-coordinate Fe complexes containing a $\text{N}(\text{SiMe}_3)$ and bulky β -diketimate ligands ($1.99 - 2.03 \text{ \AA}$).²⁰ The N–N bond lengths ($1.319(5) - 1.327(5) \text{ \AA}$) are slightly longer than commonly observed for formazanate complexes ($1.30 - 1.31 \text{ \AA}$)^{14e,15}, which was also observed for the reported homoleptic iron formazanate complex.¹⁷ A notable difference is the significantly longer Fe–N bonds ($1.955(4) - 1.966(4) \text{ \AA}$) in **2** when compared with the homoleptic bis(formazanate)iron(II) species ($1.817(2) - 1.833(2) \text{ \AA}$).¹⁷ The solid state structure of complex **3** (Figure 2, right) shows that the iron center adopts a distorted trigonal pyramidal geometry ($\tau_4 = 0.70$ and 0.72)²⁰ for both molecules in the asymmetric unit. As expected, coordination of an additional ligand causes all Fe–N bond

lengths in **3** to be elongated (1.979(2) – 2.000(2) Å) compared to **2**. Interestingly, the bulky amide forces the bound THF into a geometry wherein the C atoms lie over the formazanate ligand.

Cyclic voltammetry of complex **3** in THF revealed two reversible one-electron reduction events at $E_{1/2}^{\text{red1}} = -1.65$ V and $E_{1/2}^{\text{red2}} = -2.47$ V vs Fc⁺/Fc (Figure S29). The first reduction occurs at a potential ~0.8 V more positive than the corresponding three coordinate β -diketiminato complexes ($E_{1/2}^{\text{red}} = -2.4$ to -2.6 V vs Fc⁺/Fc).²¹ The nature of this reduction will be discussed below. The addition of CoCp*₂ to a solution of **2** or **3** in hexanes gave the one-electron reduced species **4**, which precipitated as a green solid (Scheme 2). The ¹H NMR spectrum of **4** in THF-*d*₈ at 298 K shows the expected number of paramagnetically shifted resonances between +105 and –25 ppm consistent with a single C_{2v} symmetric species (Figure S9). Intraligand bond lengths in redox-active ligands can often be used to determine the locus of a redox event.^{4–7,15,22} Unfortunately, we were unable to grow crystals of complex **4** that were suitable for X-ray structure determination. Although complex **4** is stable in the solid state for months, it decomposes in THF solution at room temperature within hours to form a mixture of species, concomitant with a color change from dark green to brown. X-ray diffraction of crystals obtained from this mixture revealed a complex wherein the formazanate has undergone N–N bond scission (Figure S37). However, we were unable to isolate sufficient amounts of this unusual product for full characterization, and thus we sought a more stable analogue. Since the C–H bonds in CoCp*₂ and [CoCp*₂]⁺ are susceptible to deprotonation or C–H activation,²³ and because of potential stabilizing effects of alkali cations,²⁴ we used an equivalent of elemental Na to reduce complex **3** in THF, which resulted in a green solution similar in color to **4**. Addition of 2 equiv of 12-crown-4 and cooling the THF mixture gave **5** as a green crystalline solid in 67% yield (Scheme 2). The ¹H NMR (Figure S15) and zero-field Mössbauer spectra (vide infra) of **5** are nearly identical to those for complex **4**. However, unlike **4**, complex **5** is stable in solution for days under inert atmosphere. Single crystals suitable for X-ray structure determination were obtained by cooling a THF solution of **5**. The solid state structure reveals that the Na is sandwiched between two crown ethers, and the resulting cation is separated from the anionic formazanate complex (Figure 3). Although the anion in **5** is isostructural to neutral complex **2**, there are some remarkable differences in the bond lengths and angles. The iron amide bond (Fe1–N14) is longer in complex **5** by *ca.* 0.05 Å, suggestive of a more electron-rich Fe center. The N–N bonds within the formazanate are elongated (1.377(6) and 1.345(6) Å) and the Fe–N (formazanate) bonds (1.951(5) and 1.958(5) Å) are shortened compared to **2** (Figure 2), which is consistent with either formazanate-centered reduction or significant π -backdonation into the formazanate ligand.

Magnetism and EPR spectroscopy.

The temperature dependence of the magnetic susceptibility of solid samples of complexes **2**–**5** was studied in the range 2–290 K using a SQUID magnetometer, and at room temperature in solution using the Evans method. The room temperature magnetic moments of complex **2** and **3** in C₆D₆ solution were $4.8 \pm 0.1 \mu_B$ and $4.7 \pm 0.1 \mu_B$, respectively, in agreement with a high-spin iron(II) configuration for both complexes ($S = 2$, spin-only value $\mu_{s.o.} = 4.9 \mu_B$). In the solid state, three-coordinate **2** and four-coordinate **3** have nearly identical, temperature

independent magnetic moments of $5.1 \mu_B$ over the range 30 – 290 K (Figures S4 and S8), which is near the solution values and consistent with a high-spin iron(II) ($S = 2$) ion in each complex. Below 30 K, the μ_{eff} for **2** decreases under the influence of zero-field splitting. The sign and magnitude of the zero-field parameters were determined from simulations of $\mu_{\text{eff}}(T)$ in combination with isofield magnetization measurements at 1, 4, and 7 T (Figure S4 inset). Global fitting of the magnetic data yielded $D = -3.8 \text{ cm}^{-1}$ and rhombicity $E/D = 0.19$. The temperature profile is different for **3**, where there is a slight rise in μ_{eff} below 30 K, presumably due to intermolecular interactions. The best fit was obtained for $D = 3.1 \text{ cm}^{-1}$ and $E/D = 0.30$, where the sign of D is opposite to that for three-coordinate **2**, though the sign loses meaning so close to full rhombicity. The similarity in the magnitude of these spin-Hamiltonian parameters for **2** and **3** reveal only a very minor change to the electronic structure upon THF coordination.

Room temperature magnetic moments of $3.8 \pm 0.1 \mu_B$ and $3.9 \pm 0.1 \mu_B$ were determined from THF solutions of **4** and **5**, respectively, suggesting $S = 3/2$ ground states for both compounds ($\mu_{\text{s.o.}} = 3.9 \mu_B$). Both **4** and **5** exhibited temperature independent effective magnetic moments of $4.0 \mu_B$ over the range 50 – 290 K (Figures S14 and S18), consistent with these values. Much like **2**, **4** shows a low-temperature downturn in μ_{eff} due to zero-field splitting but the more gentle decrease is modelled with a larger $D = -16.8 \text{ cm}^{-1}$ and large rhombicity $E/D = 0.29$. For **5**, on the other hand, the downturn in μ_{eff} is more abrupt, suggesting intermolecular interactions like those observed in **3**. Nevertheless an excellent fit was obtained from modelling the combined $\mu_{\text{eff}}(T)$ and isofield magnetization plots (including an intermolecular interaction approximation), which gave $D = -10.1 \text{ cm}^{-1}$ and $E/D = 0.26$. These solid state measurements were augmented with an EPR analysis of the odd-electron complexes **4** and **5**. Frozen solution spectra recorded in THF at 10 K highlighted the aforementioned solution instability of **4**. EPR spectra of **4** (Figure S12) show a rhombic $S = 3/2$ signal, and a strong rhombic $S = 1/2$ signal that originates from $(\text{formazanate})_2\text{Fe}^{\text{I}}$,¹⁷ which is one of the decomposition product of **4** (see SI for more details). The $S = 3/2$ signal has peaks at resonant field positions consistent with an $S = 3/2$ species having $g = (2.17, 2.24, 1.87)$, $D = -5.7 \text{ cm}^{-1}$, and $E/D = 0.19$ (Figure S12). The greater solution stability displayed by **5** yielded an EPR spectrum (Figure 4) wholly consisting of an $S = 3/2$ signal with $g_{\text{eff}} = (2.68, 2.03, 2.34)$, $D = 16 \text{ cm}^{-1}$ and $E/D = 0.33$.

Minor deviations in g and E/D may be due to structural differences of solid and solution samples (D was taken from magnetic data because the second Kramers doublet could not be determined due to relaxation issues). The increased quality of the $S = 3/2$ signal in the sample of **5** is consistent with the greater solution stability of this compound. Our assignment of **5** is bolstered by the agreement of g values and zfs parameters between EPR and magnetometry.

Mössbauer spectroscopy.

The zero-field Mössbauer spectra of solid samples of **2** and **3** at 80 K (Figure 5) each showed a quadrupole doublet, with **2** at $\delta = 0.63 \text{ mm s}^{-1}$ and $|E_Q| = 2.48 \text{ mm s}^{-1}$, and **3** at $\delta = 0.71 \text{ mm s}^{-1}$ and $|E_Q| = 1.46 \text{ mm s}^{-1}$. These parameters are within the range previously observed for three-²⁵ and four-coordinate²⁶ high-spin iron(II) diketimate complexes. The

smaller quadrupole splitting and larger isomer shift of **3** in comparison to **2** are consistent with the increase in coordination number.²⁷ The zero-field Mössbauer spectrum of a solid sample of **4** at 80 K shows an asymmetric doublet with $\delta = 0.48 \text{ mm s}^{-1}$ and $|E_Q| = 1.25 \text{ mm s}^{-1}$ (Figure 5, bottom), which shows no significant changes upon warming to 173 K. The zero-field Mössbauer spectrum of a solid sample of **5** at 80 K is nearly identical to that of **4**, with $\delta = 0.49 \text{ mm s}^{-1}$ and $|E_Q| = 1.30 \text{ mm s}^{-1}$ (Figure S16). The asymmetric broadening of the doublets is common for Kramers systems as a result of intermediate rates of spin relaxation.²⁸ Interestingly, the change in isomer shift between **2/3** and **4/5** is comparable, but opposite in direction from the reduction of the homoleptic iron formazanate complex reported by Otten and co-workers,¹⁷ which undergoes metal-centered reduction.

Though it is tempting to use the shift in Mössbauer parameters to assign the metal oxidation state, previous results advise caution. In earlier work we described a di(μ -hydrido)diiron(II) complex²⁹ with a high-spin electronic configuration ($S = 2$) supported by bulky β -diketiminate ligand that gave a mononuclear three-coordinate iron(I) hydride complex with an $S = 3/2$ electronic configuration.³⁰ The change in observed isomer shift in the zero-field Mössbauer spectra of solid samples of the di(μ -hydrido)diiron(II) complex ($\delta = 0.59 \text{ mm s}^{-1}$) and mononuclear three-coordinate iron(I) hydride ($\delta = 0.40 \text{ mm s}^{-1}$) at 80 K is comparable to the decrease in isomer shift between **2/3** and **4/5**. On the other hand, we have also observed similar changes in isomer shift from changes in ligand oxidation state. Namely, oxidation of a four-coordinate high-spin iron(II) complex ($S = 2$) with a dianionic redox-active tetrazene ligand ($\delta = 0.81 \text{ mm s}^{-1}$) gives a complex where a tetrazene radical anion ($S = 1/2$) is strongly antiferromagnetically coupled to a high-spin iron(II) center ($S = 2$) to give an overall $S = 3/2$ electronic configuration ($\delta = 0.69 \text{ mm s}^{-1}$).³¹ Consequently, the observed changes in isomer shift between **2/3** and **4/5** could be attributed to either ligand- or metal-centered reduction.

UV-Vis and X-ray absorption spectroscopy, and computations.

The iron(II) complexes **2** and **3** in hexane solution each show an intense absorption in the visible region at 524 nm ($19 \times 10^3 \text{ cm}^{-1}$), which we assign as the formazanate $\pi-\pi^*$ transition by analogy to intense bands at similar energies uncoordinated formazans.³² The absorption in these compounds is red-shifted by ca. 30 nm ($\sim 1 \times 10^3 \text{ cm}^{-1}$) relative to the protonated formazanate ligand.^{14e} In contrast, the electronic spectrum of **5** recorded at in THF at ambient temperature (Figure 6) exhibits prominent absorption bands at 414 and 696 nm ($\epsilon = 25 \times 10^3$ and $10 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$). Absorption bands with comparable intensity and position have been observed in reduced triaryl-formazanates bound to redox-inert atoms (carbon:³³ $\lambda = 398$ and 721 nm , $\epsilon = 7.5 \times 10^3$ and $3.5 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$; boron:^{14a} $\lambda = 454$ and 716 nm , $\epsilon = 15 \times 10^3$ and $7.5 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$; zinc:^{14a} $\lambda = 462$ and 755 nm , $\epsilon = 20 \times 10^3$ and $10 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$), and are diagnostic of a formazanate radical dianion. Together with the Mössbauer and EPR spectrum and magnetism of **5**, the absorption spectrum supports an electronic configuration where a ligand-centered radical ($S_L = 1/2$) is strongly antiferromagnetically coupled to a high-spin iron(II) center ($S_{\text{Fe}} = 2$) to give an overall $S_{\text{total}} = 3/2$ ground state.

The Fe K-edge X-ray absorption spectra of **2** – **4** are overlaid in Figure 7. They are characterized by electric dipole forbidden but quadrupole-allowed $1s \rightarrow 3d$ pre-edge transitions that appear at the base of the dipole-allowed $1s \rightarrow np$ rising edge that dominates a K-edge spectrum. Since the ligand field typically increases with increasing oxidation state for a given ligand set, the pre-edge energy often provides a useful marker for oxidation state. The centers of mass for pre-edge peaks in spectra obtained for **2** – **4** are identical within experimental error at 7112.2 ± 0.1 eV. As a ~ 1 eV increase in energy is expected per unit of oxidation, the similar edge positions indicate that the physical Fe oxidation state is the same for the Fe ion in neutral **2** and **3**, and the monoanion in **4**. Moreover, it is noteworthy that the pre-edge peak is of the same intensity for each complex. Electric quadrupole allowed $1s \rightarrow 3d$ pre-edge transitions gain intensity through $4p$ mixing into the $3d$ orbitals and therefore are sensitive to the symmetry of the complex.³⁴ Here, the uniform peak height is consistent with the dominant trigonal symmetry in this series where the inclusion of a THF ligand in **3** has little effect on its electronic structure (vide supra). In neutral **2** and **3**, there is a shoulder to higher energy at ~ 7113.2 eV. In contrast, the shoulder in **4** lies to lower energy at 7111.5 eV, and is tentatively assigned as transition to an orbital with significant ligand character.³⁵

Although the rising K-edge is frequently used as an additional metric of physical oxidation state in first-row metals, the rising-edge energies are also influenced by coordination geometry and ligand identity.³⁶ In the present case, similar edge positions are displayed by three-coordinate **2** and **4** of 7117.0 and 7117.4 eV, respectively, while the edge is shifted by 0.7 eV to higher energy at 7118.1 eV for four-coordinate **3**. Thus, these do not change the conclusion from the pre-edge features, that all the complexes have the same oxidation state at iron.

Density functional theory calculations were carried out to facilitate spectroscopic and electronic structure interpretations. In particular, time-dependent DFT (TDDFT) calculations are known to reproduce metal K-edge XAS with remarkable fidelity after accounting for inaccurate modeling of the $1s$ core potential with either a scalar or linear energy shift.³⁷ Time-dependent (TD) DFT calculations (B3LYP hybrid density functional and CP(PPP) basis set on Fe with ZORA-def2-TZVP(-f) on all remaining atoms) were performed on the crystallographic coordinates of each complex. For **2** and **3**, these reproduce the pre-edge region with good fidelity (Fig 7 inset), but the TD-DFT calculation for **4** produces a pre-edge with an energy maximum that is underestimated by 1 eV. Inspection of the frontier orbitals calculated for **4** using this level of theory revealed a straightforward unrestricted Kohn-Sham solution consistent with ($S = 3/2$) iron(I). Given the consistency of the spectroscopic data described above with an iron(II) configuration, we evaluated a broken-symmetry (BS) option where the four electrons from high-spin iron(II) are allowed to participate in a magnetic interaction with an $S = 1/2$ formazanate radical anion (BS 4,1). While this approach reproduces the Fe K pre-edge with remarkable fidelity, the BS calculation converges to a *ferromagnetic* $S = 5/2$ solution, which conflicts with the magnetic measurements. Thus, the single-reference DFT methods fail to acceptably reproduce physical/spectroscopic properties of **4**.

We hypothesized that the ground states of the complex ions in **4** and **5** are multiconfigurational, explaining the poor performance of DFT. Thus, we carried out a

multireference configuration interaction (MRCI) calculation using the spectroscopy oriented configuration interaction (SORCI) method.³⁸ We employed an active space of 11 electrons and 9 orbitals [CAS(11,9)] chosen to include five Fe 3d orbitals and four electrons from the formazanate π system. Five quartet, doublet, and sextet states were calculated. The chosen active space gave 90% reference weights for all states. The resulting state energies are given in Table 1. SORCI appropriately predicts a ground-state quartet, with the lowest spin-forbidden excited state—a sextet—occurring at ca. 4300 cm⁻¹ to higher energy. The UV-vis absorption spectrum calculated by SORCI reproduces the intense absorption at 14,000 cm⁻¹.

Inspection of the quartet ground state exposes substantial multiconfigurational character. Five configurations have greater than 5% participation in the ground state, with the two leading configurations contributing ca. 25% each. Details of these two principal leading configurations are shown in Figure 8. Evaluating the composition of each configuration reveals that the ground state is best described as a nearly equal mix of configurations falling into two electronic structure categories. In one category, high-spin iron(II) participates in antiferromagnetic coupling with the unpaired electron residing in an a₂-symmetry formazanate π^* MO. The second category represents high-spin iron(I) without ligand participation.

In further agreement with our assignment, the homoleptic formazanate iron complex reported by Otten and co-workers,¹⁷ which is proposed to undergo metal-centered reduction, lacks the characteristic absorption bands for ligand-centered reduction that **5** displays at 414 and 696 nm. Overall, the electronic absorption spectra show that the one-electron reduction of the formazanate complexes involves substantial ligand participation.

Reactivity of the reduced complexes.

No differences are found in the ¹H NMR spectra of complexes **4** and **5**, except for resonances attributable to the counteranion. Yet, complex **4** decomposes within hours in solution whereas no signs of decomposition were observed for solutions of **5** over the course of several days (vide supra). To investigate whether this was from reaction between the anion and CoCp*₂ or from sodium stabilization that slows decomposition, NaBAR^F₄ was added to a solution of complex **4** and [CoCp*₂]PF₆ was added to a solution of **5**. In both cases decomposition was observed to the same mixture of unidentified species, showing that the decomposition of the one-electron reduced complex is brought about by the presence of the CoCp*₂ cation. Though we have not been able to determine the fate of the CoCp*₂ cation, it is possible that decomposition occurs through a pathway involving C-H activation through deprotonation of one of the Cp* ligands.²⁴

Reduced low-coordinate iron complexes often abstract halide atoms from sp² and sp³ carbon atoms. Considering the significantly less negative reduction potential of **5**, we wanted to evaluate its competence in the activation of C-X (X = halogen bonds). Addition of alkyl and aryl iodides to solutions of **5** results in rapid formation of iron(II) iodide-amide complex **6**, which can also be prepared by reaction of **5** with 0.5 equiv of I₂ or by addition of 1 equiv NaI to solutions of **2** or **3** (Scheme 3). Addition of trityl chloride to a solution of **5** showed formation of triphenylmethyl radical (observed as Gomberg's dimer)³⁹ and the chloride analogue of **6**.

Accessing the ferric oxidation state was possible by a reaction of **2** or **3** with 0.5 equiv of I₂ giving complex **7**, which has a high-spin iron(III) ($S = 5/2$) configuration (see Supporting Information). Accordingly, we expected that treatment of **5** with one equiv of I₂, or **6** with 0.5 equiv of I₂ in THF, would also give **7**. However, these reactions result in the formation of complex **8** that still has a high-spin iron(II) ($S = 2$) center in 82% isolated yield. The only additional reaction product in the oxidations of **5** and **6** is NaI. To test our hypothesis that complex **7** is formed in these reactions but undergoes a decomposition pathway involving NaI we modified the reaction conditions to enable monitoring by ¹H NMR spectroscopy. Indeed, the addition of NaI to a THF-*d*₈ solution of complex **7** gave **8** in 65% spectroscopic yield together with **6**. Similarly, the addition of I₂ to a THF-*d*₈ solution of complex **5** gave **8** in 70 % spectroscopic yield together with **6**. In addition, formation of varying amounts of I-N(SiMe₃)₂ was observed in both reactions, and its presence was confirmed by spiking the mixture with an independently synthesized sample of I-N(SiMe₃)₂.⁴⁰

In the reaction of **7** with NaI an intermediate with very broad resonances is observed, and the apparent concentration is more pronounced at higher NaI concentration. We propose that this intermediate is a five-coordinate Fe(III) species that is formed from **7** and NaI. This five-coordinate species either reductively eliminates I-N(SiMe₃)₂ in analogy to the reaction known for a Ni(IV) complex,⁴¹ or loses •N(SiMe₃)₂⁴² as observed for lanthanide complexes.⁴³ In the latter scenario the formation of I-N(SiMe₃)₂ can be explained by abstraction of an I atom from Fe by •N(SiMe₃)₂.

In contrast, the β-diketiminato analogue of complex **7** (see Supplementary Information) does not react with stoichiometric or excess NaI. This difference suggests that the formazanate ligand is beneficial for N-I elimination. Both of the potential pathways, reductive elimination and Fe-N homolysis, are thermodynamically more favorable when the lower oxidation state is favored, indicating that the substitution of diketiminato for formazanate successfully modulates the electron density of the iron center in a way that enables new reactivity.

Conclusions

This manuscript has described the synthesis of low-coordinate iron formazanate complexes that can reversibly undergo reduction at a less negative potential compared to analogous iron β-diketiminato species. Distinguishing the high-spin iron(I) description from the antiferromagnetically coupled iron(II)-ligand radical description is challenging, and the distinctive absorption spectra support the latter model. Moreover, multiconfigurational calculations support the presence of a substantial character of iron(II) with a formazanate radical dianion. Thus, in contrast with the earlier reported bis(formazanate) complexes that have predominant iron(I) character,¹⁷ these highly electron-rich amido complexes have electron density on the supporting ligand. To our knowledge, **5** is the first well-characterized complex of a redox-active metal with a formazanate radical dianion. Despite the less negative reduction potential, **5** activates carbon halogen bonds mimicking the reactivity of low valent β-diketiminato iron complexes. In an important difference, the less-electron donating formazanate does not stabilize the higher ferric oxidation state, and therefore a different reaction pathway is followed, leading to N-I elimination.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by The Netherlands Organization for Scientific Research (Rubicon Postdoctoral Fellowship 680-50-1517 to D.L.J.B.) and the National Institutes of Health (Grant GM-065313 to P.L.H.). K.M. L. thanks the A. P. Sloan Foundation and the National Science Foundation (CHE-1454455) for funding. P.L.H. thanks the Alexander von Humboldt Foundation for support. We thank Prof. Gary Brudvig for access to an EPR spectrometer.

References

1. a) Bourget-Merle L; Lappert MF; Severn JR *Chem. Rev* 2002, 102, 3031–3066; [PubMed: 12222981] b) Kundu S; Stieber SCE; Ferrier MG; Kozimor SA; Bertke JA; Warren TH *Angew. Chem. Int. Ed*, 2016, 55, 10321–10325.
2. Selected examples MacLeod KC; Lewis RA; DeRosha DE; Mercado BQ; Holland PL *Angew. Chem. Int. Ed* 2017, 56, 1069–1072; Pfirrmann S; Limberg C; Herwig C; Stösser R; Ziemer B *Angew. Chem. Int. Ed* 2009, 48, 3357–3361; Dugan TR; Bill E; MacLeod KC; Christian GJ; Cowley RE; Brennessel WW; Ye S; Neese F; Holland PL *J. Am. Chem. Soc* 2012, 134, 20352–20364; [PubMed: 23181620] Horn B; Limberg C; Herwig C; Braun B *Chem. Commun* 2013, 49, 10923; Dai X; Kapoor P; Warren TH *J. Am. Chem. Soc* 2004, 126, 4798–4799. [PubMed: 15080682]
3. For exceptions see: Eisenstein O; Hitchcock PB; Khvostov AV; Lappert MF; Maron L; Perrin L; Protchenko AV *J. Am. Chem. Soc*, 2003, 125, 10790–10791. [PubMed: 12952450] Avent AG; Hitchcock PB; Khvostov AV; Lappert MF; Protchenko AV *Dalton Trans*, 2004, 33, 2272–2280.
4. a) Broere DLJ; Plessius R; van der Vlugt JI *Chem. Soc. Rev* 2015 44, 6886–6915; [PubMed: 26148803] b) Luca OR; Crabtree RH *Chem. Soc. Rev* 2013, 42, 1440–1459; [PubMed: 22975722] c) Praneeth VKK; Ringenberg MR; Ward TR *Angew. Chem. Int. Ed* 2012, 51, 10228–10234; d) van der Vlugt JI *Eur. J. Inorg. Chem.* 2012, 363–375; e) Lyaskovskyy V; de Bruin B; *ACS Catal.* 2012, 2, 270–279; f) Kaim W *Inorg. Chem.* 2011, 50, 9752–9765; [PubMed: 21744793] g) Chirik PJ; Wieghardt K *Science* 2010, 327, 794–795. [PubMed: 20150476]
5. a) Johnson SA; Higgins RF; Abu-Omar MM; Shores MP; Bart SC *Organometallics* 2017, 36, 3491–3497; b) Wong JL; Hernández Sánchez RH; Glancy Logan J; Zarkesh RA; Ziller JW; Heyduk AF *Chem. Sci.* 2013, 4, 1906–1910; c) Kraft SJ; Fanwick PE; Bart SC *J. Am. Chem. Soc.* 2012, 134, 6160–6168; [PubMed: 22468980] d) Haneline MR; Heyduk AF *J. Am. Chem. Soc.* 2006, 128, 8410–8411. [PubMed: 16802801]
6. a) Jacquet J, Cheaib K, Ren Y, Vezin H, Orio M, Blanchard S, Fensterbank L, M. Desage-El Murr *Chem. Eur. J.* 2017, 23, 15030–15034; b) Fujita D, Sugimoto H, Shiota Y, Morimoto Y, Yoshizawa K, Itoh S, *Chem. Commun.* 2017, 53, 4849–4852; c) Jacquet J, Blanchard S, Derat E, Desage-El Murr M, Fensterbank L, *Chem. Sci.* 2016, 7, 2030–2036; [PubMed: 29899928] d) Broere DLJ, Modder DK, Blokker E, Siegler MA, van der Vlugt JI, *Angew. Chem. Int. Ed.* 2016, 55, 2406–2410; e) Broere DLJ, de Bruin B, Reek JNH, Lutz M, Dechert S, van der Vlugt JI, *J. Am. Chem. Soc.* 2014, 136, 11574–11577; [PubMed: 24927362]
7. Hu L; Chen HJ *Am. Chem. Soc.* 2017, 139, 15564–15567.
8. a) Hoyt JM; Schmidt VA; Tondreau AM; Chirik PJ *Science* 2015, 349, 960–963; [PubMed: 26315433] b) Bouwkamp MW; Bowman AC; Lobkovsky E; Chirik PJ *J. Am. Chem. Soc.* 2006, 128, 13340–13341; [PubMed: 17031930] c) Hoyt JM; Sylvester KT; Semproni SP; Chirik PJ *J. Am. Chem. Soc.* 2013, 135, 4862–4877; [PubMed: 23448301] d) Russell SK; Lobkovsky E; Chirik PJ *J. Am. Chem. Soc.* 2011, 133, 8858–8861. [PubMed: 21598972]
9. Khusniyarov MM; Bill E; Weyhermüller T; Bothe E; Wieghardt K *Angew. Chem., Int. Ed.* 2011, 50, 1652–1655.
10. Marshak MP; Chambers MB; Nocera DG *Inorg. Chem.* 2012, 51, 11190–11197. [PubMed: 23013488]

11. Takaichi J; Morimoto Y; Ohkubo K; Shimokawa C; Hojo T; Mori S; Asahara H; Sugimoto H; Fujieda N; Nishiwaki N; Fukuzumi S; Itoh S Inorg. Chem, 2014, 53, 6159–6169. [PubMed: 24884152]
12. For a review on the non-innocence of β -diketiminatate ligands see: Camp C; Arnold J; Dalton Trans 2016, 45, 14462–14498. [PubMed: 27353604]
13. a)Chang M-C; Otten E Chem. Commun. 2014, 50, 7431–7433;b)Chang M-C; Chantzis A; Jacquemin D; Otten E Dalton Trans. 2016, 45, 9477–9484; [PubMed: 27188989] c)Barbon SM; Price JT; Reinkeluers PA; Gilroy JB Inorg. Chem. 2014, 53, 10585–10593; [PubMed: 25226240] d)Mondol R; Snoeken DA; Chang M-C; Otten E Chem. Commun. 2017, 53, 513–516;e)Gilroy JB; Ferguson MJ; McDonald R; Patrick BO; Hicks RG Chem. Commun 2007, 126–128.
14. a)Chang M-C; Dann T; Day DP; Lutz M; Wildgoose GG; Otten E Angew. Chem. Int. Ed. 2014, 53, 4118–4122;b)Chang M-C Roewen P; Travieso-Puente R; Lutz M; Otten E; Inorg. Chem. 2015, 54, 379–388. [PubMed: 25493709]
15. a)Mandal A; Schwederski B; Fiedler J; Kaim W; Lahiri GK Inorg. Chem. 2015, 54, 8126–8135; [PubMed: 26244848] b)Kabir E; Wu C-H; Wu JI; Teets TS Inorg. Chem. 2016, 55, 956–963; [PubMed: 26702999] c)Gilroy JB; Patrick BO; McDonald R; Hicks RG Inorg. Chem. 2008, 47, 1287–1294; [PubMed: 18211010] d)Hong S; Hill LMR; Gupta AK; Naab BD; Gilroy JB; Hicks RG; Cramer CJ; Tolman WB Inorg. Chem. 2009, 48, 4514–4523. [PubMed: 19425614]
16. Travieso-Puente R; Broekman JOP; Chang M-C; Demeshko S; Meyer F; Otten E; J. Am. Chem. Soc. 2016, 138, 5503–5506. [PubMed: 27074728]
17. Gilroy JB; Ferguson MJ; McDonald R; Patrick BO Hicks RG; Chem. Commun. 2007,126–128.
18. Broere DLJ; Ori I; Brosnahan A; Holland PLH Inorg. Chem. 2017, 56, 3140–3143. [PubMed: 28248496]
19. Panda A; Stender M; Wright RJ; Olmstead MM; Klavins P; Power PP Inorg. Chem. 2002, 41, 3909–3916. [PubMed: 12132915]
20. Yang L; Powell DR; Houser RP Dalton Trans. 2007, 955–964. [PubMed: 17308676]
21. Chiang KP; Barrett PM; Ding F; Smith JM; Kingsley S; Brennessel WW; Clark MM; Lachicotte RJ; Holland PL Inorg. Chem. 2009, 48, 5106–5116. [PubMed: 19438179]
22. a)Brown SN Inorg. Chem. 2012, 51, 1251–1260; [PubMed: 22260321] b)Broere DLJ; Plessius R; Tory J; Demeshko S; de Bruin B; Siegler MA; Hartl F; van der Vlugt JI Chem. Eur. J. 2016, 22, 13965–13975; [PubMed: 27531163] c)Broere DLJ; van Leest NP; Siegler MA; van der Vlugt JI Inorg. Chem. 2016, 55, 8603–8611; [PubMed: 27525360] d)Sproules S; Wieghardt K. Coord. Chem.Rev. 2010, 254, 1358–1382.
23. MacLeod KC; McWilliams SF; Mercado BQ; Holland PL Chem. Sci. 2016, 7, 5736–5746; [PubMed: 28066537] b)Ohki Y; Murata A; Imada M; Tatsumi K Inorg. Chem. 2009, 48, 4271–4273. [PubMed: 19341307]
24. a)MacLeod KC; Menges FS; McWilliams SF; Craig SM; Mercado BQ; Johnson MA; Holland PL J. Am. Chem. Soc 2016, 138, 11185–11191; [PubMed: 27571271] b)McWilliams SF; Rodgers KR; Lukat-Rodgers G; Mercado BQ; Grubel K; Holland PL Inorg. Chem 2016, 55, 2960–2968; [PubMed: 26925968] c)Grubel K; Brennessel WW; Mercado BQ; Holland PL J. Am. Chem. Soc. 2014, 136, 16807–16816; [PubMed: 25412468] d)Connor GP; Holland PL Catalysis Today 2017, 286, 21–40. [PubMed: 28344387]
25. McWilliams SF; Brennan-Wydra E; MacLeod KC Holland PL ACS Omega 2017, 2, 2594–2606. [PubMed: 28691111]
26. a)Arnet NA; Dugan TR; Menges FS; Mercado BQ; Brennessel WW; Bill E; Johnson MA; Holland PL J. Am. Chem. Soc. 2015, 137, 13220–13223; [PubMed: 26457740] b)Lewis RA; MacLeod KC; Mercado BQ; Holland PL Chem. Commun. 2014, 50, 11114–11117.
27. a)Kleinlein C; Bendelsmith AJ; Zheng S-L; Betley TA Angew.Chem. Int.Ed 2017, 56, 12197–12201;b)Dunsford JJ; Evans DJ; Pugh T; Sha SN; Chilton NF; Ingleson MJ Organometallics 2016, 35, 1098–1106;c)Evans DJ; Hughes DL; Silver J Inorg. Chem. 1997, 36, 747–748.
28. Gülich P; Bill E; Trautwein AX Mössbauer spectroscopy and transition metal chemistry, Springer, Heidelberg, 2011.

29. Initial synthesis: Smith JM; Lachiocotte RJ; Holland PL J. Am. Chem. Soc. 2003i, 125, 15752–15753; [PubMed: 14677959] Mössbauer spectrum and alternative synthesis: Dugan TR; Holland PL J. Organomet. Chem. 2009, 694, 2825–2830.
30. Chiang KP; Scarborough CC; Horitani M; Lees NS; Ding K; Dugan TR; Brennessel WW; Bill E; Hoffman BM; Holland PL Angew. Chem. Int. Ed. 2012, 51, 3658–3662.
31. Cowley RE; Bill E; Neese F; Brennessel WW; Holland PL Inorg. Chem. 2009, 48, 4828–4836. [PubMed: 19397284]
32. a)Tezcan H; Can S; Tezcan R Dyes Pigm. 2002, 52, 121;b)Ibrahim YA; Abbas AA; Elwaby AHMJ; Heterocycl. Chem. 2004, 41, 135.
33. Gilroy JB; McKinnon SDJ; Koivisto BD; Hicks RG Org. Lett. 2007, 9, 4837–4840. [PubMed: 17927192]
34. Shulman RG; Yafet Y; Eisenberger P; Blumberg WE Proc. Natl. Acad. Sci. U.S.A. 1976, 73, 1384–1388. [PubMed: 5720]
35. Banerjee P; Sproules S; Weyhermüller T; DeBeer George S; Wieghardt K Inorg. Chem, 2009, 48, 5829–5847. [PubMed: 20507101]
36. a)MacMillan SN; Lancaster KM ACS Catal. 2017, 7, 1776–1791;b)Kau LS; Spira-Solomon DJ; Penner-Hahn JE; Hodgson KO; Solomon EI J. Am. Chem. Soc. 1987, 109, 6433–6442.
37. DeBeer George S; Petrenko T; Neese FJ Phys. Chem. A 2008, 112, 12936–12943.
38. Neese FJ Chem. Phys. 2003, 119, 9428.
39. Lankamp H; Nauta WT MacLean C Tet. Lett. 1968, 9, 249–254.
40. Bailey RE; West RJ Organomet. Chem. 1965, 4, 430–439.
41. Laarhoven JJ; Mulder PJ Phys. Chem. B 1997, 101, 73–77.
42. Cook MD; Roberts BP; Singh KJ Chem. Soc, Perkin Trans 2, 1983, 635–643.
43. a)Lewis AJ; Carroll PJ; Schelter EJ J. Am. Chem. Soc 2013, 135, 511–518. [PubMed: 23215157] (c)Mullane KC; Lewis AJ; Yin H; Carroll PJ; Schelter EJ Inorg. Chem 2014, 53, 9129–9139. [PubMed: 25111809]

Figure 1.

Conceptual representation of how the redox-active formazanate ligand can provide access to a more accessible reduced state.

Figure 2.

Displacement ellipsoid plots (50% probability) of complex 2 and 3. Hydrogen atoms have been omitted for clarity. Selected bond lengths (Å) and angles (°) for 2: Fe2–N18 1.903(3); Fe2–N15 1.964(4); Fe2–N45 1.966(4); N18–Fe2–N15 132.5(2); N18–Fe2–N45 135.2(2); N15–Fe2–N45 91.7(1). For 3: Fe2–N18 1.937(2); Fe2–N15 1.979(2); Fe2–N45 1.984(2); Fe2–O20 2.145(1); N18–Fe2–N15 130.40(7); N18–Fe2–N45 127.99(6); N15–Fe2–N45 91.59(6).

Figure 3.

Displacement ellipsoid plot (50% probability) of complex **5**. Hydrogen atoms and a THF molecule have been omitted. Selected bond lengths (Å) and angles (°): Fe1–N11 1.935(5); Fe1–N14 1.951(5); Fe1–N41 1.958(5); N11–N21 1.377(6); N31–N41 1.345(6); N11–Fe1–N14 134.9(2); N11–Fe1–N41 96.5(2); N14–Fe1–N41 128.6(2).

Figure 4.

X-band EPR spectrum of **5** in THF at 10 K. Experimental data are represented by the black line; simulation is depicted by the red trace. Asterisk denotes trace impurity contributing <0.5% to the total signal. Experimental conditions: frequency 9.3701 GHz; power 0.2 mW; modulation 2.0 mT.

Figure 5. Zero-field Mössbauer spectrum of a solid samples of complex **2** (top), **3** (middle), **4** (bottom) at 80 K. The black circles are experimental data with the fit represented by the black line. The thin grey trace is the residual of the fit.

Figure 6. Experimental (solid) and calculated (dotted, SORCI) absorption spectra for complex **3** (pink) and **5** (green) in THF solution.

Figure 7. Overlay of the normalized Fe K-edge XAS spectra of **2** (red), **3** (blue), and **4** (green). Inset shows an expansion of the pre-edge region with a comparison of the experimental spectra above the corresponding calculated spectra obtained from B3LYP TD-DFT calculations. Calculated intensity in arbitrary units.

Figure 8.

Calculated SORCI averaged atomic natural orbitals (AANOs) comprising the CAS(11,9) reference as well as the two leading configurations comprising the ground-state quartet of the complex ion in **4**. AANOs comprising principally Fe 3d character (the Fe ligand field) are boxed, and the antiferromagnetic (AF) interaction between Fe 3d_{xy} and the a₂ formazanate (Fz) π* is indicated with a red bracket. Orbitals are plotted at an isovalue of 0.03 au.

Scheme 1.
Synthesis of complexes 2 and 3.

Scheme 2.
Synthesis of complexes **4** and **5**.

Table 1.SORCI-calculated [CAS(11,9)] states for the Fe complex anion in **4**.

State	Spin multiplicity	Energy (cm ⁻¹)
0	4	0
1	4	823
2	6	3953
3	4	4297.7
4	6	4402.3
5	4	5689.2
6	6	5792.2
7	6	8840
8	2	12537.3
9	2	14095
10	4	14254
11	2	17470
12	2	18342.5
13	6	19032.2
14	2	28867