Efficacy and Safety of High-Dose Rifampin in Pulmonary Tuberculosis

A Randomized Controlled Trial

Gustavo E. Velásquez^{1,2}, Meredith B. Brooks², Julia M. Coit², Henry Pertinez^{3,4}, Dante Vargas Vásquez⁵, Epifanio Sánchez Garavito⁶, Roger I. Calderón⁷, Judith Jiménez⁷, Karen Tintaya⁷, Charles A. Peloquin⁸, Elna Osso², Dylan B. Tierney⁹, Kwonjune J. Seung^{9,10}, Leonid Lecca^{2,7}, Geraint R. Davies^{3,4*}, and Carole D. Mitnick^{2,9,10*}

¹Division of Infectious Diseases and ⁹Division of Global Health Equity, Brigham and Women's Hospital, Boston, Massachusetts; ²Department of Global Health and Social Medicine, Harvard Medical School, Boston, Massachusetts; ³Institute of Infection and Global Health and ⁴Institute of Translational Medicine, University of Liverpool, Liverpool, United Kingdom; ⁵Hospital Nacional Hipólito Unanue, Lima, Peru; ⁶Hospital Nacional Sergio Bernales, Lima, Peru; ⁷Partners in Health/Socios en Salud Sucursal Peru, Lima, Peru; ⁸College of Pharmacy and Emerging Pathogens Institute, University of Florida, Gainesville, Florida; and ¹⁰Partners in Health, Boston, Massachusetts

ORCID ID: 0000-0003-1438-0692 (G.E.V.).

Abstract

Rationale: We examined whether increased rifampin doses could shorten standard therapy for tuberculosis without increased toxicity.

Objectives: To assess the differences across three daily oral doses of rifampin in change in elimination rate of *Mycobacterium tuberculosis* in sputum and frequency of rifampin-related adverse events.

Methods: We conducted a blinded, randomized, controlled phase 2 clinical trial of 180 adults with new smear-positive pulmonary tuberculosis, susceptible to isoniazid and rifampin. We randomized 1:1:1 to rifampin at 10, 15, and 20 mg/kg/d during the intensive phase. We report the primary efficacy and safety endpoints: change in elimination rate of *M. tuberculosis* log₁₀ colony-forming units and frequency of grade 2 or higher rifampin-related adverse events. We report efficacy by treatment arm and by primary (area under the plasma concentration–time curve [AUC]/minimum inhibitory concentration [MIC]) and secondary (AUC) pharmacokinetic exposure.

Measurements and Main Results: Each 5-mg/kg/d increase in rifampin dose resulted in differences of -0.011 (95% confidence interval, -0.025 to $+0.002$; $P = 0.230$) and -0.022 (95% confidence interval, -0.046 to -0.002 ; $P = 0.022$) log₁₀ cfu/ml/d in the modified intention-to-treat and per-protocol analyses, respectively. The elimination rate in the per-protocol population increased significantly with rifampin AUC_{0–6} ($P = 0.011$) but not with AUC_{0–6}/MIC_{99.9} ($P = 0.053$). Grade 2 or higher rifampin-related adverse events occurred with similar frequency across the three treatment arms: 26, 31, and 23 participants (43.3%, 51.7%, and 38.3%, respectively) had at least one event ($P = 0.7092$) up to 4 weeks after the intensive phase. Treatment failed or disease recurred in 11 participants (6.1%).

Conclusions: Our findings of more rapid sputum sterilization and similar toxicity with higher rifampin doses support investigation of increased rifampin doses to shorten tuberculosis treatment.

Clinical trial registered with www.clinicaltrials.gov (NCT 01408914).

Keywords: tuberculosis; rifampin; randomized controlled trial; treatment efficacy; adverse drug event

(Received in original form December 15, 2017; accepted in final form May 30, 2018)

*Co–senior authors.

Supported by NIH/National Institute of Allergy and Infectious Diseases grants U01 AI091429 (C.D.M. and G.R.D.) and L30 AI120170 and UM1 AI068636 (G.E.V.). G.E.V. received support from the Ronda Stryker and William Johnston Fellowship in Global Health and Social Medicine and the Dr. Lynne Reid/Drs. Eleanor and Miles Shore Fellowship at Harvard Medical School, the Burke Global Health Fellowship at the Harvard Global Health Institute, and the AIDS Clinical Trials Group Minority HIV Investigator Mentoring Program. The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the NIH or the institutions with which the authors are affiliated. The funding source had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the report for publication.

Author Contributions: G.R.D. and C.D.M. designed the study. G.E.V., M.B.B., G.R.D., and C.D.M. interpreted the data and wrote the first draft of the manuscript. M.B.B., J.M.C., D.V.V., E.S.G., R.I.C., J.J., K.T., C.A.P., E.O., D.B.T., K.J.S., L.L., G.R.D., and C.D.M. implemented the study. D.V.V., E.S.G., R.I.C., J.J., K.T., and C.A.P. collected the data. G.E.V., M.B.B., H.P., and G.R.D. performed the analysis. J.M.C. assisted with data management. All authors revised the manuscript critically for important intellectual content and gave final approval of the version to be published.

Correspondence and requests for reprints should be addressed to Gustavo E. Velásquez, M.D., M.P.H., Department of Global Health and Social Medicine, Harvard Medical School, 641 Huntington Avenue, Boston, MA 02115. E-mail: gvelasquez@bwh.harvard.edu.

This article has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org.

Am J Respir Crit Care Med Vol 198, Iss 5, pp 657–666, Sep 1, 2018

Copyright © 2018 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.201712-2524OC on June 29, 2018

Internet address: www.atsjournals.org

At a Glance Commentary

Scientific Knowledge on the

Subject: The standard of care for patients with new pulmonary tuberculosis is a 6-month, four-drug regimen that includes rifampin for the full course of therapy. Attempts to further shorten tuberculosis therapy in the 1970s with higher, intermittent doses of rifampin were unsuccessful because of an apparent increase in toxicity. Renewed interest in tuberculosis treatment shortening developed earlier this century, with trials of fluoroquinolones, rifapentine, and a shortened regimen in patients at lower risk for poor outcomes. We sought to systematically examine the concept that increased daily doses of rifampin could shorten standard therapy for tuberculosis and improve treatment outcomes without increased toxicity.

What This Study Adds to the

Field: This blinded, randomized, controlled phase 2 clinical trial assessed differences across three daily oral doses of rifampin (10, 15, and 20 mg/kg) in change in elimination rate of *Mycobacterium tuberculosis* in sputum and frequency of rifampin-related adverse events. We found that doses of 15 and 20 mg/kg/d resulted in more rapid change in counts of *M. tuberculosis* and a similar frequency of rifampin-related adverse events. This is the first controlled study to show both dose and exposure response of rifampin on sputum sterilization. Our findings support the continued investigation of higher doses of rifampin, beyond 20 mg/kg, for potential treatment shortening.

Tuberculosis is the leading infectious cause of death worldwide, killing 1.7 million in 2016 (1). Standard treatment for the annual estimated 10.4 million patients with new pulmonary tuberculosis is a 6-month, four-drug regimen including rifampin throughout (2).

Combining rifampin and pyrazinamide in the early 1970s shortened tuberculosis treatment from 18 to 6 months (3). Further attempts to shorten tuberculosis therapy in

the 1970s were guided by perceived cost constraints: any increase in the dose was matched by decreased frequency of administration. The resulting higher, intermittent doses of rifampin led to increased toxicity (4–6). Ultimately, the 600-mg daily dose of rifampin was selected as the standard (7). Interest in shortening tuberculosis treatment has been renewed over the last 20 years, with preclinical and clinical studies of fluoroquinolones and one trial of a further-shortened regimen in patients with noncavitary disease and rapid bacteriological response (8–14). Before the initiation of the present study, a phase 2 trial of treatment shortening with another rifamycin, rifapentine, was published; further work on treatment shortening with rifapentine was subsequently pursued (15–17).

Daily dose optimization of rifampin remained a priority. Unlike moxifloxacin and rifapentine, rifampin is an established first-line drug, available throughout the world, from multiple suppliers, now for pennies a capsule. It is also available in combination formulations. Of the four current first-line agents, rifampin has the most potent sterilizing activity and has been established as key to the continuation phase of treatment (18, 19). At the 600-mg daily dose, rifampin is well tolerated (20–23). However, concern about the toxicity of rifampin at higher doses persists pursuant to trials of intermittent dosing (6, 20, 24).

Although multiple controlled trials began to evaluate higher rifampin doses in African populations (25–27), and a single observational study evaluated rifampin concentrations in routine conditions in Peru (28), none evaluated, under controlled conditions, the concentration of rifampin and its concentration-dependent activity in Latin American patients. Prior evidence indicates important population variability in these measures for antituberculous drugs (29–32). No other late-stage clinical study has explored efficacy as a function of the pharmacokinetic/pharmacodynamic parameter thought to best predict rifampin efficacy (area under the plasma concentration–time curve [AUC]/minimum inhibitory concentration [MIC]) (22, 33–35). The present clinical trial, entitled “Evaluation of High-Dose Rifampin in Patients with New, Smear-Positive Tuberculosis” (HIRIF), was designed to examine the concept that increased rifampin doses could shorten standard therapy for tuberculosis and

improve treatment outcomes without increased toxicity. The primary objectives of this study were to assess the differences across three daily oral doses of rifampin (10, 15, and 20 mg/kg) in plasma concentrations of rifampin, change in elimination rate of *Mycobacterium tuberculosis* in sputum, and frequency of grade 2 or higher adverse events (AEs) related to rifampin. Here we report the primary efficacy and safety endpoints. The pharmacokinetic results have been previously published (36), and some of the results of this study have been previously reported in the form of abstracts (37–39).

Methods

Design

HIRIF was a blinded, randomized, controlled phase 2 clinical trial registered under an investigational new drug application with the U.S. Food and Drug Administration (106635) and with ClinicalTrials.gov (NCT01408914) investigating the pharmacokinetics, efficacy, and safety of higher doses of rifampin in patients with pulmonary tuberculosis. Participants were identified and referred for enrollment from health centers in Lima, Peru between September 2013 and February 2015. Eligible participants were previously untreated adults (18–60 yr old) with smear-positive disease ($\geq 2+$) and strains susceptible to isoniazid and rifampin. Exclusion criteria included a contraindication to rifampin and certain extrapulmonary manifestations of tuberculosis. All participants provided written informed consent. Additional details on study methods are provided in the published protocol (40) and the online supplement.

Intervention

Participants were randomly allocated (1:1:1) in blocks of varying size to receive rifampin at 10, 15, or 20 mg/kg/d during the 8-week intensive phase (for dosing by weight band, see Table E1 in the online supplement). The randomization sequence was generated by the unblinded study statistician and programmed into the electronic data capture system. All other study staff (including laboratory staff) were blinded, except pharmacy staff who received the arm assignment from the electronic data capture

system and prepared study-drug kits. Standard doses of rifampin and companion drugs (isoniazid, pyrazinamide, and ethambutol) were delivered through fixed-dose combinations (FDCs) supplied by Macleods Pharmaceuticals (2). These were supplemented by 150-mg capsules of rifampin and matched placebo donated by Sanofi. During the 18-week continuation phase, participants received isoniazid supplied by Micro Labs Limited and rifampin, both dosed at 10 mg/kg three times weekly. Participants received pyridoxine at 50 mg, three times weekly,

throughout treatment. Treatment was ambulatory and directly observed.

Protocol-defined safety halts occurred when rifampin-related serious AEs (SAEs) met prespecified halting criteria (Table E2) (40). During safety halts, all subjects were treated with a standard regimen according to local guidelines (with an intensive-phase rifampin dose of 10 mg/kg/d). When the sponsor and the data safety and monitoring board determined that the study could resume, participants in the intensive phase of treatment reinitiated their assigned experimental doses.

Outcomes

The primary pharmacokinetic/pharmacodynamic (PK/PD) endpoint, AUC of rifampin in plasma at steady state divided by the MIC for 99.9% of *M. tuberculosis* (MIC_{99.9}), has been previously reported (36). The primary efficacy endpoint was the change in elimination rate of *M. tuberculosis* log₁₀ colony-forming units per milliliter in overnight pooled sputum samples collected throughout the intensive phase and cultured on 7H11 Middlebrook medium. Secondary efficacy endpoints included proportion of culture

Figure 1. Study flow for efficacy analyses. *One patient had an indeterminate Hain result. †Clinical or radiological signs suggestive of pericardial, pleural, or miliary TB. ‡Serum alanine aminotransferase more than two times the upper limit of normal. §Hemoglobin concentration <7.0 g/dl or platelet count <150,000/mm³. ¶Screening was not completed in four patients because of the implementation of protocol-defined study halts, and in one patient because of an inoperative Hain machine. CFU = colony-forming units; ET = early termination; INH = isoniazid; LTFU = lost to follow-up; PK = pharmacokinetics; RIF = rifampin; TB = tuberculosis.

conversion in Löwenstein-Jensen (LJ) medium at 8 weeks, and proportion of unfavorable outcomes (treatment failure, recurrence after cure, or death) at 12 months. We classified recurrent disease by identical 24-locus mycobacterial interspersed repetitive units–variable number tandem repeat (MIRU-VNTR) sequencing as relapsed disease, and by nonidentical MIRU-VNTR as reinfection. The primary safety endpoint was the frequency of grade 2 or higher rifampin-related AEs, per the Division of Microbiology and Infectious Diseases (National Institute of Allergy and Infectious Diseases) Adult Toxicity Table (41), during the intensive phase and up to 4 weeks thereafter.

Statistical Analysis

We defined the intention-to-treat (ITT) population as all participants who received at least one dose of study medication. The modified intention-to-treat (mITT) population (prespecified as primary for efficacy) included those in the ITT population with more than one nonmissing, detectable, and plausible quantitative culture result. The per-protocol (PP) population included those in the mITT population whose experimental rifampin dose was not affected by study halts. Sensitivity analyses were performed on all participants with at least one log₁₀ cfu/ml count. We analyzed log₁₀ cfu/ml

using linear and nonlinear mixed effects models in NONMEM 7.2 (ICON plc), Perl-speaks-NONMEM 4.4.0 (Uppsala University), and R version 3.3.2 (R Foundation for Statistical Computing). Monophasic and biphasic structural models were fitted, by treatment arm and pharmacokinetic exposure, using the M3 partial likelihood method and Laplacian estimation with interaction.

We used the one-sided Cochran-Armitage test for trend with 5% significance to compare proportions of participants who experienced at least one grade 2 or higher rifampin-related AE. We used the log-rank test to compare the time to AEs. We performed similar tests for secondary analyses of rifampin-related hepatic AEs and SAEs. A sensitivity analysis excluded AEs that occurred during study halts in the 15- and 20-mg/kg arms. We performed all safety analyses in Stata/SE 14.2 (StataCorp LLC).

Results

We screened 351 patients for eligibility; of these, 180 were randomized to receive study drug, 60 to each arm (Figure 1). Three study halts occurred during the study. The data safety and monitoring board recommended restarting enrollment and study dosing after each of the halts. Details on the study halts are provided in Table E2. The trial

ended when all enrolled participants completed follow-up.

The mITT dose–efficacy analysis included 174 participants. Six participants were excluded for inadequate data: four had only one colony-forming unit observation; one had all colony-forming unit observations below the limit of detection (<10 cfu/ml); and one had implausibly discrepant colony-forming units in two samples obtained 9 days apart. The PP dose–efficacy analysis included 132 patients in the mITT population whose intensive-phase rifampin dosing was unaffected by study halts. The mITT and PP exposure–efficacy analyses included 163 and 126 patients from the respective dose–efficacy populations who completed a PK sampling visit (Figure 1). All 180 participants were included in the safety analysis. Most participants were young (median age, 25 yr) and male (63.3%). Median weight was slightly higher in the 15-mg/kg arm, 55.4 kg (interquartile range [IQR], 49.2–60.5 kg) compared with 53.5 kg (IQR, 50.6–59.6 kg) and 54.9 kg (IQR, 50.3–61.7 kg) in the 10- and 20-mg/kg arms, respectively. More than one-half of participants had a baseline smear grade of 3+ (51.7%); slightly more patients in the 15-mg/kg arm (60.0%) than in the 10-mg/kg (46.7%) and 20-mg/kg (48.3%) arms had a 3+ smear grade. The mean baseline mycobacterial load was 5.2 log₁₀ cfu/ml. The mean baseline loads were 4.8

Table 1. Participant Characteristics by Treatment Arm*

Characteristic	Rifampin Dose			Total (N = 180)
	10 mg/kg (n = 60)	15 mg/kg (n = 60)	20 mg/kg (n = 60)	
Age, yr, median (IQR)	24 (21–37)	25 (20–35)	27 (22–37)	25 (21–37)
Female	21 (35.0)	19 (31.7)	26 (43.3)	66 (36.7)
Weight, kg, median (IQR) [†]	53.5 (50.6–59.6)	55.4 (49.2–60.5)	54.9 (50.3–61.7)	54.1 (50.5–60.8)
BMI, kg/m ² , median (IQR) [†]	21.2 (19.2–23.3)	21.2 (19.5–23.9)	21.4 (19.9–24.2)	21.2 (19.6–23.7)
Cavitary disease (n = 178)	23 (39.0)	22 (37.3)	21 (35.0)	66 (37.1)
HIV positive	2 (3.3)	2 (3.3)	1 (1.7)	5 (2.8)
Diabetes (n = 179)	0 (0)	0 (0)	1 (1.7)	1 (0.6)
Smear grade at baseline				
2+	32 (53.3)	24 (40.0)	31 (51.7)	87 (48.3)
3+	28 (46.7)	36 (60.0)	29 (48.3)	93 (51.7)
Baseline log ₁₀ cfu/ml, median (IQR) (n = 175)	5.0 (4.4–5.8)	5.0 (4.5–6.2)	5.1 (4.6–5.8)	5.0 (4.5–5.9)
Baseline TTP, d, median (IQR)	4.3 (1.6–5.0)	4.0 (0.1–5.2)	4.4 (0.1–5.0)	4.3 (0.1–5.0)
MIC _{99.9} , median (IQR)	0.2 (0.1–0.2)	0.2 (0.1–0.3)	0.2 (0.1–0.2)	0.2 (0.1–0.2)
AUC _{0–6} /MIC _{99.9} , median (IQR) (n = 168)	115.7 (59.1–197.6)	202.0 (144.1–460.3)	284.4 (175.3–399.2)	193.8 (112.0–344.4)
Rifampin dose, mg, median (IQR)	450 (450–600)	900 (600–900)	975 (750–1,200)	750 (600–900)

Definition of abbreviations: AUC_{0–6} = area under the plasma concentration–time curve from 0 to 6 h; BMI = body mass index; IQR = interquartile range; MIC_{99.9} = minimum inhibitory concentration for 99.9% of *Mycobacterium tuberculosis*; TTP = time-to-positivity in BACTEC MGIT 960.

*Values shown represent n (%) unless otherwise specified.

[†]Values for weight and BMI shown are those obtained at the randomization visit.

and 5.5 log₁₀ cfu/ml among participants with 2+ and 3+ smear grades, respectively. Cavitory disease was present in 66 (37.1%). HIV coinfection and diabetes were rare, present in five participants (2.8%) and one participant (0.6%). Other than weight and smear grade, the distributions of sociodemographic characteristics, comorbidities, clinical indicators of disease severity, and mycobacterial load were similar across arms. Participants received median rifampin doses of 450 mg (IQR, 450–600 mg), 900 mg (IQR, 600–900 mg), and 975 mg (IQR, 750–1,200 mg) in the 10-, 15-, and 20-mg/kg arms, respectively (Table 1).

Efficacy

Efficacy outcomes are shown in Tables 2 and 3, and Figure 2. A monophasic model of bacterial elimination best described the data. We estimated the difference in elimination rate between the treatment arms as -0.011 (95% confidence interval [CI], -0.025 to 0.002) log₁₀ cfu/ml/d for each 5-mg/kg increase in rifampin dose in the mITT population ($P = 0.230$). In the PP population, the difference was -0.022 (95% CI, -0.046 to -0.002 ; $P = 0.022$). The elimination rate in the mITT population did not increase with plasma rifampin AUC_{0–6} (area under the plasma concentration–time curve from 0 to 6 h) (-0.001 log₁₀ cfu/ml/d per 1 log increase in AUC_{0–6}; 95% CI, -0.003 to 0.001 ; $P = 0.750$) or AUC_{0–6}/MIC_{99.9} (-0.002 log₁₀ cfu/ml/d per 1 log increase in AUC_{0–6}/MIC_{99.9}; 95% CI, -0.005 to 0.002 ; $P = 0.330$). The elimination rate in the PP population increased with plasma rifampin AUC_{0–6} (-0.017 log₁₀ cfu/ml/d per 1 log increase in AUC_{0–6}; 95% CI, -0.029 to

-0.007 ; $P = 0.011$) but not with AUC_{0–6}/MIC_{99.9} (-0.010 log₁₀ cfu/ml/d per 1 log increase in AUC_{0–6}/MIC_{99.9}; 95% CI, -0.021 to 0.000 ; $P = 0.053$). These results were not affected by adjustment for important baseline covariates including age, sex, and extent of radiological disease. The proportion of participants with 8-week culture conversion in LJ medium was nearly identical across the three treatment arms (76.7%, 73.3%, and 75.0%, respectively). Eleven participants (6.1%) experienced unfavorable outcomes by the end of the 12-month follow-up (7.5% of the 147 participants in whom 12-mo status could be assessed). Five participants (2.8%) experienced treatment failure. Of these, three (60%) were in the 10-mg/kg arm. Six more (3.3%) experienced recurrent disease, three (50%) in the 10-mg/kg arm; two (33%) were confirmed relapses by MIRU-VNTR sequencing. There were no deaths during the study. A detailed description of all treatment failure and recurrence outcomes is provided in Table E3.

Safety

The frequency of grade 2 or higher rifampin-related AEs was similar across treatment arms: 26, 31, and 23 participants (43.3%, 51.7%, and 38.3%) in the 10-, 15-, and 20-mg/kg arms, respectively, had at least one event ($P = 0.7092$) (Tables 4 and E4). Time to first grade 2 or higher rifampin-related AE was also similar across arms ($P = 0.3610$; Figure E1). The frequency and time to grade 2 or higher rifampin-related hepatic AEs and the frequency of SAEs were similar across arms (Tables 4 and 5). Results were similar when we excluded AEs that occurred during

study halts in the experimental arms (Table E5).

The distribution of participants with at least one AE stratified by organ system is shown in Table E6. The most common AEs were hepatic (24.4%), gastrointestinal (14.4%), musculoskeletal (12.2%), and respiratory (11.7%). Nine SAEs occurred in eight participants during the entire study. Of these, six SAEs occurred in five participants during the safety period; four were rifampin-related (Table 5). A detailed description of all SAEs is provided in Table E7.

Discussion

The present study is the first clinical trial to show dose and exposure response of higher doses of rifampin under conditions of combination therapy. Higher rifampin doses and exposure resulted in an increase in the rate of sputum culture sterilization. These findings build on pharmacokinetic results from the present study and others showing that rifampin exposure increases at least proportionally with higher doses (25, 36), by clearly demonstrating that the elimination rate is also related to rifampin AUC_{0–6}. These dose- and exposure-related efficacy findings are consistent with the results of the PanACEA (Pan-African Consortium for the Evaluation of Antituberculosis Antibiotics) MAMS-TB-01 trial (NCT01785186). This latter study found that rifampin doses of 20 mg/kg are unlikely to permit treatment shortening and that the optimal dose of rifampin is likely higher than 20 mg/kg (26). The statistically nonsignificant increase in culture negativity in the 20-mg/kg arm of

Table 2. Decrease in Viable Colony-Forming Unit Counts of *Mycobacterium tuberculosis* by Rifampin Dose or Exposure to Rifampin

Dose/Exposure Variable	n	Δ log ₁₀ cfu/ml/d	95% CI*	P Value
5-mg/kg increase in rifampin dose				
mITT population	174	-0.011	-0.025 to 0.002	0.230
PP population	132	-0.022	-0.046 to -0.002	0.022
1 log increase in rifampin AUC _{0–6}				
mITT population	163	-0.001	-0.003 to 0.001	0.750
PP population	126	-0.017	-0.029 to -0.007	0.011
1 log increase in rifampin AUC _{0–6} /MIC _{99.9}				
mITT population	163	-0.002	-0.005 to 0.002	0.330
PP population	126	-0.010	-0.021 to 0.000	0.053

Definition of abbreviations: AUC_{0–6} = area under the plasma concentration–time curve from 0 to 6 hours; CI = confidence interval; MIC_{99.9} = minimum inhibitory concentration for 99.9% of *Mycobacterium tuberculosis*; mITT = modified intention-to-treat; PP = per protocol.

*The 95% CIs were obtained using the sandwich estimator.

Table 3. Secondary Efficacy Outcomes*

Outcome	Rifampin Dose			Total (N = 180)
	10 mg/kg (n = 60)	15 mg/kg (n = 60)	20 mg/kg (n = 60)	
Culture conversion in Löwenstein-Jensen medium at 8 wk				
Converted	46 (76.7)	44 (73.3)	45 (75.0)	135 (75.0)
Did not convert	3 (5.0)	5 (8.3)	7 (11.7)	15 (8.3)
Contaminated Week 8 cultures	4 (6.7)	1 (1.7)	0 (0)	5 (2.8)
Week 8 cultures not available	7 (11.7)	10 (16.7)	8 (13.3)	25 (13.9)
Early discontinuation	6 (10.0)	9 (15.0)	8 (13.3)	23 (12.8)
Lost to follow-up	1 (1.7)	1 (1.7)	0 (0)	2 (1.1)
Treatment outcome at 12 mo				
Recurrence-free cure	44 (73.3)	46 (76.7)	46 (76.7)	136 (75.6)
Treatment failure	3 (5.0)	1 (1.7)	1 (1.7)	5 (2.8)
Recurrence after cure	3 (5.0)	1 (1.7)	2 (3.3)	6 (3.3)
Relapse [†]	2 (3.3)	0 (0)	0 (0)	2 (1.1)
Reinfection [‡]	0 (0)	1 (1.7)	0 (0)	1 (0.6)
Recurrence [§]	1 (1.7)	0 (0)	2 (3.3)	3 (1.7)
Death	0 (0)	0 (0)	0 (0)	0 (0)
Outcome not evaluable	10 (16.7)	12 (20.0)	11 (18.3)	33 (18.3)
Early discontinuation	8 (13.3)	11 (18.3)	10 (16.7)	29 (16.1)
Lost to follow-up	2 (3.3)	1 (1.7)	1 (1.7)	4 (2.2)

*Values shown represent n (%).

[†]Two patients had relapse diagnosed by identical pretreatment and post-treatment *Mycobacterium tuberculosis* isolates, with 24 of 24 matching MIRU-VNTR (mycobacterial interspersed repetitive units–variable number tandem repeat) loci.

[‡]One patient had reinfection diagnosed by nonidentical pretreatment and post-treatment *M. tuberculosis* isolates, with 9 of 20 matching MIRU-VNTR loci.

[§]Recurrent tuberculosis was diagnosed without positive culture (and without MIRU-VNTR) in three patients: two were diagnosed by a positive sputum smear, and one was diagnosed clinically.

the RIFATOX (Evaluation of High-Dose Rifampicin Toxicity in Pulmonary Tuberculosis) trial (ISRCTN55670677) also suggests that this dose may not be sufficient for treatment shortening (42).

The present study characterized the consistent impact of both rifampin dose and exposure on efficacy. Other analyses were not statistically significant: the analysis of 8-week culture conversion in LJ medium,

the mITT analysis of dose and exposure as measured by AUC_{0-6} and $AUC_{0-6}/MIC_{99.9}$ on the primary endpoint of change in $\log_{10}$ cfu/ml, and the PP analysis of exposure as measured by $AUC_{0-6}/MIC_{99.9}$. There are several potential explanations for these findings. First, the study was not powered for the 8-week binary, relatively insensitive, secondary endpoint. Second, the mITT efficacy estimates were affected by protocol-defined study halts. The use of standard rifampin dosing among all participants during study halts may have diminished the effect of the higher doses, biasing the effect estimate of dose toward the null. This explanation is supported by the significant effect of dose detected in the PP analysis, which includes only the 132 participants (73.3%) who received their assigned rifampin dose for the full intensive phase. Study halts may have also biased the effect of exposure toward the null in the mITT population. Because rifampin exposure was not measured during study halts, the estimates of rifampin exposure in experimental arm participants who received standard rifampin dosing during study halts likely overestimate overall rifampin exposure. Given the effect of study halts on both dose- and exposure-response

Figure 2. Predictions of bacillary elimination for the three treatment arms derived from the partial likelihood model showing observed data (dark gray) and model predictions (light gray) above and below the limit of detection of colony counting, respectively.

Table 4. Participants Experiencing Grade 2 or Higher Rifampin-related Adverse Events and Hepatic Adverse Events during the Safety Period, by Treatment Arm*

Variable	Rifampin Dose			P Value
	10 mg/kg (n = 60)	15 mg/kg (n = 60)	20 mg/kg (n = 60)	
≥1 grade 2+ RIF-related AE	26 (43.3)	31 (51.7)	23 (38.3)	0.7092 [†]
Time to first grade 2+ RIF-related AE, d, median (IQR)	29.5 (14–39)	26 (9–42)	25 (13–37)	0.3610 [‡]
≥1 grade 2+ hepatic RIF-related AE	16 (26.7)	14 (23.3)	14 (23.3)	0.6645 [†]
Time to first grade 2+ RIF-related hepatic AE, d, median (IQR)	31.5 (14–40.5)	36.5 (26–57)	42.5 (26–63)	0.8357 [‡]

Definition of abbreviations: AE = adverse event; IQR = interquartile range; RIF = rifampin.

*Values shown represent n (%) unless otherwise specified. The safety period was defined as 12 weeks after randomization or 4 weeks after the last experimental rifampin dose, whichever occurred later.

[†]One-sided Cochran-Armitage test for trend with 5% significance.

[‡]Global log-rank test.

estimates in the mITT population, we believe that the PP population is less biased than the mITT population for the efficacy analysis, even though the PP population was the prespecified secondary analysis population.

Third, 1:1:1 randomization yielded median rifampin doses of 450, 900, and 975 mg in the 10-, 15-, and 20-mg/kg arms, respectively. Because median weights for all three arms fell in the 53- to 55-kg range, we expected that median rifampin doses would approach 600, 900, and 1,200 mg, the assigned rifampin doses for the 55- to 70-kg weight band. However, more than one-half of study participants in the 10-mg/kg arm were in the two lowest rifampin weight bands, more than one-half in the 15-mg/kg arm were in the two highest weight bands, and the 20-mg/kg arm had the same number of participants in the two lowest and two highest weight bands. Consequently, the median difference in dose between the 15- and 20-mg/kg arms was only 75 mg. These subtle differences in assigned rifampin doses (and consequent exposures) could have

biased our results, reducing the apparent activity of the 20-mg/kg arm and the ability to distinguish a dose-response effect.

The effect of rifampin AUC_{0–6}/MIC_{99.9} (the primary PK/PD endpoint) on sputum culture sterilization was of only borderline statistical significance. Prior preclinical studies have suggested that rifampin AUC/MIC is a strong predictor of concentration-dependent killing (22, 33). Clinical studies of patients receiving combination therapy have shown mixed results. Others have shown that rifampin AUC alone is an important predictor of clinical outcomes (32) and that its predictive ability may be bolstered by MIC information (34). However, a more recent study found no relationship between rifampin AUC/MIC and 2-month culture conversion (35). The distribution of susceptible rifampin MICs in our sample was higher than in the latter study (35), albeit within previously published ranges (34, 43). We did not find that rifampin MICs varied across levels of AUC to suggest that the effect of rifampin

AUC_{0–6}/MIC_{99.9} was attenuated by selection bias. Our findings suggest that rifampin AUC_{0–6}/MIC_{99.9} was not as strong a predictor of sputum sterilization as rifampin AUC_{0–6}, and that there may be a role for higher doses of rifampin in populations with susceptible rifampin MICs on the higher end of the distribution. In addition, the present analysis of rifampin MIC_{99.9} does not capture the activity of rifampin metabolites, which may contribute to treatment efficacy.

Despite our inclusion criterion of baseline sputum smear grade of 2+ or higher, we found a mean baseline mycobacterial load of 5.2 log₁₀ cfu/ml, approximately 10-fold lower than in previous studies (8, 44). Low baseline loads have been reported in other settings, such as Hong Kong, Tanzania, and South Africa (25, 27, 45, 46). In the present study, low baseline loads may have contributed to the selection of a monophasic model of elimination by rifampin dose. This is distinct from the biphasic model selected to best represent the data in a study of

Table 5. Participants Experiencing Serious Adverse Events by Treatment Arm*

Variable	Rifampin Dose			P Value [†]
	10 mg/kg (n = 60)	15 mg/kg (n = 60)	20 mg/kg (n = 60)	
≥1 SAE during entire study	2 (3.3)	1 (1.7)	5 (8.3)	0.0919
≥1 SAE during safety period [‡]	1 (1.7)	1 (1.7)	3 (5.0)	0.1333
≥1 RIF-related SAE during safety period [‡]	1 (1.7)	1 (1.7)	2 (3.3)	0.2679

Definition of abbreviations: RIF = rifampin; SAE = serious adverse event.

*Values shown represent n (%).

[†]One-sided Cochran-Armitage test for trend with 5% significance.

[‡]The safety period was defined as 12 weeks after randomization or 4 weeks after the last experimental rifampin dose, whichever occurred later.

fluoroquinolone-containing regimens (8). Our findings are consistent with those of a 14-day early bactericidal activity study by the TB Alliance, which did not show a pronounced initial fall in the mycobacterial load among patients receiving standard therapy who had a mean baseline load of $5.399 \log_{10}$ cfu/ml (47), and with an observation that the 0- to 2-day early bactericidal activity of isoniazid-based treatments is strongly correlated with pretreatment mycobacterial load (46).

We found no difference in the secondary efficacy outcomes of 8-week culture conversion or in the frequency of treatment failure and disease recurrence across arms. The frequencies of treatment failure and recurrence we observed are comparable to those of standard therapy in other clinical trials (16, 19, 48). However, these relatively rare events occurred disproportionately in the 10-mg/kg arm, in which the median rifampin dose was 450 mg daily. We previously reported that 81% of the patients in the 20-mg/kg arm and only 33% of those of in the 10-mg/kg arm achieved a maximum drug concentration in plasma ($C_{\max}$) greater than 8 mg/L, the low end of the targeted range (36). Others have reported that standard rifampin dosing through FDCs has resulted in suboptimal $C_{\max}$ values less than 8 mg/L (35). Taken together, these findings suggest that standard doses of rifampin at 10 mg/kg may have compromised treatment efficacy and that the 20-mg/kg dose should be considered for dosing to maximize efficacy

pending results of studies evaluating even higher doses. This has important implications for the use of FDCs, which are currently recommended over separate drug formulations in the treatment of drug-susceptible tuberculosis (49). Current guidelines recommend maximum rifampin doses of 600 mg daily (2), or 750 mg daily when using FDCs containing 150 mg of rifampin each (50). Under programmatic settings, supplementation with loose rifampin would be required to achieve higher doses.

We did not observe an increased risk of toxicity with rifampin doses up to 20 mg/kg delivered over 8 weeks. We found similar results when we excluded adverse events that occurred during study halts in the experimental arms. Our power calculation assumed a 10% frequency of grade 2 or higher rifampin-related AEs in the control arm. With 43.3% of control arm participants having at least one event, there was increased power over what was anticipated: we had more than 80% power to rule out a toxicity difference of 20% between the control and experimental arms. Our safety findings are consistent with data from several studies. The maximum-dose tolerability study, HIGHRIF1 (NCT01392911), found no SAEs among participants who received rifampin doses up to 35 mg/kg over 14 days (25). The RIFATOX phase 2 trial found that rifampin doses up to 20 mg/kg delivered over 16 weeks did not result in a significant increase in AEs (42). Finally, the more recent

MAMS-TB-01 trial found that a rifampin dose of 35 mg/kg delivered as part of combination therapy over 12 weeks resulted in similar frequencies of grade 3 or higher AEs to standard therapy (26). Taken together, there is a growing body of evidence from controlled studies to suggest that concerns about the toxicity of rifampin may have taken on too much importance, relative to efficacy. In a future publication, we plan to pool data from this trial with those from another phase 2 trial of high-dose rifampin conducted by PanACEA, HIGHRIF2 (NCT00760149) (27), with the goal of improving statistical power for efficacy and safety evaluations.

In summary, this is the first controlled trial to show dose and exposure response of higher doses of rifampin on sputum culture sterilization under conditions of combination therapy. Rifampin doses of up to 20 mg/kg/d were safe compared with the standard dose. These findings support reconsideration of currently recommended standard dosing guidelines and the continued investigation of higher doses of rifampin, beyond 20 mg/kg, for potential treatment shortening. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Acknowledgment: The authors thank the patients and their families who made this study possible. The authors also thank the health care workers at the participating health centers in Lima Ciudad and Lima Este.

References

- World Health Organization. Global tuberculosis report 2017. Geneva, Switzerland: World Health Organization; 2017. Report No. WHO/HTM/TB/2017.23.
- World Health Organization. Treatment of tuberculosis: guidelines, 4th ed. Geneva: World Health Organization; 2010. Report No. WHO/HTM/TB/2009.420.
- Fox W, Ellard GA, Mitchison DA. Studies on the treatment of tuberculosis undertaken by the British Medical Research Council tuberculosis units, 1946–1986, with relevant subsequent publications. *Int J Tuberc Lung Dis* 1999;3(10, Suppl 2): S231–S279.
- National Research Institute of Tuberculosis, Warsaw; World Health Organization. A comparative study of daily followed by twice or once weekly regimens of ethambutol and rifampicin in retreatment of patients with pulmonary tuberculosis: the results at 1 year: a cooperative tuberculosis chemotherapy study in Poland. *Tubercle* 1975;56:1–26.
- Zierski M, Bek E, Bergson H, Kucharska A. A comparative study of daily followed by twice- or once-weekly regimens of ethambutol and rifampicin in the retreatment of patients with pulmonary tuberculosis: second report. *Tubercle* 1976;57:105–113.
- Hong Kong Tuberculosis Treatment Services; Brompton Hospital; British Medical Research Council. A controlled trial of daily and intermittent rifampicin plus ethambutol in the retreatment of patients with pulmonary tuberculosis: results up to 30 months. *Tubercle* 1975;56:179–189.
- van Ingen J, Aarnoutse RE, Donald PR, Diacon AH, Dawson R, Plemper van Balen G, et al. Why do we use 600 mg of rifampicin in tuberculosis treatment? *Clin Infect Dis* 2011;52:e194–e199.
- Rustomjee R, Lienhardt C, Kanyok T, Davies GR, Levin J, Mthiyane T, et al.; Gatifloxacin for TB (OFLOTUB) Study Team. A phase II study of the sterilising activities of ofloxacin, gatifloxacin and moxifloxacin in pulmonary tuberculosis. *Int J Tuberc Lung Dis* 2008;12:128–138.
- Nuermberger EL, Yoshimatsu T, Tyagi S, O'Brien RJ, Vernon AN, Chaisson RE, et al. Moxifloxacin-containing regimen greatly reduces time to culture conversion in murine tuberculosis. *Am J Respir Crit Care Med* 2004;169:421–426.
- Nuermberger EL, Yoshimatsu T, Tyagi S, Williams K, Rosenthal I, O'Brien RJ, et al. Moxifloxacin-containing regimens of reduced duration produce a stable cure in murine tuberculosis. *Am J Respir Crit Care Med* 2004;170:1131–1134.
- Burman WJ, Goldberg S, Johnson JL, Muzanye G, Engle M, Mosher AW, et al. Moxifloxacin versus ethambutol in the first 2 months of treatment for pulmonary tuberculosis. *Am J Respir Crit Care Med* 2006;174:331–338.

12. Dorman SE, Johnson JL, Goldberg S, Muzanye G, Padayatchi N, Bozeman L, *et al.*; Tuberculosis Trials Consortium. Substitution of moxifloxacin for isoniazid during intensive phase treatment of pulmonary tuberculosis. *Am J Respir Crit Care Med* 2009;180: 273–280.
13. Johnson JL, Hadad DJ, Dietze R, Maciel ELN, Sewali B, Gitta P, *et al.* Shortening treatment in adults with noncavitary tuberculosis and 2-month culture conversion. *Am J Respir Crit Care Med* 2009;180: 558–563.
14. Conde MB, Efron A, Loredó C, De Souza GRM, Graça NP, Cezar MC, *et al.* Moxifloxacin versus ethambutol in the initial treatment of tuberculosis: a double-blind, randomised, controlled phase II trial. *Lancet* 2009;373:1183–1189.
15. Dorman SE, Goldberg S, Stout JE, Muzanyi G, Johnson JL, Weiner M, *et al.*; Tuberculosis Trials Consortium. Substitution of rifapentine for rifampin during intensive phase treatment of pulmonary tuberculosis: study 29 of the Tuberculosis Trials Consortium. *J Infect Dis* 2012; 206:1030–1040.
16. Jindani A, Harrison TS, Nunn AJ, Phillips PPJ, Churchyard GJ, Charalambous S, *et al.*; RIFAQUIN Trial Team. High-dose rifapentine with moxifloxacin for pulmonary tuberculosis. *N Engl J Med* 2014; 371:1599–1608.
17. ClinicalTrials.gov. TBTC study 31: rifapentine-containing tuberculosis treatment shortening regimens (S31/A5349) [accessed 2017 Dec 15]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02410772>.
18. Mitchison DA. The action of antituberculosis drugs in short-course chemotherapy. *Tubercle* 1985;66:219–225.
19. Jindani A, Nunn AJ, Enarson DA. Two 8-month regimens of chemotherapy for treatment of newly diagnosed pulmonary tuberculosis: international multicentre randomised trial. *Lancet* 2004; 364:1244–1251.
20. Long MW, Snider DE Jr, Farer LS. U.S. Public Health Service cooperative trial of three rifampin–isoniazid regimens in treatment of pulmonary tuberculosis. *Am Rev Respir Dis* 1979;119:879–894.
21. Jindani A, Aber VR, Edwards EA, Mitchison DA. The early bactericidal activity of drugs in patients with pulmonary tuberculosis. *Am Rev Respir Dis* 1980;121:939–949.
22. Jayaram R, Gaonkar S, Kaur P, Suresh BL, Mahesh BN, Jayashree R, *et al.* Pharmacokinetics–pharmacodynamics of rifampin in an aerosol infection model of tuberculosis. *Antimicrob Agents Chemother* 2003; 47:2118–2124.
23. Sirgel FA, Fourie PB, Donald PR, Padayatchi N, Rustonjee R, Levin J, *et al.* The early bactericidal activities of rifampin and rifapentine in pulmonary tuberculosis. *Am J Respir Crit Care Med* 2005;172: 128–135.
24. Poole G, Stradling P, Worlledge S. Potentially serious side effects of high-dose twice-weekly rifampicin. *BMJ* 1971;3:343–347.
25. Boeree MJ, Diacon AH, Dawson R, Narunsky K, du Bois J, Venter A, *et al.*; PanACEA Consortium. A dose-ranging trial to optimize the dose of rifampin in the treatment of tuberculosis. *Am J Respir Crit Care Med* 2015;191:1058–1065.
26. Boeree MJ, Heinrich N, Aarnoutse R, Diacon AH, Dawson R, Rehal S, *et al.*; PanACEA Consortium. High-dose rifampicin, moxifloxacin, and SQ109 for treating tuberculosis: a multi-arm, multi-stage randomised controlled trial. *Lancet Infect Dis* 2017; 17:39–49.
27. Aarnoutse RE, Kibiki GS, Reither K, Semvua HH, Haraka F, Mtabho CM, *et al.*; PanACEA Consortium. Pharmacokinetics, tolerability, and bacteriological response of rifampin administered at 600, 900, and 1,200 milligrams daily in patients with pulmonary tuberculosis. *Antimicrob Agents Chemother* 2017;61: e01054–17.
28. Requena-Méndez A, Davies G, Ardrey A, Jave O, López-Romero SL, Ward SA, *et al.* Pharmacokinetics of rifampin in Peruvian tuberculosis patients with and without comorbid diabetes or HIV. *Antimicrob Agents Chemother* 2012;56:2357–2363.
29. McIlleron H, Wash P, Burger A, Norman J, Folb PI, Smith P. Determinants of rifampin, isoniazid, pyrazinamide, and ethambutol pharmacokinetics in a cohort of tuberculosis patients. *Antimicrob Agents Chemother* 2006;50:1170–1177.
30. Chigutsa E, Visser ME, Swart EC, Denti P, Pushpakom S, Egan D, *et al.* The SLCO1B1 rs4149032 polymorphism is highly prevalent in South Africans and is associated with reduced rifampin concentrations: dosing implications. *Antimicrob Agents Chemother* 2011;55: 4122–4127.
31. Pasipanodya JG, Srivastava S, Gumbo T. Meta-analysis of clinical studies supports the pharmacokinetic variability hypothesis for acquired drug resistance and failure of antituberculosis therapy. *Clin Infect Dis* 2012;55:169–177.
32. Pasipanodya JG, McIlleron H, Burger A, Wash PA, Smith P, Gumbo T. Serum drug concentrations predictive of pulmonary tuberculosis outcomes. *J Infect Dis* 2013;208:1464–1473.
33. Gumbo T, Louie A, Deziel MR, Liu W, Parsons LM, Salfinger M, *et al.* Concentration-dependent *Mycobacterium tuberculosis* killing and prevention of resistance by rifampin. *Antimicrob Agents Chemother* 2007;51:3781–3788.
34. Gumbo T, Pasipanodya JG, Wash P, Burger A, McIlleron H. Redefining multidrug-resistant tuberculosis based on clinical response to combination therapy. *Antimicrob Agents Chemother* 2014;58: 6111–6115.
35. Rockwood N, Pasipanodya JG, Denti P, Sirgel F, Lesosky M, Gumbo T, *et al.* Concentration-dependent antagonism and culture conversion in pulmonary tuberculosis. *Clin Infect Dis* 2017;64: 1350–1359.
36. Peloquin CA, Velásquez GE, Lecca L, Calderón RI, Coit J, Milstein M, *et al.* Pharmacokinetic evidence from the HIRIF trial to support increased doses of rifampin for tuberculosis. *Antimicrob Agents Chemother* 2017;61:e00038–17.
37. Milstein MB, Velásquez GE, Coit JM, Jimenez J, Tintaya K, Osso EM, *et al.* Tuberculosis clinical trials and safety halts: evaluating for bias in effect estimates [abstract OA-220-14]. Presented at the 48th World Conference on Lung Health of the International Union against Tuberculosis and Lung Disease (the Union). October 10–14, 2017, Guadalajara, Mexico. *Int J Tuberc Lung Dis* 2017;21(11, Suppl 2): S350–S351.
38. Milstein M, Coit J, Velásquez GE, Osso E, Vasquez DV, Garavito ES, *et al.* Active screening for hepatotoxicity: identifying time periods for targeted monitoring in tuberculosis patients [abstract PD-722-13]. Presented at the 48th World Conference on Lung Health of the International Union against Tuberculosis and Lung Disease (the Union). October 10–14, 2017, Guadalajara, Mexico. *Int J Tuberc Lung Dis* 2017;21(11, Suppl 2):S275–S276.
39. Velásquez GE, Brooks MB, Coit JM, Vargas Vasquez D, Sánchez Garavito E, Calderón RI, *et al.* A randomized controlled trial of high-dose rifampin for pulmonary tuberculosis [abstract 39LB]. Presented at the Conference on Retroviruses and Opportunistic Infections. March 4–7, 2018, Boston, MA.
40. Milstein M, Lecca L, Peloquin C, Mitchison D, Seung K, Pagano M, *et al.* Evaluation of high-dose rifampin in patients with new, smear-positive tuberculosis (HIRIF): study protocol for a randomized controlled trial. *BMC Infect Dis* 2016;16:453.
41. National Institute of Allergy and Infectious Diseases. Division of Microbiology and Infectious Diseases adult toxicity table, November 2007, Draft [accessed 2017 Dec 15]. Available from: <https://www.niaid.nih.gov/sites/default/files/dmidadulttox.pdf>.
42. Jindani A, Borgulya G, de Patiño IW, Gonzales T, de Fernandes RA, Shrestha B, *et al.*; International Consortium for Trials of Chemotherapeutic Agents in Tuberculosis, St George's University of London. A randomised phase II trial to evaluate the toxicity of high-dose rifampicin to treat pulmonary tuberculosis. *Int J Tuberc Lung Dis* 2016;20:832–838.
43. Schön T, Jurén P, Giske CG, Chrysanthou E, Sturegård E, Werngren J, *et al.* Evaluation of wild-type MIC distributions as a tool for determination of clinical breakpoints for *Mycobacterium tuberculosis*. *J Antimicrob Chemother* 2009; 64:786–793.
44. Ruslami R, Nijland H, Aarnoutse R, Alisjahbana B, Soeroto AY, Ewalds S, *et al.* Evaluation of high- versus standard-dose rifampin in Indonesian patients with pulmonary tuberculosis. *Antimicrob Agents Chemother* 2006;50:822–823.

45. Sirgel FA, Donald PR, Odhiambo J, Githui W, Umapathy KC, Paramasivan CN, *et al.* A multicentre study of the early bactericidal activity of anti-tuberculosis drugs. *J Antimicrob Chemother* 2000;45: 859–870.
46. De Jager V, van der Merwe L, Venter A, Donald PR, Diacon AH. Time trends in sputum mycobacterial load and two-day bactericidal activity of isoniazid-containing antituberculosis therapies. *Antimicrob Agents Chemother* 2017;61:e02088-16.
47. Diacon AH, Dawson R, von Groote-Bidlingmaier F, Symons G, Venter A, Donald PR, *et al.* Bactericidal activity of pyrazinamide and clofazimine alone and in combinations with pretomanid and bedaquiline. *Am J Respir Crit Care Med* 2015;191:943–953.
48. Merle CS, Fielding K, Sow OB, Gninafon M, Lo MB, Mthiyane T, *et al.*; OFLOTUB/Gatifloxacin for Tuberculosis Project. A four-month gatifloxacin-containing regimen for treating tuberculosis. *N Engl J Med* 2014;371:1588–1598.
49. World Health Organization. Guidelines for treatment of drug-susceptible tuberculosis and patient care, 2017 update. Geneva, Switzerland: World Health Organization; 2017. Report No. WHO/HTM/TB/2017.05.
50. World Health Organization. Operational guide for national tuberculosis control programmes on the introduction and use of fixed-dose combination drugs. Geneva, Switzerland: World Health Organization; 2002. Report No. WHO/CDS/TB/2002.308.