Published in final edited form as:

Crit Care Med. 2017 February ; 45(2): e146–e153. doi:10.1097/CCM.0000000000002063.

Chloride content of fluids used for large volume resuscitation is associated with reduced survival

Ayan Sen, MD, MS, FCCP*, Christopher M Keener, MS, Florentina E Sileanu, MS, Emily Foldes, MS, Gilles Clermont, MD, CM, MS, Raghavan Murugan, MD, MS, FRCP, FCCP, and John A Kellum, MD, MCCM

Center for Critical Care Nephrology and CRISMA Center, Department of Critical Care Medicine, University of Pittsburgh, Pittsburgh, PA, United States

Abstract

Objective—We sought to investigate if the chloride content of fluids used in resuscitation was associated with short and long-term outcomes.

Design—We identified patients who received large volume fluid resuscitation (LVR), defined as greater than 60ml/kg over a 24 hour period. Chloride load was determined for each patient based on the chloride ion concentration of the fluids they received during LVR multiplied by the volume of fluids. We compared the development of hyperchloremic acidosis (HCA), AKI and survival among those with higher and lower chloride loads.

Setting—University Medical Center

Patients—Patients admitted to intensive care units (ICUs) from 2000–2008

Interventions—None

Main Results—Among 4,710 patients receiving LVR HCA was documented in 523 (11%). Crude rates of HCA, AKI and hospital mortality all increased significantly as chloride load increased ($p < 0.001$). However, chloride load was no longer associated with HCA or AKI after controlling for total fluids, age, and baseline severity. Conversely, each 100 mEq increase in chloride load was associated with a 5.5% increase in the hazard of death even after controlling for total fluid volume, age and severity ($p = 0.0015$) over one year.

Corresponding Author: John A Kellum MD, MCCM, Professor of Critical Care Medicine, Medicine, Bioengineering, and Translational & Clinical Science, Director, Center for Critical Care Nephrology, 604, Scaife Hall, 3550 Terrace Street, Pittsburgh, PA 15261. kellum@pitt.edu, Telephone: 412-647-8110, Fax: 412-647-2645.

*Current affiliation: Department of Critical Care Medicine, Mayo Clinic, Scottsdale, AZ, United States

Disclosures

JAK has received consulting fees and research grants from Baxter Healthcare Corporation. None of the other authors report any disclosures.

Copyright form disclosures: Dr. Kellum disclosed other support (Work was funded by Baxter Healthcare) and received funding from Baxter. His institution received funding from Baxter. Dr. Sen's institution received funding from Baxter Healthcare Corporation. Dr. Keener's institution received funding from Baxter. Dr. Sileanu disclosed work for hire. Her institution received funding from Baxter. Dr. Foldes' institution received funding from Baxter Healthcare Corporation. Dr. Clermont received support for article research from the National Institutes of Health (NIH). His institution received funding from Baxter, Astute Medical, NIH, and NSF. Dr. Murugan's institution received support from Baxter Inc. (Grant paid to University of Pittsburgh).

Conclusions—Chloride load is associated with significant adverse effects on survival out to one year even after controlling for total fluid load, age and baseline severity of illness. However the relationship between chloride load and development of HCA or AKI is less clear and further research is needed to elucidate the mechanisms underlying the adverse effects of chloride load on survival.

Keywords

saline; chloride; metabolic acidosis; acute kidney injury; critical illness; resuscitation; survival

Use of crystalloids containing supra-physiologic chloride concentrations has been associated with hyperchloremia, acidosis and more frequent complications when compared to resuscitation with balanced crystalloids (1–6). Hyperchloremia has been shown to be independently associated with increased mortality in the intensive care unit (ICU) and postsurgical patients, and the amount of chloride received via intravenous infusion has been identified as an important and potentially modifiable cause of hyperchloremic acidosis (HCA) (7–9). In a prospective, sequential period study, chloride restriction was associated with a significant reduction in the incidence of acute kidney injury (AKI) and the use of renal replacement therapy (RRT) (6). A later extended analysis of patients receiving chloride-restricted fluids showed persistent decrease of AKI and RRT rates but also found evidence of other confounders such as patient characteristics, clinician expertise or the Hawthorne effect, which could help explain fluctuations in AKI incidence (6, 10). However, in a cluster randomized, double-crossover trial, there was no difference in the incidence of AKI, in-hospital mortality or use of RRT within 90 days after enrollment in a heterogeneous population of ICU patients who received a buffered crystalloid or saline (11). The exposure to saline in this study was small (about 2L) and patients were relatively low risk (APACHE II 14.1, mortality 8%, need for RRT 3%).

Evidence from experimental models of sepsis in animals who receive fluid resuscitation have shown that HCA induces hemodynamic instability (12) and inflammatory response (13, 14), worsens renal dysfunction (13, 14), and decreases survival (12, 14, 15). Fluid choice may therefore impact outcomes in patients requiring infusion of large volumes of IV fluids. While studies have examined mortality in patients receiving crystalloids with different chloride concentrations, the relative contributions of chloride load and fluid volume to mortality risk remain unclear (16–19) as does the relationship between chloride load, HCA, AKI and mortality. Furthermore, the effects of HCA on long-term outcomes are largely unknown.

Materials and Methods

After receiving approval from the University of Pittsburgh Institutional Review Board, we interrogated the High Density Intensive Care (HiDenIC-8) database which contains data from all patients admitted to intensive care units (ICUs) from 2000 to 2008 merging several computerized sources. Patient records have also been linked to national databases (National Death Index database, the Social Security Administration Death Master File, US Renal Data Systems) to ascertain vital status and dialysis out to 1 year.

We defined large-volume resuscitation (LVR) as 60ml/kg or greater in a 24hr period and included patients receiving this volume in our analysis cohort. Patients with a history of hemodialysis or kidney transplant, a baseline creatinine ≥ 4 mg/dl recorded prior to hospital admission, those with insufficient data to determine AKI status or those who developed HCA or AKI prior to LVR or received LVR in the OR were excluded. Figure 1 outlines the selection of the study population.

We determined baseline demographic, comorbidity, and severity of illness characteristics for all patients in the LVR cohort, including derived variables which have been described previously and include “suspected sepsis”, “hypotensive index”, and the Acute Physiology Score (APS-III) derived from the “acute physiology” portion of the Acute Physiology, Age, Chronic Health Evaluation (APACHE III) score (20, 21). We calculated the total chloride ion concentration of each fluid used for resuscitation in order to assess the relationship between the chloride load and amount of fluid administered during LVR. Total chloride load was determined for each patient as follows: $(C_1 \cdot V_1) + (C_2 \cdot V_2) + (C_3 \cdot V_3) + (C_n \cdot V_n) = \text{Chloride Load (mEq)}$

Where C_n = chloride concentration (mEq) per liter of fluid n and V_n is the volume of that fluid administered in the 24 hours beginning with the start of LVR. For example, if a patient received 3 litres of 0.9% saline and 2 litres of lactated Ringer’s, total chloride load would be $(154 \cdot 3 = 462) + (109 \cdot 2 = 218) = 680$ mEq.

Outcomes included hyperchloremic metabolic acidosis (HCA), AKI and mortality. HCA was determined as explained in Table S1. Briefly, we defined metabolic acidosis as an arterial base deficit > 2 mEq/l and defined HCA as predominate etiology of metabolic acidosis after excluding other causes. AKI was determined using the widely validated KDIGO criteria (Table S2) (22). Baseline creatinine was determined as the lowest creatinine in the one year preceding the index hospital admission. If no baseline creatinine was available and no history of chronic kidney disease, we estimated the creatinine using the MDRD equation as previously described (23). We evaluated hospital mortality as well as mortality at 30 days, 90 days, and 1 year following ICU admission.

Statistical Analyses

Fisher’s Exact Test and the Wilcoxon Rank Sum Test were used for unadjusted analysis to assess the relationship between chloride load and the outcome variables. For binary outcomes, adjusted analyses were performed via logistic regression using the Hosmer-Lemeshow goodness-of-fit test for model calibration and the area under the ROC curve for model discrimination. Due to non-linear relationship between chloride load and the logit of the probability for outcomes, chloride load was analyzed categorically on the basis of quartiles. Such models were adjusted for the total volume of fluid given during LVR. Alternatively, the chloride load was divided by the total volume of LVR fluids. Time-to-event outcomes were conducted in both unadjusted and adjusted analyses. For the unadjusted analyses, Kaplan-Meier plots were assessed. Adjusted analyses were performed using the Cox proportional hazards model. Where log-log plots or Schoenfeld residual plots indicated possible violation of the proportional hazards assumption, covariates were

interacted with various transformations of time. Statistical analyses were performed using SAS (version 9.4), STATA (version SE 14), and R (3.2.2) software.

Results

Baseline Characteristics

Of the 4,710 patients in our analysis cohort (Figure 1) baseline characteristics and processes of care variables for patients with varying chloride loads are shown in Table 1. Patients who received larger chloride loads were sicker with higher APS III scores (76.0 [55.0–101.0] for quartile 4 compared to 56.0 [40.0–76.0] for quartile 1). They were more likely to be on mechanical ventilation and vasopressors and had longer ICU stays (p-values all <0.001). There was, as expected, also a relationship between chloride load and total mean volume of fluid administered. For example, the fourth quartile of chloride load received more than twice the total volume to that of the first quartile (mean of 8 liters [6.8–10; 95% CI] as compared to 3.7 liters [3.2–4.2]).

Unadjusted outcomes

The relationships (unadjusted) between chloride load and each of the outcomes examined are shown in Table 2. Crude rates of HCA, AKI and hospital mortality all increased significantly as chloride load increased ($p<0.001$). Importantly, the unadjusted association between chloride load and mortality persisted out to 1 year (our maximum follow-up) where mortality for the first (lowest) quartile of chloride load was 35% but 47% for the fourth quartile ($p<0.001$).

Chloride load and HCA

Chloride load was strongly associated with the development of HCA in univariate analysis. However, after controlling for total fluids and APS III, the relationship was attenuated and no longer significant across the quartiles ($p=0.23$). Although the omnibus comparison between the models was not significant, the odds of developing HCA increased with increasing chloride load with patients in the fourth quartile having 32% higher odds of developing HCA than patients in the first quartile ($p<0.001$) (Table 3, model A).

Chloride load and AKI

Chloride load (quartiles) was associated with increasing odds of AKI even after controlling for total fluids ($p=0.03$) (Table 3, model B). An adjusted ordinal logistic regression model yielded an odds ratio of 1.09 ($p<0.0001$) for a 100 mEq increase in chloride load. Adjusting for APS III, the odds ratio for AKI was 1.04 ($p<0.0001$). However, when chloride load was modeled as a continuous variable, it was no longer associated with AKI once controlling for APS-III.

HCA and AKI

Using ordinal logistic regression, we did find that HCA (unadjusted) was associated with AKI (OR 1.24, $p=0.025$). However, age and APS III adjustment was not performed as the proportional odds assumption was no longer valid after adding these variables ($p<0.0001$).

Therefore, the association of HCA with AKI could not be ascertained after controlling for age and APS III in multivariate analysis.

Chloride load and Mortality

The relationship between chloride load and mortality remained consistent at the various time points assessed. At 30 days following ICU admission, the odds of mortality increased dramatically with increasing chloride load with a greater than 70% increase in the odds of mortality for a patient receiving a chloride load in the fourth quartile compared to the first quartile (Table 3, model C). This relationship persisted after controlling for the total volume of fluids and baseline severity of illness (APS-III). Similar results were observed at hospital discharge and at 90 days (Table S3). The model assessing the relationship between chloride load, in quartiles, and mortality at 1-year showed good fit (Hosmer Lemeshow p -value=0.76) and the continuous model evaluating the relationship between chloride load and total volume remained significant ($p<0.001$). Chloride load remained significantly associated with increased mortality whether analyzed continuously or by quartiles and even after controlling for total fluids, age and baseline severity of illness (APS-III) ($p=0.005$; Table S3). When compared to the first quartile, the fourth quartile of chloride load had an odds ratio of 1.49 (1.16–1.92). A Cox-proportional hazards model adjusting for total volume of fluid given, age and APS III revealed a hazard ratio of 1.005 ($p=0.0015$). We also fit a separate model that included interaction terms between the model variables and time. The hazards ratios for the interaction model are summarized using a curve over time with a reference line for the hazard ratio under the proportional hazards assumption (Figure 2).

Discussion

Hyperchloremia has been identified in up to 80% of patients admitted to a mixed medical–surgical ICU (24). When large amounts of saline are infused, the kidney is slow to excrete the excess chloride load (25). In animal studies, sustained renal vasoconstriction was specifically related to hyperchloremia, which was potentiated by previous salt depletion and related to the tubular reabsorption of chloride (26, 27). Consistent with prior studies we have shown increased risk for mortality and complications (AKI) with increased use of fluids containing high chloride concentrations. However, we are the first to show that chloride load is associated with a long-term mortality hazard, at least as far as one year. In patients receiving LVR with a chloride load above the median (i.e. third and fourth quartiles), there was decreased survival compared to the first quartile. Specifically, compared to the lowest quartile of chloride load the highest quartile was 1.7 and 1.49 times as likely to die by 90 days and 1 year respectively after controlling for fluid load, age and baseline severity. When examined as a continuous variable each 100 mEq increase in chloride load was associated with 5.5% increase in mortality over one year (HR 1.055 ($p=0.0015$) again after adjusting for volume, age and severity of illness.

We have also shown that the relationship between chloride load and mortality is complex. We found a relationship between chloride load and AKI in univariate models but multivariate models that included baseline severity of illness (APS-III) did not perform acceptably. Moreover continuous chloride load was no longer significantly associated with

AKI once controlling for APS-III. Thus, AKI may not be the only, or even predominant, effect of chloride load to explain its relationship with mortality.

A similar complex relationship may exist for HCA. Importantly, the HiDenIC-8 database, which included detailed laboratory data, afforded us the opportunity to examine the contribution of HCA to clinical outcomes whereas previous studies were unable to do so (7, 9). We hypothesized that in patients receiving LVR, higher chloride load would be associated with greater incidence of HCA and this is indeed what we found. However, this association was no longer significant after controlling for total fluids administered and APS-III. Admittedly, our ascertainment of HCA was not complete because clinicians only characterized acid-base variables sufficiently to determine HCA in less than 40% of patients. Nevertheless, in those patients who had their acid-base status characterized, HCA was not linked to chloride load once we controlled for other variables. Total volume was a strong predictor of HCA and is obviously a determinant of chloride load. One could argue that we should not have attempted to control for total volume in our models since greater volumes of solutions like 0.9% saline may lead to adverse effects whereas small volumes may be well tolerated—and it would be important to understand this relationship. However, our intent was to understand whether fluid composition, rather than volume per se, was associated with harm and we did demonstrate such an association between chloride load and AKI and also with mortality. The fact that this relationship could not be shown with HCA is notable.

One possible explanation for our findings is that supra-physiologic chloride solutions may complicate existing disease but may be well-tolerated in less severely ill patients. Although LVR patients are not healthy, by definition, patients with relatively intact renal function and acid-base balance may be able to handle the chloride load whereas patients with underlying renal dysfunction or metabolic acidosis already at the start of LVR may not be. The use of saline has long been known to be associated with an increased risk of hyperchloremic metabolic acidosis (28), but it has only recently been shown that these metabolic changes can result in decreased renal blood flow and renal cortical hypoperfusion, as demonstrated in healthy volunteers (29). Yet, these volunteers could tolerate these effects and did not develop AKI.

Alternative explanations for the association between chloride load and mortality have been sought. For example, immunologic effects of metabolic acidosis have been described (30, 31), as well as lung injury (32). Acidosis has been associated with lung inflammation by increase in lung myeloperoxidase activity, and thereby, increased lung injury scores in animal models (32).

A clinical trial by Yunos et al demonstrated a decreased AKI incidence and use of RRT in ICU patients with implementation of a chloride-restricted strategy (10). However, this study did not explore the relationship between high chloride load and development of HCA. In the SPLIT trial, a cluster randomized double-crossover trial, no difference in outcomes were seen using saline vs buffered crystalloid solution (plasmalyte) (11). However, in this study, the mean crystalloid infusion was around 2 liters which is considerably lower than our starting point of LVR that included 60 ml/kg over a 24 hour period. The authors do suggest that the confidence intervals around the point estimate of treatment effect were wide and the

possibility of a clinically significant effect of AKI was not excluded. In addition, the patients in this trial were of lower acuity than our LVR population. Higher risk patients may still benefit from avoiding chloride-rich fluids based on our results. A meta-analysis of 21 studies that included randomized clinical trials and observational studies found a significantly higher risk of AKI (RR 1.64, 95% CI 1.27 to 2.13; $P < 0.001$) and hyperchloremia/metabolic acidosis (RR 2.87, 1.95 to 4.21; $P < 0.001$) but no mortality effect (33).

Our study has important strengths. Most importantly, our ascertainment of exposure (chloride load) and clinical outcomes including HCA and AKI are based on primary data (documented infusion of fluid, laboratory volumes and urine volume) not on billing or other administrative codes. We carefully determined fluid composition and characterized acid-base disorders (rather than relying only on hyperchloremia) to a level of detail not performed (to our knowledge) in any previous study on this topic. Similarly, we ascertained AKI using changes in both creatinine and urine output using the latest international guidelines (22). We have previously shown that failure to include urine output criteria significantly underestimates AKI (34). Finally, we were able to link our data with national databases to determine long-term hazards.

Our study also has important limitations. We did not include patients receiving LVR in the OR since we could not ascertain fluid administration to the same level of detail nor could we adequately account for blood loss. In nearly two-thirds of patients receiving LVR we could not determine if HCA occurred because clinicians did not characterize their acid-base status. We focused on patients receiving at least 60ml/kg in a 24 hour period because this is where we would anticipate the most effects of chloride load given a large exposure and a susceptible population. However, it is possible that some effect might be observed with smaller volumes in less critically ill patients. Our results cannot address this question nor can we state what the minimal exposure to supra-physiologic chloride solutions would be in order to observe any hazard. Our methods do not address what the optimal alternative fluid to saline would be. Our patients received a variety of alternative fluids and we did not attempt to analyze differences among these fluids (Table S4). Our database is now roughly 10–15 years old and practice has changed in this time. However we do not think that fluid resuscitation has been relatively unchanged, at least in North America where saline remains the preferred fluid. Finally, because this was not a randomized trial we cannot entirely control for bias. Patients with higher illness severity received more fluids and thus more chloride. However, the mortality signal remained after controlling for both baseline severity (APS III) and total fluid volume.

Conclusions

Among patients receiving large volume resuscitation (>60 ml/kg over a 24 hours), increasing chloride load is associated with significant adverse effects on survival out to one year even after controlling for total fluid administered and baseline severity of illness. These results add to a growing body of literature suggesting that large volumes of high-chloride solutions may be deleterious. However, the mechanisms responsible for these effects remain unclear. We recommend avoiding supra-physiologic chloride solutions when resuscitation volumes are likely to exceed 60 ml/kg.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by funds from Baxter Healthcare Corporation. Baxter reviewed the content of this article, but the data collection, data analysis and interpretation, and writing are solely the responsibility of the authors.

References

- Berger MM, Pictet A, Revelly JP, et al. Impact of a bicarbonated saline solution on early resuscitation after major burns. *Intensive Care Med.* 2000; 26(9):1382–1385. [PubMed: 11089771]
- Cho YS, Lim H, Kim SH. Comparison of lactated Ringer's solution and 0.9% saline in the treatment of rhabdomyolysis induced by doxylamine intoxication. *Emerg Med J.* 2007; 24(4):276–280. [PubMed: 17384382]
- Cieza JA, Hinostroza J, Huapaya JA, et al. Sodium chloride 0.9% versus Lactated Ringer in the management of severely dehydrated patients with choleraform diarrhoea. *J Infect Dev Ctries.* 2013; 7(7):528–532. [PubMed: 23857387]
- Mahler SA, Conrad SA, Wang H, et al. Resuscitation with balanced electrolyte solution prevents hyperchloremic metabolic acidosis in patients with diabetic ketoacidosis. *Am J Emerg Med.* 2011; 29(6):670–674. [PubMed: 20825879]
- Raghunathan K, Shaw A, Nathanson B, et al. Association between the choice of IV crystalloid and in-hospital mortality among critically ill adults with sepsis*. *Crit Care Med.* 2014; 42(7):1585–1591. [PubMed: 24674927]
- Yunos NM, Bellomo R, Hegarty C, et al. Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *Jama.* 2012; 308(15):1566–1572. [PubMed: 23073953]
- McCluskey SA, Karkouti K, Wijesundera D, et al. Hyperchloremia after noncardiac surgery is independently associated with increased morbidity and mortality: a propensity-matched cohort study. *Anesth Analg.* 2013; 117(2):412–421. [PubMed: 23757473]
- Boniatti MM, Cardoso PR, Castilho RK, et al. Is hyperchloremia associated with mortality in critically ill patients? A prospective cohort study. *J Crit Care.* 2011; 26(2):175–179. [PubMed: 20619601]
- Yunos NM, Kim IB, Bellomo R, et al. The biochemical effects of restricting chloride-rich fluids in intensive care. *Crit Care Med.* 2011; 39(11):2419–2424. [PubMed: 21705897]
- Yunos NM, Bellomo R, Glassford N, et al. Chloride-liberal vs. chloride-restrictive intravenous fluid administration and acute kidney injury: an extended analysis. *Intensive Care Med.* 2015; 41(2):257–264. [PubMed: 25518951]
- Young P, Bailey M, Beasley R, et al. Effect of a Buffered Crystalloid Solution vs Saline on Acute Kidney Injury Among Patients in the Intensive Care Unit: The SPLIT Randomized Clinical Trial. *Jama.* 2015; 314(16):1701–1710. [PubMed: 26444692]
- Kellum JA, Song M, Almasri E. Hyperchloremic acidosis increases circulating inflammatory molecules in experimental sepsis. *Chest.* 2006; 130(4):962–967. [PubMed: 17035425]
- Zhou F, Peng ZY, Bishop JV, et al. Effects of fluid resuscitation with 0.9% saline versus a balanced electrolyte solution on acute kidney injury in a rat model of sepsis*. *Crit Care Med.* 2014; 42(4):e270–278. [PubMed: 24335444]
- Wan L, Bellomo R, May CN. The effect of normal saline resuscitation on vital organ blood flow in septic sheep. *Intensive Care Med.* 2006; 32(8):1238–1242. [PubMed: 16775719]
- Kiraly LN, Differding JA, Enomoto TM, et al. Resuscitation with normal saline (NS) vs. lactated ringers (LR) modulates hypercoagulability and leads to increased blood loss in an uncontrolled hemorrhagic shock swine model. *J Trauma.* 2006; 61(1):57–64. discussion 64–55. [PubMed: 16832250]

16. Waters JH, Gottlieb A, Schoenwald P, et al. Normal saline versus lactated Ringer's solution for intraoperative fluid management in patients undergoing abdominal aortic aneurysm repair: an outcome study. *Anesth Analg.* 2001; 93(4):817–822. [PubMed: 11574339]
17. Williams EL, Hildebrand KL, McCormick SA, et al. The effect of intravenous lactated Ringer's solution versus 0.9% sodium chloride solution on serum osmolality in human volunteers. *Anesth Analg.* 1999; 88(5):999–1003. [PubMed: 10320158]
18. Shaw AD, Bagshaw SM, Goldstein SL, et al. Major complications, mortality, and resource utilization after open abdominal surgery: 0.9% saline compared to Plasma-Lyte. *Ann Surg.* 2012; 255(5):821–829. [PubMed: 22470070]
19. Young JB, Utter GH, Schermer CR, et al. Saline versus Plasma-Lyte A in initial resuscitation of trauma patients: a randomized trial. *Ann Surg.* 2014; 259(2):255–262. [PubMed: 23732264]
20. Knaus WA, Wagner DP, Draper EA, et al. The APACHE III prognostic system. Risk prediction of hospital mortality for critically ill hospitalized adults. *Chest.* 1991; 100(6):1619–1636. [PubMed: 1959406]
21. Sileanu FE, Murugan R, Lucko N, et al. AKI in low-risk versus high-risk patients in intensive care. *Clin J Am Soc Nephrol.* 2015; 10(2):187–196. [PubMed: 25424992]
22. KDIGO AW. Kidney Disease: Improving Global Outcomes (KDIGO) Clinical Practice Guideline for Acute Kidney Injury. *Kidney Int Suppl* (2011). 2012; 2(1):1–141.
23. Zavada J, Hoste E, Cartin-Ceba R, et al. A comparison of three methods to estimate baseline creatinine for RIFLE classification. *Nephrol Dial Transplant.* 2010; 25(12):3911–3918. [PubMed: 20100732]
24. Moviat M, van Haren F, van der Hoeven H. Conventional or physicochemical approach in intensive care unit patients with metabolic acidosis. *Crit Care.* 2003; 7(3):R41–45. [PubMed: 12793889]
25. Veech RL. The toxic impact of parenteral solutions on the metabolism of cells: a hypothesis for physiological parenteral therapy. *Am J Clin Nutr.* 1986; 44(4):519–551. [PubMed: 3094358]
26. Wilcox CS. Regulation of renal blood flow by plasma chloride. *J Clin Invest.* 1983; 71(3):726–735. [PubMed: 6826732]
27. Matavelli LC, Zhou X, Varagic J, et al. Salt loading produces severe renal hemodynamic dysfunction independent of arterial pressure in spontaneously hypertensive rats. *Am J Physiol Heart Circ Physiol.* 2007; 292(2):H814–819. [PubMed: 16997890]
28. Kellum JA. Saline-induced hyperchloremic metabolic acidosis. *Crit Care Med.* 2002; 30(1):259–261. [PubMed: 11902280]
29. Chowdhury AH, Cox EF, Francis ST, et al. A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and plasma-lyte(R) 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg.* 2012; 256(1):18–24. [PubMed: 22580944]
30. Rhee P, Wang D, Ruff P, et al. Human neutrophil activation and increased adhesion by various resuscitation fluids. *Crit Care Med.* 2000; 28(1):74–78. [PubMed: 10667502]
31. Junger WG, Hoyt DB, Hamreus M, et al. Hypertonic saline activates protein tyrosine kinases and mitogen-activated protein kinase p38 in T-cells. *J Trauma.* 1997; 42(3):437–443. discussion 443–445. [PubMed: 9095111]
32. Pedoto A, Caruso JE, Nandi J, et al. Acidosis stimulates nitric oxide production and lung damage in rats. *Am J Respir Crit Care Med.* 1999; 159(2):397–402. [PubMed: 9927349]
33. Krajewski ML, Raghunathan K, Paluszkiwicz SM, et al. Meta-analysis of high- versus low-chloride content in perioperative and critical care fluid resuscitation. *Br J Surg.* 2015; 102(1):24–36. [PubMed: 25357011]
34. Kellum JA, Sileanu FE, Murugan R, et al. Classifying AKI by Urine Output versus Serum Creatinine Level. *J Am Soc Nephrol.* 2015; 26(9):2231–2238. [PubMed: 25568178]

Figure 1.
Study Cohort Selection

Figure 2. Hazard ratio plot

This figure displays the hazard ratio for a 100 unit increase in chloride load for a Cox proportional hazards model adjusting for total volume of fluid given, age, and APACHE-III (dotted line), as well as the hazard ratio at various days when an interaction term is incorporated to relax the proportional hazards assumption (solid curve) and the same set of adjusting covariates. The simple proportional hazard model has a hazard ratio of 1.055 (p-value = 0.0015). The joint significance of the main effect for chloride load and the interaction term between chloride load and log time has a p-value of 0.0016.

Table 1

Baseline Characteristics and Process of Care by Chloride Load (N=4,710)

Characteristics	Q1	Q2	Q3	Q4	p-value
mEq	323–491	492–635	636–848	849–5432	
Age, years					
18 to 44	316 (26.9%)	287 (24.4%)	249 (21.1%)	300 (25.5%)	
45 to < 65	443 (37.7%)	487 (41.3%)	504 (42.8%)	516 (43.9%)	<0.001
65 to < 75	169 (14.4%)	185 (15.7%)	204 (17.3%)	191 (16.2%)	
75	248 (21.1%)	219 (18.6%)	221 (18.8%)	169 (14.4%)	
Males	422 (35.9%)	559 (47.5%)	640 (54.3%)	729 (61.9%)	<0.001
Race					
White	921 (78.2%)	920 (78.1%)	933 (79.2%)	868 (73.7%)	
Black	94 (8.0%)	88 (7.5%)	86 (7.3%)	104 (8.8%)	0.05
Other	162 (13.8%)	170 (14.4%)	159 (13.5%)	205 (17.4%)	
BMI, Median (IQR)	20.8 (18.5–23.6)	23.5 (20.8 – 26.1)	25.1 (22.5 – 28.3)	26.6 (23.0 – 30.8)	<0.001
APS III^{a,b}, Median (IQR)	56.0 (40.0 – 76.0)	62.0 (44.0 – 81.0)	64.0 (45.0 – 83.0)	76.0 (55.0 – 101.0)	<0.001
Comorbid condition					
Diabetes	137 (11.6%)	166 (14.1%)	186 (15.8%)	174 (14.8%)	0.028
Cardiac Disease	149 (12.7%)	150 (12.7%)	134 (11.4%)	143 (12.1%)	0.73
Chronic Renal Disease	61 (5.2%)	61 (5.2%)	66 (5.6%)	45 (3.8%)	0.21
Vascular Disease	96 (8.2%)	91 (7.7%)	104 (8.8%)	74 (6.3%)	0.13
Multiple Comorbidity	605 (51.4%)	586 (49.7%)	552 (46.9%)	516 (43.8%)	0.001
Nephrotoxins^{a,c}	560 (47.6%)	545 (46.3%)	558 (47.4%)	538 (45.7%)	0.77
Surgical admission	653 (59.1%)	670 (61.3%)	633 (58.2%)	625 (58.1%)	0.40
Mechanical ventilation^a	784 (66.6%)	842 (71.5%)	860 (73.0%)	953 (81.0%)	<0.001
Serum creatinine, mg/dl, Median (IQR)					
Baseline Creatinine	0.8 (0.6 – 1.2)	0.9 (0.7 – 1.2)	0.9 (0.7 – 1.2)	0.9 (0.7 – 1.1)	0.022
Admission Creatinine	0.9 (0.7 – 1.2)	0.9 (0.7 – 1.3)	1.0 (0.8 – 1.4)	1.1 (0.8 – 1.7)	<0.001
Reference Creatinine	0.8 (0.6 – 1.0)	0.8 (0.7 – 1.0)	0.9 (0.7 – 1.0)	0.9 (0.7 – 1.1)	<0.001
Hypotensive index^{a,d} > 0	431 (36.6%)	573 (48.7%)	645 (54.8%)	825 (70.5%)	<0.001
Fluids in Liters, Median (IQR)	3.7 (3.2 – 4.2)	4.4 (3.9 – 5.0)	5.4 (4.8 – 6.0)	8.0 (6.8 – 10.0)	<0.001

Characteristics	Q1	Q2	Q3	Q4	p-value
mEq	323–491	492–635	636–848	849–5432	
Vasopressors ^a	290 (24.6%)	383 (32.5%)	485 (41.2%)	695 (59.0%)	<0.001
Hospital Length of Stay, days	15.0 (9.0 – 26.0)	15.5 (8.0 – 28.0)	16.0 (8.0 – 27.0)	16.0 (7.0 – 28.0)	0.45
ICU Length of Stay, days	6.0 (3.0 – 11.0)	6.0 (4.0 – 14.0)	7.0 (4.0 – 13.0)	8.0 (4.0 – 16.0)	<0.001
Sepsis ^{a,e}	180 (15.3%)	222 (18.8%)	284 (24.1%)	382 (32.5%)	<0.001

^aMeasured within 24 hours of ICU admission

^bAPS-III, Acute Physiology Score

^cNephrotoxins include angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, vancomycin, aminoglycoside, calcineurin inhibitor, nonsteroidal, diuretics

^dArea under the curve for severity and duration of hypotension (see the Materials and Methods)

^e“Suspected sepsis” defined as ordering of blood cultures and antibiotics (see the Materials and Methods)

Table 2

Outcomes stratified by chloride load

Quartile	Q1	Q2	Q3	Q4	P-value
mEq	323–491	492–635	636–848	849–5432	
HCA within 24 hours of LVR					
Missing	758 (64.4%)	748 (63.5%)	737 (62.6%)	651 (55.3%)	
No	335 (28.5%)	341 (28.9%)	321 (27.2%)	296 (25.1%)	<0.001
Yes	84 (7.1%)	89 (7.6%)	120 (10.2%)	230 (19.5%)	
Max KDIGO					
No AKI	329 (27.9%)	245 (20.8%)	226 (19.2%)	163 (13.9%)	
Stage 1	221 (18.8%)	211 (17.9%)	165 (14.0%)	151 (12.8%)	
Stage 2	371 (31.5%)	421 (35.7%)	433 (36.8%)	362 (30.8%)	<0.001
Stage 3	256 (21.7%)	301 (25.5%)	354 (30.1%)	499 (42.5%)	
Hospital Mortality	226 (19.2%)	285 (24.2%)	298 (25.3%)	417 (35.4%)	<0.001
Mortality after ICU Admission					
30 Days	207 (17.6%)	266 (22.6%)	289 (24.5%)	387 (32.9%)	<0.001
90 Days	294 (24.9%)	343 (29.1%)	374 (31.7%)	480 (40.8%)	<0.001
365 Days	407 (34.6%)	432 (36.7%)	481 (40.8%)	553 (46.9%)	<0.001

Table 3

Logistic regression models for chloride load quartiles and clinical outcomes

A. Hyperchloremic acidosis (N=2,526)*		
Effect	Odds Ratio	p-value[§]
Chloride load Q2	0.88 (0.63,1.22)	
Chloride load Q3	1.10 (0.78,1.53)	0.23
Chloride load Q4	1.32 (0.86,2.01)	
Fluids in Liters	1.15 (1.08,1.22)	<.0001
Age	1.00 (1.00,1.01)	0.27
APS-III	1.02 (1.02,1.03)	<.0001
B. Acute kidney injury (N=4,706)**		
Chloride load Q2	1.30 (1.06,1.59)	
Chloride load Q3	1.33 (1.07,1.66)	0.03
Chloride load Q4	1.41 (1.02,1.96)	
Fluids in Liters	1.04 (0.98,1.09)	0.18
Age	1.03 (1.02,1.03)	<.0001
APS-III	1.03 (1.02,1.03)	<.0001
C. Mortality by 30 days (N=4,708)***		
Chloride load Q2	1.26 (1.02,1.56)	
Chloride load Q3	1.35 (1.09,1.69)	0.002
Chloride load Q4	1.71 (1.30,2.26)	
Fluids in Liters	1.00 (0.97,1.04)	0.95
Age	1.03 (1.03,1.04)	<.0001
APS-III	1.02 (1.02,1.02)	<.0001

* AUC = 0.8212, Hosmer-Lemeshow goodness of fit test p-value=0.022;

** AUC = 0.7581, Hosmer-Lemeshow goodness of fit test p-value = 0.4105;

*** AUC = 0.7260, Hosmer-Lemeshow goodness of fit test p-value = 0.8277;

[§]P-values for effect of chloride load reflect the omnibus test across all 4 quartiles (comparator is always quartile 1)