Published in final edited form as:

Org Biomol Chem. 2018 August 22; 16(33): 5979–5986. doi:10.1039/c8ob01529e.

Enantioselective Total Synthesis of Decytospolide A and Decytospolide B Using an Achmatowicz Reaction

Arun K. Ghosh, Hannah M. Simpson, and Anne M. Veitschegger

Department of Chemistry and Department of Medicinal Chemistry, Purdue University, 560 Oval Drive, West Lafayette, Indiana 47907, United States. akghosh@purdue.edu

Abstract

Enantioselective syntheses of decytospolide A and decytospolide B are described here. The current synthesis highlights an Achmatowicz rearrangement of an optically active furanyl alcohol followed by reduction of the resulting dihydropyranone hemiacetal with $\text{BF}_3 \cdot \text{OEt}_2$ and Et_3SiH to provide the saturated tetrahydropyranyl alcohol directly. This reduction was investigated with a variety of other Lewis acids. The synthesis also features Noyori asymmetric transfer hydrogenation and Friedel-Craft acylation. Overall, the synthesis provides ready access to the natural products and may be useful in the preparation of bioactive derivatives.

Abstract

Decytospolide A (R = H)

Decytospolide B (R = Ac)

Enantioselective syntheses of decytospolides are described using an Achmatowicz rearrangement, transfer hydrogenation and Friedel-Craft acylation as the key steps.

Introduction

Functionalized tetrahydropyrans are important structural features present in many bioactive natural products.^{1,2} Over the years, several methods have been developed for the synthesis of substituted tetrahydropyran rings.^{3,4,5} However, there are limitations with respect to readily available starting materials, stereo- and regiochemical issues and lack of potential for incorporation of multiple substitutions within the tetrahydropyran ring system. Additionally, many of these transformations rely on transition metals such as a Pd-catalyzed

Correspondence to: Arun K. Ghosh.

† Electronic Supplementary Information (ESI) available: 1H- and 13C-NMR spectra of new compounds. See DOI: 10.1039/x0xx00000x

decarboxylative allylation and an indium catalysed Prins cyclization.^{6,7} The Achmatowicz reaction, an oxidative ring enlargement of a furanyl alcohol, has been developed into a very practical reaction with immense potential.⁸ In recent years, the Achmatowicz reaction has been utilized in the synthesis of a variety of natural products.^{10,11} We have utilized this reaction in the synthesis of a number of bioactive natural products containing functionalized tetrahydropyran rings.^{10,12,13} In particular, the Achmatowicz reaction of furanyl alcohol **1** (Figure 1) provides dihydropyranone hemiacetal **2** which upon reduction, typically with triethylsilane in the presence of trifluoroacetic acid (TFA) provided a variety of 2,6-disubstituted dihydropyranone derivatives **3**.^{10,11} Such enones have been utilized in the synthesis of bioactive natural products, including the potent anticancer agent, herboxidiene **4**.^{10,13}

Thus far, dihydropyranone hemiacetal reduction provides access to a range of enones under a variety of reaction conditions.^{10,11} However, the potential for reduction of an Achmatowicz reaction product enone hemiacetal to a saturated tetrahydropyran derivative and further reduction to the alcohol functionality has been scarcely explored. For further development of the Achmatowicz reaction as well as its application, we sought to synthesize 2,6-disubstituted tetrahydropyranyl alcohols as exemplified by alcohol **5** using a silane in the presence of a Lewis acid. There have been limited studies for this transformation.¹⁴ This transformation would provide easy access to natural products containing substituted tetrahydropyran rings with three contiguous chiral centers. Such functionalized tetrahydropyran rings are imbedded in a variety of bioactive molecules, including decytospolides A (**6**) and B (**7**) and their derivatives.

The decytospolides contain three asymmetric centers surrounding a central tetrahydropyran ring flanked by two alkyl chains. Both natural products were recently isolated by Zhang and co-workers from the endophytic fungus, *Cytospora* sp. No ZW02, from *Ilex canariensis*, an evergreen shrub from the Canary Islands.¹⁵ The chemical structure of both decytospolides was determined by extensive NMR studies and HRMS analysis. The absolute configuration was established through Mosher ester analysis.^{15,16} Decytospolide B exhibited moderate cytotoxicity in A549 and QGY cancer cell lines with IC₅₀ values of 14.8 and 46.8 µg/mL, respectively. Since decytospolide A did not show appreciable cytotoxicity, the acyl group in decytospolide B may be responsible for its moderate anticancer activity. There is potential for further improvement through modification of this acyl group. Several syntheses of these natural products have been reported.^{6,7,17–19} We, however, planned to assemble the functionalized tetrahydropyran ring using the Achmatowicz reaction as the key step. Herein, we report our synthesis of decytospolides A and B using the Achmatowicz, Noyori reduction, and Friedel-Craft reactions as the key steps. We particularly sought to synthesize 2,5,6-trisubstituted tetrahydropyran derivatives in a highly stereoselective manner in optically active form in a one-pot operation directly from the Achmatowicz product, the dihydropyranone hemiacetal.

Results and discussion

Our synthetic strategies to decytospolides A and B is shown in Scheme 1. We planned to synthesize trisubstituted tetrahydropyran alcohol **8** in an optically active manner by

reduction of dihydropyran hemiacetal **9** which would be obtained directly from furanyl alcohol **10** using an Achmatowicz rearrangement. We particularly planned to synthesize alcohol **8** stereoselectively from dihydropyran hemiacetal **9**.^{10,11} Furan derivative **10** would be synthesized from furan **11** by a Friedel-Crafts acylation followed by asymmetric reduction.²⁰

The synthesis of furanyl alcohol **10** in optically active form is shown in Scheme 2. Deprotonation of furan **11** with *n*-BuLi in THF at 0 °C to 23 °C followed by addition of commercially available (*R*)-butylene oxide furnished the corresponding alcohol through epoxide opening. The resulting alcohol was acetylated with acetic anhydride in the presence of Et₃N and DMAP to afford furan derivative **12** in 84% yield over two steps. We specifically planned to install the C3-acetoxy group with defined stereochemistry to avoid forming a mixture of diastereomers and provide easy access to C3 stereo-defined derivatives for biological evaluation. Furan derivative **12** underwent a Friedel-Craft acylation by reaction with hexanoyl chloride in the presence of SnCl₄ at 0 °C for 30 min to provide ketone derivative **13** in 75% yield.²¹ For enantioselective reduction of the ketone, we planned an asymmetric transfer hydrogenation reaction developed by Noyori and co-workers.^{22,23} Therefore, reaction of **13** with a catalytic (1 mol%) amount of Noyori's catalyst, (*R,R*) RuCl(mesitylene)-Ts-DPEN, in the presence of Et₃N and formic acid in CH₂Cl₂ at 55 °C for 12 h, furnished alcohol **10** in 93% yield. The asymmetric reduction proceeded with high diastereoselectivity as alcohol **10** was isolated as the single product by ¹H- and ¹³C-NMR analysis (diastereoselectivity > 20:1).

The synthesis of the decytospolides is shown in Scheme 3. Initially, the Achmatowicz reaction of **10** was carried out with a catalytic amount of VO(acac)₂ and ^tBuOOH; however, the reaction was sluggish.^{24,25} Achmatowicz reaction with oxone in the presence of NaHCO₃ in a mixture (4:1) of THF and water at 23 °C for 30 min smoothly converted **10** to the corresponding dihydropyranone hemiacetal (**9**).²⁶ The resulting hemiacetal was initially subjected to reduction with Et₃SiH and TFA at -45 °C for 3 h. This resulted in a mixture of dihydropyranone **14** and a small amount of alcohol **8**. Further optimization of the reaction with an excess of BF₃•OEt₂ at -45 °C for 3 h provided alcohol **8** as the exclusive product in 47% yield over 2-steps. The stereochemistry at C8 was determined by ¹H NMR coupling with the proton at C9, which has two J values (2.4 and 9 Hz). Having J = 9 Hz indicates a trans relationship between the protons at C8 and C9 and since the stereocenter at C9 was set by the Noyori reduction, the stereochemistry at C8 was assigned based on that. We then, investigated silane reduction of the dihydropyranone hemiacetal **9** in the presence of a number of other Lewis acids. The results are shown in Table 1. As can be seen, the use of BF₃•OEt₂ and SnCl₄ as the Lewis acids provided alcohol **8** exclusively in 56% and 82% yield, respectively (entries 3 and 4). Reductions with Lewis acids Sc(OTf)₃ and Cu(OTf)₂ yielded only trace amounts of dihydropyranone **14**, while using TiCl₄ as the Lewis acid gave 14% of **14**.

To complete the synthesis of the decytospolides, alcohol **8** was protected as a TBS-ether with TBSOTf in the presence of 2,6-lutidine in CH₂Cl₂ at 0 °C to 23 °C for 12 h. The acetate was hydrolyzed with K₂CO₃ in MeOH at 23 °C to provide the corresponding

alcohol. Oxidation of the resulting alcohol with Dess-Martin periodinane in the presence of NaHCO_3 in CH_2Cl_2 furnished ketone **15** in 89% yield over 3-steps. Removal of the silyl ether was carried out with tetrabutylammonium fluoride (TBAF) in THF at 23 °C for 12 h to provide decytospolide A (**6**) $\{[\alpha]_D^{23}+18.4$ (c 0.45, CHCl_3) $\}$ in quantitative yield. Treatment of decytospolide A (**6**) with acetic anhydride in the presence of pyridine and DMAP in CH_2Cl_2 at 23 °C for 1.5 h furnished decytospolide B (**7**) $\{[\alpha]_D^{23}+28.4$ (c 0.75, CHCl_3) $\}$ in quantitative yield. The ^1H - and ^{13}C -NMR spectra of synthetic decytospolides are in complete agreement with the spectra reported for the natural decytospolide A $\{[\alpha]_D^{20}+6.1$ (c 0.08, CHCl_3) $\}$ and decytospolide B $\{[\alpha]_D^{20}+26.6$ (c 0.02, CHCl_3) $\}$.¹⁴

We also investigated an alternative route to the decytospolides in an effort to synthesize the saturated tetrahydropyranol directly following the Achmatowicz reaction. As shown in Scheme 4, commercially available furural, was converted to furanyl methyl acetate **17** by a one-carbon homologation using a Jovic reaction as developed by Snowden and co-workers.^{27,28} Methyl ester **17** is also commercially available. Methyl ester **17** was reacted with hexanoyl chloride in the presence of SnCl_4 as described above to provide furanyl-ketone **18** in 57% yield. Reduction of ketone **18** using Noyori's catalyst $\text{RuCl}(\text{mesitylene})[(R,R)\text{-Ts-DPEN}]$ (1 mol%) in the presence of Et_3N and formic acid at 50 °C for 12 h afforded optically active furanyl alcohol **19** in 83% yield and 99% ee determined by chiral HPLC analysis (Please see experimental section for details).^{20,21} Achmatowicz reaction of **19** with oxone at 23 °C for 30 min furnished the corresponding dihydropyranone hemiacetal which was subjected to reduction with excess Et_3SiH in the presence of $\text{BF}_3\cdot\text{OEt}_2$ at -40 °C for 16 h to provide saturated tetrahydropyranol derivative **20** as a single product by ^1H -NMR analysis. Presumably, the reduction of enone first provided the ketone which was reduced by Lewis acid chelation followed by axial delivery of hydride.

Tetrahydropyranol derivative **20** was readily converted to the decytospolides as shown in Scheme 5. Reaction of methyl ester **20** with $\text{NH}(\text{OMe})\text{Me}\cdot\text{HCl}$ in the presence of $i\text{-PrMgCl}$ in THF at -30 °C for 5 h provided Weinreb amide derivative **21**. Treatment of Weinreb amide **21** with EtMgBr in THF at 0 °C to 23 °C for 5 h afforded decytospolide A, **6** in 97% yield $\{[(\alpha)_D^{23}+7.8$ (c , 1.33, CHCl_3) $\}]$. Acylation of **6** with acetic anhydride in the presence of pyridine and DMAP furnished decytospolide B, **7** $\{[\alpha]_D^{23}+22.8$ (c , 1.96, CHCl_3) $\}$ in 86% yield. The ^1H -NMR and ^{13}C -NMR spectra of these synthetic decytospolides are in complete agreement with reported spectra for the natural products.¹⁴

Conclusions

In summary, we have accomplished an enantioselective total syntheses of decytospolides A and B. The synthesis features an Achmatowicz rearrangement of an optically active furanyl alcohol which was obtained conveniently by use of the Friedel-Craft reaction followed by a Noyori asymmetric transfer hydrogenation reaction as the key steps. The synthesis highlights a highly stereoselective reduction of the Achmatowicz product, a dihydropyranone hemiacetal to the saturated tetrahydropyranol derivative using $\text{BF}_3\cdot\text{OEt}_2$ and Et_3SiH . Reduction presumably proceeds through Lewis Acid chelation followed by delivery of an axial hydride. The current work will provide access to structural variants of

these natural products for further studies. Further studies and applications are in progress in our laboratories.

Experimental Section

Chemicals and reagents were purchased from commercial suppliers and used without further purification. Anhydrous solvents were obtained as follows: dichloromethane and toluene from calcium hydride, diethyl ether and tetrahydrofuran from sodium/benzophenone, and methanol from activated magnesium. All other solvents were reagent grade. All moisture-sensitive reactions were either carried out in flame- or oven-dried (120 °C) glassware under an argon atmosphere. TLC analysis was conducted using glass-backed thin-layer silica gel chromatography plates (60 Å, 250 µm thickness, F254 indicator). Column chromatography was performed using Silicycle 230–400 mesh, 60 Å pore diameter silica gel. ^1H and ^{13}C NMR spectra were recorded on either Bruker ARX400, Bruker DRX-500, Bruker AV500HD, or Bruker Avance-III-800 spectrometers. Chemical shift (δ values) are reported in parts per million and are referenced to the residual solvent signal (CDCl_3 ^1H singlet = 7.26, ^{13}C triplet = 77.16). Characteristic splitting patterns due to spin-spin coupling are identified as follows: s = singlet, d = doublet, t = triplet, q = quartet, quint = quintet, sept = septet, m = multiplet, dd = doublet of doublets, ddd = doublet of doublet of doublets, td = triplet of doublets, dq = doublet of quartets, brs = broad singlet, app = apparent. All coupling constants are measured in hertz (Hz). Optical rotations were recorded by a Perkin Elmer 341 polarimeter. IR spectra were recorded on a Perkin Elmer Spectrum Two FT-IR Spectrometer. LRMS and HRMS spectra were recorded at the Purdue University Department of Chemistry Mass Spectrometry Center. HPLC data was obtained on an Agilent 1290 Infinity II.

(*R*)-1-(Furan-2-yl)butan-2-yl acetate (**12**):

To furan (2.02 mL, 27.74 mmol) dissolved in THF (23 mL) at 0 °C was added *n*-BuLi (15.6 mL, 24.97 mmol) dropwise upon which a bright yellow color developed. After stirring for 1 h at this temperature, (*R*)-(+)-butylene oxide (1.21 mL, 13.87 mmol) was added and the reaction was slowly warmed to room temperature. After stirring for 12 h, the deep red solution was quenched with satd. NH_4Cl , extracted with EtOAc, washed with brine and dried over Na_2SO_4 . Purification by column chromatography (10% EtOAc/Hexanes) gave 1.69 g (87% yield) of the resulting furan alcohol as a yellow oil.

To (*R*)-furan alcohol (868 mg, 6.19 mmol) dissolved in CH_2Cl_2 (21 mL) at 0 °C was added acetic anhydride (1.2 mL, 12.4 mmol), Et_3N (1.3 mL, 9.3 mmol) and a few crystals of DMAP. The reaction was allowed to warm to room temperature. After 6 h, the reaction was quenched with satd. NaHCO_3 , extracted with CH_2Cl_2 , washed with brine and dried over Na_2SO_4 . Purification by column chromatography (5% to 10% EtOAc/Hexanes) afforded 1.1 g (97% yield) of acetate **12** as a clear oil $[\alpha]_{\text{D}}^{20} +19.6$ (c 0.73, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 7.31 (dd, J = 1.9, 0.9 Hz, 1H), 6.28 (dd, J = 3.2, 1.9 Hz, 1H), 6.05 (dd, J = 3.2, 0.9 Hz, 1H), 5.04 (m, 1H), 2.87 (d, J = 6.3 Hz, 2H), 2.02 (s, 3H), 1.65–1.55 (m, 2H), 0.92 (t, J = 7.5 Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 170.7, 151.9, 141.5, 110.4, 107.0, 73.8, 32.5, 26.7, 21.3, 9.7; FT-IR (neat) ν_{max} = 2965, 2925, 2852, 1738, 1507, 1461, 1436, 1376, 1239, 1012, 739 cm^{-1}

(R)-1-(5-Hexanoylfuran-2-yl)butan-2-yl acetate (13):

Hexanoic acid (604 mg, 5.20 mmol) in an excess of thionyl chloride was refluxed overnight. The thionyl chloride was removed by distillation, strictly keeping the system under argon. The remaining hexanoyl chloride was used immediately for the subsequent reaction. To hexanoyl chloride (699 mg, 5.2 mmol) in CH₂Cl₂ (15 mL) at 0 °C was added SnCl₄ (8.7 mL, 8.7 mmol, 1 M in CH₂Cl₂) dropwise. After stirring for 1 h at this temperature, acetate **12** (789 mg, 4.33 mmol) dissolved in CH₂Cl₂ (5 mL) was added to the reaction mixture via cannula and remained stirring at 0 °C. After 30 min, the red-brown solution was quenched with ice, extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (10% to 20% EtOAc/Hexanes) gave 909 mg (75% yield) of furan derivative **13** as a clear oil; [α]_D²⁰ +16.2 (*c* 0.86, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 7.07 (d, *J* = 3.5 Hz, 1H), 6.21 (d, *J* = 3.5 Hz, 1H), 5.07 (quint, *J* = 6.4 Hz, 1H), 2.99–2.91 (m, 2H), 2.74 (t, *J* = 7.9 Hz, 2H), 2.02 (s, 3H), 1.72–1.66 (m, 2H), 1.65–1.59 (m, 2H), 1.34–1.31 (m, 4H), 0.92 (t, *J* = 7.4 Hz, 3H), 0.89 (t, *J* = 6.5 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 189.4, 170.6, 157.0, 152.1, 118.3, 109.9, 73.2, 38.4, 32.8, 31.6, 26.9, 24.3, 22.6, 21.2, 14.0, 9.7; FT-IR (neat) ν_{max} = 2961, 2931, 2866, 1741, 1674, 1588, 1516, 1374, 1238, 1023 cm⁻¹

(R)-1-(5-((R)-1-Hydroxyhexyl)furan-2-yl)butan-2-yl acetate (10):

To furan derivative **13** (280 mg, 1 mmol) in CH₂Cl₂ (5 mL) was sequentially added Et₃N (1.0 mL, 7.5 mmol), formic acid (0.28 mL, 7.5 mmol) and Noyori catalyst, (*R,R*) RuCl(mesitylene)-Ts-DPEN (6.2 mg, 0.01 mmol), and the reaction was set to reflux at 55 °C. After refluxing for 12 h, the orange solution was diluted with water, extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (10% to 20% EtOAc/Hexanes) provided 261 mg (93% yield) of alcohol **10** as a clear oil. This alcohol was obtained as a single diastereomer by ¹H NMR (ratio >20:1) [α]_D²⁰ +13.5 (*c* 0.84, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.10 (d, *J* = 3.1 Hz, 1H), 5.97 (d, *J* = 3.1 Hz, 1H), 5.04 (m, 1H), 4.59 (q, *J* = 6.8 Hz, 1H), 2.88–2.80 (m, 2H), 2.01 (s, 3H), 1.92 (d, *J* = 5.2 Hz, 1H), 1.83–1.79 (m, 2H), 1.65–1.55 (m, 2H), 1.42 (m, 1H), 1.31–1.30 (m, 5H), 0.93–0.86 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) δ : 170.7, 155.9, 151.3, 107.6, 106.7, 73.8, 68.0, 35.6, 32.6, 31.7, 26.8, 25.4, 22.7, 21.3, 14.2, 9.7; FT-IR (neat) ν_{max} = 3447, 2958, 2933, 2859, 1739, 1559, 1464, 1433, 1373, 1242, 1021, 964, 792 cm⁻¹; LRMS-ESI (+) *m/z* 305.1 [M + Na]⁺.

(R)-1-((2R,5S,6R)-5-Hydroxy-6-pentyltetrahydro-2H-pyran-2-yl)butan-2-yl acetate (8):

To furanyl alcohol **10** (317 mg, 1.12 mmol) dissolved in THF (8 mL) and H₂O (2 mL) at 0 °C was added KBr (6.7 mg, 0.06 mmol), NaHCO₃ (47 mg, 0.56 mmol) and oxone (826 mg, 1.34 mmol) after which a light yellow color developed. After stirring at 0 °C for 30 min, the reaction was quenched with saturated throughout NaHCO₃, extracted with EtOAc, washed with brine and dried over Na₂SO₄. The resulting crude hemiacetal **9** was used directly for the subsequent reaction.

To crude hemiacetal dissolved in CH₂Cl₂ (10 mL) at –45 °C was added Et₃SiH (0.89 mL, 5.6 mmol) and TFA (1.3 mL, 16.8 mmol) dropwise upon which a yellow color developed. The reaction stirred at this temperature for 3 h, then was allowed to warm to room temperature. After stirring for 30 min at 23 °C, the reaction was cooled to 0 °C and

additional Et₃SiH (0.895 mL, 5.6 mmol) was added followed by BF₃·OEt₂ (0.415 mL, 3.36 mmol). After stirring for 30 min at 0 °C, the reaction was quenched slowly with satd. NaHCO₃ until the effervescence ceased. The reaction was extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (10% to 30% EtOAc/Hexanes) provided 151 mg (47% yield over two steps) of tetrahydropyran derivative **8** as a clear oil. It was obtained as a single diastereomer by ¹H NMR analysis (>20:1) [α]_D²⁰ +2.7 (*c* 0.49, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 5.01 (m, 1H), 3.30–3.22 (m, 2H), 2.95 (td, *J* = 9.0, 2.4 Hz, 1H), 2.06 (m, 1H), 2.03 (s, 3H), 1.79 (m, 1H), 1.65–1.53 (m, 6H), 1.43–1.24 (m, 9H), 0.90–0.85 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) δ : 170.7, 82.2, 74.1, 72.9, 70.9, 40.0, 33.3, 32.2, 32.1, 32.0, 31.8, 27.8, 25.0, 22.7, 21.3, 14.2, 9.5; FT-IR (neat) $\nu_{\max}$ = 3451, 2929, 2859, 1739, 1718, 1461, 1436, 1373, 1242, 1080, 1056, 1024, 954 cm⁻¹; LRMS-ESI (+) *m/z* 287.1 [M+H]⁺.

1-((2*R*,5*S*,6*R*)-5-((*tert*-Butyldimethylsilyl)oxy)-6-pentyltetrahydro-2*H*-pyran-2-yl)butan-2-one (15):

To tetrahydropyran derivative **8** (141 mg, 0.49 mmol) dissolved in CH₂Cl₂ (5 mL) at 0 °C was added 2,6-lutidine (0.23 mL, 1.97 mmol) and TBSOTf (0.34 mL, 1.48 mmol) and the reaction was warmed to 23 °C. After stirring for 12 h, the reaction was quenched with satd. NaHCO₃, extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (5% to 10% EtOAc/Hexanes) gave 200 mg (quantitative) of the resulting silyl ether as a clear oil.

To the silyl ether (186 mg, 0.46 mmol) in MeOH (3 mL) at 0 °C was added K₂CO₃ (6.4 mg, 0.05 mmol) upon which a yellow color developed. The reaction was warmed to room temperature. After 12 h, the reaction was diluted with H₂O and EtOAc, extracted with EtOAc, washed with brine and dried over Na₂SO₄. Purification by column chromatography (5% to 20% EtOAc/Hexanes) gave 155 mg (93% yield) of the resulting alcohol as a clear oil.

To the above alcohol (144 mg, 0.40 mmol) dissolved in CH₂Cl₂ (4 mL) at 0 °C was added NaHCO₃ (202 mg, 2.41 mmol) followed by DMP (341 mg, 0.8 mmol) and the reaction was warmed to room temperature. After 12 h, the reaction was quenched with a 1:1 mixture of satd. sodium thiosulfate and saturated NaHCO₃. The mixture was extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (5% to 10% EtOAc/Hexanes) gave 138 mg (96% yield) of ketone **15** as a clear oil [α]_D²³ +32.0 (*c* 0.58, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 3.71 (m, 1H), 3.21 (m, 1H), 3.02 (m, 1H), 2.62 (dd, *J* = 14.9, 8.2 Hz, 1H), 2.53–2.41 (m, 2H), 2.36 (dd, *J* = 14.9, 4.7 Hz, 1H), 1.94 (m, 1H), 1.78–1.69 (m, 2H), 1.52–1.22 (m, 9H), 1.03 (t, *J* = 7.3 Hz, 3H), 0.86 (s, 12H), 0.04 (s, 6H); ¹³C NMR (125 MHz, CDCl₃) δ : 210.3, 82.5, 74.3, 71.4, 48.7, 37.2, 33.6, 32.2, 31.9, 31.5, 25.9, 25.1, 22.7, 18.1, 14.2, 7.6, -3.9, -4.6; FT-IR (neat) $\nu_{\max}$ = 2954, 2929, 2855, 1721, 1464, 1376, 1253, 1105, 887, 837, 774 cm⁻¹; LRMS-ESI (+) *m/z* 357.3 [M+H]⁺.

Decytospolide A (6):

To ketone **15** (126 mg, 0.35 mmol) dissolved in THF (4 mL) at 0 °C was added TBAF (0.71 mL, 0.71 mmol, 1 M in THF) and the reaction was warmed to 23 °C. After 12 h, the reaction

was concentrated and purification by column chromatography (30% to 50% EtOAc/Hexanes) provided 82 mg (96% yield) of decytospolide A (**6**) as a clear oil [α]_D²³ +18.4 (*c* 0.45, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 3.72 (m, 1H), 3.24 (td, *J* = 9.8, 4.5 Hz, 1H), 3.01 (m, 1H), 2.64 (dd, *J* = 15.0, 8.1 Hz, 1H), 2.53–2.41 (m, 2H), 2.37 (dd, *J* = 15.0, 4.8 Hz, 1H), 2.06 (m, 1H), 1.81–1.69 (m, 2H), 1.54 (brs, 1H), 1.47–1.24 (m, 9H), 1.03 (t, *J* = 7.3 Hz, 3H), 0.86 (t, *J* = 6.7 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 210.2, 82.3, 74.2, 70.6, 48.5, 37.2, 33.1, 32.1, 31.9, 31.4, 25.1, 22.7, 14.2, 7.6; FT-IR (neat) ν_{max} = 3447, 2929, 2859, 1714, 1457, 1376, 1077 cm⁻¹; LRMS-ESI (+) *m/z* 243.1 [M+H]⁺; HRMS-ESI (+) *m/z* calc'd for C₁₄H₂₇O₃ [M+H]⁺: 243.1955, found 243.1958.

Decytospolide B (**7**):

To decytospolide A (**6**) (73 mg, 0.30 mmol) dissolved in CH₂Cl₂ (3 mL) at 0 °C was added pyridine (73 μ L, 0.9 mmol), acetic anhydride (85 μ L, 0.9 mmol) and a few crystals of DMAP and the reaction was warmed to 23 °C. After 1.5 h, the reaction was diluted with H₂O, extracted with CH₂Cl₂, washed with brine and dried over Na₂SO₄. Purification by column chromatography (10% to 20% EtOAc/Hexanes) provided 77 mg (89% yield) of decytospolide B (**7**) as a clear oil [α]_D²⁰ +28.1 (*c* 0.75, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 4.43 (td, *J* = 10.0, 4.6 Hz, 1H), 3.75 (m, 1H), 3.21 (td, *J* = 9.1, 2.4 Hz, 1H), 2.66 (dd, *J* = 15.2, 8.0 Hz, 1H), 2.52–2.40 (m, 2H), 2.36 (dd, *J* = 15.3, 4.9 Hz, 1H), 2.12 (m, 1H), 2.02 (s, 3H), 1.73 (m, 1H), 1.53–1.35 (m, 4H), 1.29–1.20 (m, 6H), 1.02 (t, *J* = 7.3 Hz, 3H), 0.85 (t, *J* = 6.9 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 209.9, 170.4, 79.4, 74.3, 72.2, 48.3, 37.3, 32.0, 31.8, 30.9, 29.5, 24.9, 22.7, 21.3, 14.1, 7.6; FT-IR (neat) ν_{max} = 2933, 2859, 1739, 1714, 1457, 1373, 1235, 1080, 1042 cm⁻¹; LRMS-ESI (+) *m/z* 285.1 [M+H]⁺; HRMS-ESI (+) *m/z* calc'd for C₁₆H₂₈O₄Na [M+Na]⁺: 307.1880, found 307.1884.

Methyl 2-(5-hexanoylfuran-2-yl)acetate (**18**):

Hexanoic acid (94 μ L, 0.75 mmol) was dissolved in thionyl chloride (6 mL) and refluxed for 3 h. The excess thionyl chloride was distilled off to give the resulting hexanoyl chloride as a dark yellow oil. It was then dissolved in CH₂Cl₂ (6 mL) and cooled to 0°C. SnCl₄ (1.13 mL, 1 M in CH₂Cl₂, 1.13 mmol) was then added slowly dropwise and the resulting solution stirred at 0°C for 45 min. Methyl ester **17** (105.3 mg, 0.75 mmol) was dissolved in CH₂Cl₂ (2 mL) and the resulting solution was added slowly to the reaction over 10 min. The reaction was then stirred at 0°C for 45 min before being quenched with ice. The biphasic mixture was separated and the aqueous layer extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by column chromatography (10 to 20% EtOAc/Hexanes) to give ketone **18** (102 mg, 57%) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ : 7.12 (d, *J* = 3.5 Hz, 1H), 6.41 (d, *J* = 3.5 Hz, 1H), 3.77 (s, 2H), 3.74 (s, 3H), 2.84–2.66 (m, 2H), 1.70 (t, *J* = 7.4 Hz, 2H), 1.34 (h, *J* = 3.6 Hz, 4H), 0.90 (td, *J* = 7.1, 5.9, 3.4 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 189.33, 168.68, 152.16, 129.03, 118.12, 110.76, 77.24, 76.92, 76.60, 52.38, 38.23, 34.01, 31.39, 24.06, 22.34, 13.82; ESI-API MS: [M+H] = 239.1; HRMS-ESI (+) *m/z* calc'd for C₁₃H₁₀O₄ [M+H]⁺: 239.1280, found 239.1282.

Methyl (*R*)-2-(5-(1-hydroxyhexyl)furan-2-yl)acetate (19**):**

Ketone **18** (60.5 mg, 0.25 mmol) was dissolved in CH₂Cl₂ (7 mL) and Et₃N (0.5 mL, 3.81 mmol) was added followed by HCO₂H (143 μL, 3.81 mmol). RuCl[(*R,R*)-TsDPEN] (mesitylene) (3.9 mg, 0.006 mmol) was dissolved in CH₂Cl₂ (0.5 mL) and then added to the reaction. The reaction was then set to stir at 50°C for 12 h before being quenched with H₂O. It was then extracted with CH₂Cl₂ and the combined organic layers washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by column chromatography (10% to 30% EtOAc/Hexanes) to give optically active alcohol **19** (50.3 mg, 83%) as a clear oil. ¹H-NMR (400 MHz, CDCl₃) δ 6.15 (d, *J* = 0.9 Hz, 2H), 4.62 (t, *J* = 6.8 Hz, 1H), 3.71 (s, 3H), 3.66 (s, 2H), 1.98 (s, 1H), 1.88 – 1.74 (m, 2H), 1.43 (tdd, *J* = 10.0, 8.1, 7.5, 4.1 Hz, 1H), 1.36 – 1.22 (m, 5H), 0.88 (q, *J* = 4.9, 4.1 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ 169.76, 156.47, 146.83, 108.51, 106.61, 67.66, 52.15, 35.30, 33.84, 31.48, 25.11, 22.44, 13.90; ESI-API MS: [M+Na] = 263.1; HRMS-ESI (+) *m/z* calc'd for C₁₃H₂₀O₄Na [M+H]⁺: 263.1254, found 263.1256; 99% ee, determined by HPLC using Chiralpak IA3 and gradient of 0–10% isopropanol/hexanes (*t*_{major} = 25.7 min, *t*_{minor} = 24.8 min)

Methyl 2-((2*R*,5*S*,6*R*)-5-hydroxy-6-pentyltetrahydro-2*H*-pyran-2-yl)acetate (20**):**

Furanyl alcohol **19** (16.9 mg, 0.07 mmol) was dissolved in THF (2 mL) and H₂O (0.5 mL) and cooled to 0°C. KBr (0.4 mg, 0.003 mmol), NaHCO₃ (2.9 mg, 0.03 mmol) and oxone (51.9 mg, 0.08 mmol) were then added and the reaction slowly warmed to 23 °C. After 3 h, it was quenched with saturated NaHCO₃ and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was used immediately for the next reaction.

The crude oil was dissolved in CH₂Cl₂ (3 mL) and Et₃SiH (112 μL, 0.7 mmol) was added. The reaction was then cooled to –40°C and BF₃•OEt₂ (52 μL, 0.42 mmol) was added slowly dropwise. The reaction was then stirred at –40°C for 16 h before being warmed to 23 °C and stirred for an additional 1 h. It was then quenched with saturated NH₄Cl and extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by column chromatography (20% to 30% EtOAc/Hexanes) give tetrahydropyran derivative **20** (10.0 mg, 58%) as a clear oil. ¹H NMR (400 MHz, CDCl₃) δ : 3.80 – 3.69 (m, 1H), 3.67 (s, 3H), 3.27 (ddd, *J* = 10.4, 8.9, 4.6 Hz, 1H), 3.04 (td, *J* = 9.0, 2.4 Hz, 1H), 2.53 (dd, *J* = 14.9, 8.0 Hz, 1H), 2.40 (dd, *J* = 14.9, 5.4 Hz, 1H), 2.13 – 2.01 (m, 1H), 1.85 – 1.72 (m, 2H), 1.54 – 1.29 (m, 3H), 1.32 – 1.22 (m, 6H), 0.87 (t, *J* = 6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ 171.67, 82.12, 73.83, 70.39, 51.50, 40.77, 32.75, 31.74, 31.65, 30.84, 24.85, 22.56, 13.95; ESI-API MS: [M+H] = 245.1, [M+Na] = 267.1; HRMS-ESI (+) *m/z* calc'd for C₁₃H₂₅O₄ [M+H]⁺: 245.1747, found 245.1750.

2-((2*R*,5*S*,6*R*)-5-Hydroxy-6-pentyltetrahydro-2*H*-pyran-2-yl)-*N*-methoxy-*N*-methylacetamide (21**):**

To a solution of NH(OMe)Me•HCl (16.0 mg, 0.16 mmol) in dry THF (3 mL) and cooled to –30°C, *i*-PrMgCl (450 μL, 0.45 mmol) was added slowly dropwise and the reaction stirred at –30°C for 1 h. Methyl ester **20** (10 mg, 0.04 mmol) was dissolved in dry THF and then

added slowly, dropwise to the reaction which was stirred at -30°C for an additional 4 h before being quenched with saturated NH_4Cl . It was extracted with EtOAc and the combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The crude residue was purified by column chromatography (50% to 80% EtOAc/Hexanes) to give Weinreb amide **21** (10.4 mg, 93%) as a clear oil. ^1H NMR (800 MHz, CDCl_3) δ 3.85 – 3.80 (m, 1H), 3.71 (d, J = 2.2 Hz, 3H), 3.32 – 3.27 (m, 1H), 3.20 (s, 3H), 3.09 (ddd, J = 8.9, 6.5, 2.5 Hz, 1H), 2.85 (d, J = 13.4 Hz, 1H), 2.43 (dd, J = 15.1, 6.3 Hz, 1H), 2.13 – 2.08 (m, 1H), 1.90 – 1.79 (m, 2H), 1.55 – 1.44 (m, 2H), 1.46 – 1.39 (m, 1H), 1.41 – 1.31 (m, 1H), 1.33 – 1.27 (m, 5H), 0.90 (td, J = 7.0, 2.3 Hz, 3H); ^{13}C NMR (201 MHz, CDCl_3) δ 172.04, 82.24, 74.14, 70.58, 61.33, 37.95, 33.02, 32.03, 31.90, 31.25, 29.71, 25.01, 22.66, 14.07; ESI-API MS: $[\text{M}+\text{H}] = 274.0$, $[\text{M}+\text{Na}] = 296.1$.

Decytospolide A (6) via Weinreb amide 21:

Weinreb amide **21** (10.4 mg, 0.038 mmol) was dissolved in dry THF (2 mL) and cooled to 0°C . EtMgBr in THF solution (305 μL , 0.3 mmol) was added slowly dropwise and the reaction slowly warmed to 23°C . It was stirred for 5 h, then quenched with saturated NH_4Cl and extracted with EtOAc. The combined organic layers were then washed with brine and dried over Na_2SO_4 . The crude residue was purified by column chromatography (20% to 50% EtOAc/Hexanes) to give decytospolide A, **6** (8.9 mg, 97%) as a clear oil. ^1H NMR (400 MHz, CDCl_3) δ 3.73 (dddd, J = 10.2, 8.0, 4.8, 2.0 Hz, 1H), 3.25 (ddd, J = 10.6, 9.0, 4.6 Hz, 1H), 3.02 (td, J = 8.9, 2.5 Hz, 1H), 2.65 (dd, J = 15.0, 8.1 Hz, 1H), 2.58 – 2.29 (m, 3H), 2.07 (ddd, J = 12.0, 5.5, 2.8 Hz, 1H), 1.85 – 1.70 (m, 2H), 1.53 – 1.41 (m, 1H), 1.46 (s, 2H), 1.44 – 1.33 (m, 1H), 1.37 – 1.24 (m, 5H), 1.03 (t, J = 7.3 Hz, 3H), 0.93 – 0.83 (m, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 209.96, 82.04, 74.00, 70.42, 48.26, 36.98, 32.83, 31.84, 31.70, 31.16, 24.88, 22.51, 13.92, 7.41; ESI-API MS: $[\text{M}+\text{H}] = 243.1$, $[\text{M}+\text{Na}] = 265.1$; HRMS-ESI (+) m/z calc'd for $\text{C}_{14}\text{H}_{27}\text{O}_3$ $[\text{M}+\text{H}]^+$: 243.1955, found 243.1958.

Decytospolide B (7) via Weinreb amide 21:

Above synthetic decytospolide A (9.7 mg, 0.04 mmol) was dissolved in CH_2Cl_2 (3 mL) and cooled to 0°C . Pyridine (10 μL , 0.12 mmol), Ac_2O (4 μL , 0.04 mmol) and a few crystals of DMAP were added and the reaction warmed to 23°C . The reaction was stirred for 3 h before being quenched with water. It was extracted with CH_2Cl_2 and the combined organic layers washed with brine and dried over Na_2SO_4 . The crude residue was purified by column chromatography (10 to 30% EtOAc/Hexanes) to give **7** (9.8 mg, 86%) as a clear oil. ^1H NMR (400 MHz, CDCl_3) δ 4.44 (td, J = 10.0, 4.7 Hz, 1H), 3.76 (ddd, J = 10.9, 7.8, 4.9 Hz, 1H), 3.22 (td, J = 9.0, 2.6 Hz, 1H), 2.67 (dd, J = 15.2, 8.0 Hz, 1H), 2.58 – 2.33 (m, 3H), 2.13 (ddd, J = 11.3, 5.6, 3.3 Hz, 1H), 2.03 (s, 3H), 1.79 – 1.70 (m, 1H), 1.58 – 1.36 (m, 3H), 1.40 – 1.17 (m, 7H), 1.03 (t, J = 7.3 Hz, 3H), 0.86 (t, J = 6.9 Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 209.77, 170.22, 79.21, 74.09, 71.95, 48.09, 37.10, 31.82, 31.60, 30.70, 29.24, 24.71, 22.48, 21.10, 13.89, 7.40; ESI-API MS: $[\text{M}+\text{H}] = 285.1$, $[\text{M}+\text{Na}] = 307.2$; HRMS-ESI (+) m/z calc'd for $\text{C}_{16}\text{H}_{28}\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$: 307.1880, found 307.1884.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support of this work was provided by the National Institutes of Health and Purdue University.

Notes and references

1. Kang EJ and Lee E, Chem. Rev, 2005, 105, 4348–4378. [PubMed: 16351047]
2. Yeung KS and Paterson I, Chem. Rev, 2005, 105, 4237–4313. [PubMed: 16351045]
3. Fuwa H Mar. Drugs, 2016, 14, 65.
4. Nasir NM, Ermanis K and Clarke PA Org. Biomol. Chem 2014, 12, 3323–3335. [PubMed: 24744139]
5. Brimble MA, Kaliappan KP, Moreno-Vargas AJ, Palanichamy K, Perry MA, Rainier JD, Rychnovsky SD, Sizemore N, Stubbing LA and Vogel P, Synthesis of saturated oxygenated heterocycles. I, 5- and 6-membered rings, Springer Berlin 2014.
6. Arundale E and Mikeska LA Chem. Rev 1952, 51, 505–555.
7. Zeng J, Tan YJ, Ma J, Leow ML, Tirtorahardjo D and Liu X-W Chem. Eur. J 2014, 20, 405–409. [PubMed: 24285699]
8. Achmatowicz O, Bukowski P, Szechner B, Zwierzchowska Z and Zamojski A Tetrahedron 1971, 27, 1973–1996.
9. Achmatowicz O, Gryniewicz G and Szechner B, Tetrahedron, 1976, 32, 1051–1054.
10. Ghosh AK and Brindisi M RSC Advances 2016, 6, 111564–111598. [PubMed: 28944049]
11. Deska J, Thiel D and Gianolio E, Synthesis 2015, 47, 3435–3450.
12. Ghosh AK, Chen Z-H, Effenberger KA and Jurica MS J. Org. Chem 2014, 79, 5697–5709. [PubMed: 24873648]
13. Ghosh AK and Li J Org. Lett 2011, 13, 66–69. [PubMed: 21126066]
14. Koh P-F and Loh T-P Green Chem 2015, 17, 3746–3450.
15. Lu S, Sun P, Li T, Kurtán T, Mándi A, Antus S, Krohn K, Draeger S, Schulz B, Yi Y, Li L and Zhang W J. Org. Chem 2011, 76, 9699–9710. [PubMed: 22011230]
16. Dale JA and Mosher HS J. Am. Chem. Soc 1973, 95, 512–519.
17. Krishna PR, Nomula R and Ramana DV Tetrahedron Lett 2012, 53, 3612–3614.
18. Clarisse D and Fache F Tetrahedron Lett 2014, 55, 2221–2222.
19. Yahata K, Minami M, Watanabe K and Fujioka H Org. Lett 2014, 16, 3680–3683. [PubMed: 25006698]
20. Gore PH Chem. Rev 1955, 55, 229–281.
21. Tofi M, Koltsida K and Vassilikogiannakis G Org. Lett 2009, 11, 313–316. [PubMed: 19072227]
22. Hashiguchi S, Fujii A, Takehara J, Ikariya T and Noyori R J. Am. Chem. Soc 1995, 117, 7562–7563.
23. Fujii A, Hashiguchi S, Uematsu N, Ikariya T and Noyori R, R. J. Am. Chem. Soc 1996, 118, 2521–2522.
24. Ghosh AK and Chen Z-H Org. Lett 2013, 15, 5088–5091. [PubMed: 24050251]
25. Georgiadis M-P, Albizati KF and Georgiadis TM Org. Prep. Proc. Int 1992, 24, 95–118.
26. Li Z and Tong R J. Org. Chem 2016, 81, 4847–4855. [PubMed: 27167167]
27. Cafiero LR and Snowden TS Org. Lett 2008, 10, 3853–3856. [PubMed: 18686964]
28. Shamshina JL and Snowden TS Org. Lett 2006, 8, 5881–5884. [PubMed: 17134296]

Figure 1.
Achmatowicz reaction and structures of the decytospolides

Scheme 1.
Retrosynthetic analysis of the decytospolides

Scheme 2.

Synthesis of alcohol **10**. Reagents and Conditions: (a) (*R*)-1,2-epoxybutane, *n*-BuLi, THF, 0 °C to 23 °C; (b) Ac₂O, Et₃N, DMAP, CH₂Cl₂, (84%, 2-steps); (c) C₅H₁₁COCl, SnCl₄, CH₂Cl₂, 0 °C (75%); (d) RuCl(mesitylene), [(*R,R*)-Ts-DPEN], (1 mol%), HCO₂H, Et₃N, CH₂Cl₂, 55° C (93%).

Scheme 3.

Synthesis of the decytospolides. Reagents and Conditions: (a) Oxone, KBr, NaHCO_3 , THF/ H_2O (4:1); (b) Et_3SiH , TFA, $\text{BF}_3 \cdot \text{OEt}_2$, CH_2Cl_2 , -45°C to rt, (47%, 2-steps); (c) Et_3SiH , $\text{BF}_3 \cdot \text{OEt}_2$, CH_2Cl_2 , -45°C ; (d) TBSOTf, 2,6-lutidine, CH_2Cl_2 , 0°C to 23°C ; (e) K_2CO_3 , MeOH; (f) DMP, NaHCO_3 , CH_2Cl_2 , 0°C to 23°C , (89%, 3-steps); (g) TBAF, THF, 23°C (99%); (h) Ac_2O , pyr, DMAP, 23°C , CH_2Cl_2 (99%).

Scheme 4.

Synthesis of alcohol **20**. Reagents and conditions: (a) $\text{C}_6\text{H}_{11}\text{COCl}$, SnCl_4 , CH_2Cl_2 , $0\text{ }^\circ\text{C}$ (57%); (b) $\text{RuCl}(\text{mesitylene})[(R,R)\text{-Ts-DPEN}]$, (1 mol%), Et_3N , HCO_2H , CH_2Cl_2 , $50\text{ }^\circ\text{C}$, 12 h (83%, 99% ee); (c) KBr , NaHCO_3 , oxone, $\text{THF}/\text{H}_2\text{O}$ (4:1), $0\text{ }^\circ\text{C}$ to $23\text{ }^\circ\text{C}$, 3 h; (d) $\text{BF}_3\cdot\text{OEt}_2$, Et_3SiH , CH_2Cl_2 , $-40\text{ }^\circ\text{C}$, 16 h (58%, over 2-steps).

Scheme 5.

Synthesis of decytospolides. Reagents and conditions: (a) $\text{NH(OMe)Me} \cdot \text{HCl}$, $i\text{-PrMgCl}$, THF, -30°C , 5 h (93%); (b) EtMgBr , THF, 0°C to rt, 5 h (97%); (c) Ac_2O , pyridine, DMAP, CH_2Cl_2 , 0°C to 23°C , 3 h (86%).

Table 1.Reduction of the dihydropyranone hemiacetal **9** with various Lewis acids^a

Entry	Lewis Acid (equiv)	Et ₃ SiH (equiv)	Time	Yield for enone 14	Yield for THP 8
1	TFA [15]	5	24 h	40%	trace
2	TFA [15] BF ₃ •OEt ₂ [3]	5, then 5 more	4 h	none	47%
3	BF ₃ •OEt ₂ [6]	10	6 h	none	56%
4	SnCl ₄ [6]	10	4 h	none	82%
5	Cu(OTf) ₂ [6]	10	18 h	trace	none
6	TiCl ₄ [6]	10	18 h	14%	none
7	Sc(OTf) ₃ [6]	10	18 h	trace	none

^a All reactions were carried out in CH₂Cl₂ at –45 °C