Published in final edited form as:

Soc Sci Med. 2018 September ; 213: 165–172. doi:10.1016/j.socscimed.2018.07.049.

Why smokers avoid cigarette pack risk messages: Two randomized clinical trials in the US

Marissa G. Hall^{1,2}, Jennifer R. Mendel¹, Seth M. Noar^{3,1}, and Noel T. Brewer^{2,1}

¹Lineberger Comprehensive Cancer Center, University of North Carolina

²Department of Health Behavior, Gillings School of Global Public Health, University of North Carolina

³School of Media and Journalism, University of North Carolina

Abstract

Background.—Message avoidance (e.g., trying not to look at the message) may be motivated by reactance, a maladaptive rejection of the message. An alternative view is that avoidance indicates that a message is eliciting fear and other negative affect, thereby increasing the likelihood of behavioral change. We sought to identify which psychological mechanism (reactance or fear and other negative affect) explains message avoidance. We also examined whether avoidance was associated with more forgoing or butting out of cigarettes.

Methods.—Trial 1 randomly assigned 2,149 adult U.S. smokers to receive either pictorial warnings (intervention) or text-only warnings (control) on their cigarette packs for four weeks in 2014 and 2015. Trial 2 randomly assigned 719 adult U.S. smokers to receive either messages about toxic chemicals in cigarette smoke (intervention) or messages about not littering cigarette butts (control) for three weeks in 2016 and 2017. Negative affect included fear, anxiety, disgust, sadness, and guilt. Reactance included perceived threat to freedom, anger, and counterarguing.

Results.—Intervention messages led to greater message avoidance in both trials (both $p < .001$). In Trial 1, intervention messages elicited greater negative affect, which in turn was associated with greater avoidance (mediated effect = .21, $p < .001$). In contrast, reactance explained only a small part of the effect in Trial 1 (mediated effect = .03, $p < .001$). Similarly, in Trial 2, intervention messages elicited greater negative affect, which was associated with more avoidance (mediated effect = .12, $p < .001$); reactance did not explain any of the effect. In both trials, avoidance was associated with more forgoing or butting out of cigarettes ($p < .001$).

Conclusions.—Smokers may avoid cigarette pack risk messages because they evoke aversive types of emotion. These studies add to a growing body of evidence that, in the context of cigarette pack messages, avoidance is not a form of defensive processing but instead a sign of *deeper* processing.

Corresponding author: Marissa G. Hall, PhD, Department of Health Behavior, Gillings School of Global Public Health, University of North Carolina, 324 Rosenau Hall CB7440, Chapel Hill, NC 27599, mghall@unc.edu, 919-843-3105.

Conflict of Interest: Noel T. Brewer has served as a paid expert consultant in litigation against the tobacco industry. The other authors declare that they have no conflicts of interest.

Keywords

Cigarette pack warnings; pictorial warnings; avoidance; defensive processing; risk messages; reactance; fear appeals; negative affect.

Introduction

Tobacco use is the leading cause of disease and death worldwide (World Health Organization, 2012). Health warnings on cigarette packs are a promising, low-cost solution for combating the tobacco epidemic. Cigarette pack warnings can attract attention (Noar et al., 2017) and inform smokers (Brewer et al., 2018a; Noar et al., 2016). The addition of images to text-only warnings enhances their effectiveness; pictorial warnings increase cigarette quit attempts (Brewer et al., 2016) and, once implemented in the US, could help prevent more than 650,000 deaths over the next 50 years (Levy et al., 2016).

Health behavior and health communication theories suggest that risk communications such as cigarette pack messages may sometimes cause avoidance, an unintended outcome that is unlikely to lead to beneficial effects. For example, the Extended Parallel Process Model (Witte, 1992) posits that under some conditions, fear communications can lead to maladaptive coping and ultimately generate defensive motivation including avoidance, suggesting the risk message is being rejected and is ineffective. Similarly, the Transtheoretical Model (DiClemente et al., 1991; Prochaska & DiClemente, 1986) views consciousness raising (e.g., information seeking) as a sign of increased readiness to change behavior, with those not ready to change actively avoiding information about health behavior change.

Avoidance itself can take many forms. In the context of cigarette pack risk messages, avoidance typically includes behaviors such as putting the pack out of sight, or trying to avoid looking at the risk message when the pack is in sight (Borland et al., 2009a; Osman et al., 2017; Thrasher et al., 2016). Avoidance can also include cognitive strategies to divert attention away from the message (McQueen et al., 2013), such as putting the message out of one's mind and avoiding thinking about what the message says. We conceptualize these behavioral and cognitive actions as components of avoidance.

Research from qualitative (Hardcastle et al., 2015), observational (Borland et al., 2009b; Hammond et al., 2004; Noar et al., 2017), eye tracking (Maynard et al., 2014), and experimental studies (Brewer et al., 2018b; McCloud et al., 2017; McQueen et al., 2015) indicate that some smokers avoid cigarette pack warnings after initial exposure. Studies from Canada, the US, Australia, and Mexico suggest that about 30% of smokers regularly attempt to avoid looking at or thinking about cigarette pack warnings (Cho et al., 2016; Hammond et al., 2003; Thrasher et al., 2016). However, in contrast with the Extended Parallel Process Model's predictions, avoidance of cigarette pack warnings may not be a maladaptive response. Research shows that avoidance does not appear to undermine the effect of cigarette pack warnings (Borland et al., 2009a; Cho et al., 2016), and may instead be a marker for beneficial effects on behavior (Brewer et al., 2018b; Cho et al., 2016; Thrasher et al., 2016). For instance, two longitudinal studies have found avoidance of cigarette pack

warnings to be associated with more subsequent quit attempts (Cho et al., 2016; Thrasher et al., 2016). Ironic Process Theory offers one explanation for this finding: efforts to avoid thoughts often fail and make these thoughts even more pronounced (Wegner, 1994). Moreover, cigarette pack risk messages appear on products that regular smokers see and use very frequently, potentially making them harder to fully ignore.

Understanding *why* smokers avoid risk messages on their cigarette packs may shed light on the counterintuitive finding that avoidance of messages does not detract from, and may instead be a marker of, message effectiveness. However, studies to date have not explored the reasons why smokers avoid risk messages on cigarette packs. Using data from two large trials, we aimed to determine whether exposure to risk messages on smokers' own cigarette packs increased message avoidance. We also sought to explore the reasons why smokers avoided cigarette pack risk messages to investigate whether avoidance is a result of adaptive or maladaptive responding. Specifically, we examined whether smokers avoided risk messages because of negative affect, which is an adaptive response shown to promote quitting (Brewer et al., 2018b). We also examined whether smokers avoided risk messages because of message reactance (a maladaptive response, hypothesized (Brehm, 1966; Brehm & Brehm, 1981) and shown to undermine quitting (Brewer et al., 2018b)). We expected that negative affect and message reactance would both be positively associated with avoidance, but that the effect of negative affect would be stronger based on previous work showing avoidance being associated with greater message impact (Brewer et al., 2018b; Cho et al., 2016; Thrasher et al., 2016). Finally, we examined whether avoidance was associated with greater forgoing or butting out of cigarettes. We expected that avoidance would be associated with more forgoing or butting out, based on prior work showing avoidance is a marker for greater cigarette pack warning effectiveness (Brewer et al., 2018b; Cho et al., 2016; Thrasher et al., 2016).

Methods

Trial 1

Participants.—From September 2014 to August 2015, we recruited a convenience sample of 2,149 adult smokers in North Carolina and California, U.S. to participate in a randomized trial comparing the impact of pictorial versus text-only warnings. Details of the trial including the protocol, survey development, and participant recruitment have been previously published (Brewer et al., 2016; Brodar et al., 2016). Other papers using this dataset have explored the impact of pictorial warnings on quit attempts (Brewer et al., 2016) and reactance (Hall et al., 2017a), mediators (including avoidance) of the impact of pictorial cigarette pack warnings (Brewer et al., 2018b), the frequency and content of social interactions (Morgan et al., 2017), trajectories of pictorial warnings' impact (Parada et al., 2017), and attitudes toward regulation of tobacco products (Hall et al., 2018; Kowitt et al., 2017). Participants were age 18 or older, proficient in English, and current smokers, defined as having smoked at least 100 cigarettes during their lifetime and now smoking every day or some days. Exclusion criteria included pregnancy, current enrollment in a smoking cessation trial, smoking only roll-your-own cigarettes, smoking fewer than seven cigarettes per week, and living in the same household as another trial participant.

Procedures.—Smokers received warnings on their own cigarette packs for four weeks using a protocol developed by our research team whereby participants brought in an eight-day supply of cigarettes weekly (Brewer et al., 2015). They were randomly assigned to have one of four pictorial warnings applied to the top half of the front and back panels of their cigarette packs (intervention), or one of four text-only warnings applied to the side of their cigarette packs placed over the current Surgeon General’s warning (control), for the duration of the trial (Figure 1). Randomization created groups that did not differ on demographic characteristics (Table 1, all $p>.05$).

Participants completed two computer surveys at the first visit (at baseline and immediately after seeing their assigned warning on their cigarette packs), and one survey at each visit thereafter. Participants received a cash incentive at the end of each visit, totaling up to \$185 in North Carolina and \$200 in California. Participation incentives were higher in California due to higher cost of living. At the end of the final follow-up appointment, participants received information about local smoking cessation programs.

Measures.—The baseline survey assessed participant demographics and smoking behavior. The second baseline survey assessed negative affect and message reactance immediately after participants first saw their assigned warnings (Table 2). To measure negative affect (i.e., anxious, disgusted, guilty, sad, scared), we adapted items from several sources (Keller & Block, 1996; Nonnemaker et al., 2010; Watson et al., 1988). We originally planned to examine fear alone as a mediator given its importance in the Extended Parallel Process Model (Witte, 1992), but our previously-reported confirmatory factor analysis supports treating negative affect as a single latent factor (Hall et al., 2017a). We used the validated Brief Reactance to Health Warnings Scale to measure message reactance (Hall et al., 2016; Hall et al., 2017b). The final follow-up survey (four weeks after the start of the trial) assessed message avoidance using items adapted from two large-scale longitudinal studies (Table 2) (International Tobacco Control Policy Evaluation Project, 2003; Population Assessment of Tobacco and Health Study, 2018; Yong et al., 2014). The final follow-up survey also assessed the number of times participants reported forgoing or butting out a cigarette in the past week (Li et al., 2014). The items read: “In the last week, how often have you stopped yourself from having a cigarette because you wanted to smoke less?” and “In the last week, how often have you butted out a cigarette before you finished it because you wanted to smoke less?” Response options were never (coded as 0), 1–2 times (coded as 1.5), 3–4 times (coded as 3.5), 5–9 times (coded as 7), and 10 or more times (coded as 10). To create a composite variable, we summed participants’ responses to these the two items, ranging from 0–20.

Data Analysis.—Analyses used Stata/SE version 14.1 and Mplus version 8 with two-tailed tests and a critical alpha of .05. We report most results as standardized coefficients (β s). First, we created a composite score representing the average of the three message avoidance items and conducted between-group t -tests to compare message avoidance by trial arm.

Next, we examined the associations between items and their latent constructs using a measurement model that included negative affect, message reactance, and message avoidance. Analyses treated indicators for these three variables as ordinal. This model

exhibited adequate fit, with Root Mean Square Error of Approximation (RMSEA)=.058 (Steiger, 1990), Bentler Comparative Fit Index (CFI)=.995 (Bentler, 1990), and Tucker-Lewis Index (TLI)=.994 (Tucker & Lewis, 1973). Then, we used a structural equation model to examine the impact of trial arm on message avoidance; this model was just-identified and therefore did not produce fit statistics (Bollen, 1989). The final structural equation model, used to examine our mediation hypotheses and the association between avoidance and forgoing or butting out a cigarette, employed full information maximum likelihood estimation to handle missing data (Bollen, 1989; Kline, 2011; Peters & Enders, 2002; Sidi & Harel, 2018). This model used bootstrapped 95% confidence intervals with 1,000 repetitions (Hayes, 2009) and also exhibited acceptable fit (RMSEA=.051, CFI=.994, TLI=.993).

Trial 2

Participants.—We recruited a convenience sample of 784 adult smokers from the general population in the Bay Area in California, U.S. from September 2016 to March 2017 to participate in a trial comparing the impact of messages about toxic chemical versus messages about not littering. We randomized 719 participants who returned to Visit 2. We have previously reported that chemical messages did not lead to higher quit intentions, but did inform smokers of the chemicals in cigarettes and harms of smoking (Brewer et al., 2018a). Eligibility and exclusion criteria were the same as Trial 1, except that participants had to be 21 years or older, the legal age for buying tobacco products in California at the time of the study. Demographics did not differ by trial arm (Table 1, all $p>.05$). Additional details on participant recruitment and the trial including the protocol and surveys have been previously published (Brewer et al., 2018a).

Procedures.—Participants attended 5 visits, each 1 week apart, at the trial office in San Francisco, California, U.S. Smokers brought in an eight-day supply of cigarettes to Visits 1–4. At Visit 2, we randomly assigned participants to receive either labels with chemical messages (intervention) or litter messages (control) on the sides of their cigarette packs (i.e., on the side opposite from the existing Surgeon General’s warning; Figure 2). Participants received a new label, in random order, at each visit for a total of three new messages during the trial.

Participants completed computer surveys at the baseline visit and at each subsequent weekly visit. While participants completed the surveys, research staff placed the assigned labels on participants’ cigarette packs. At the end of each visit, upon completion of the survey, participants received a cash incentive that totaled up to \$300 across the trial. At the end of the final visit, we offered participants information and resources about smoking cessation.

Measures.—The baseline survey assessed participant demographics and smoking behavior. The final follow-up survey assessed negative affect, message reactance, message avoidance, and forgoing or butting out a cigarette using items that were nearly identical to Trial 1 (see Table 2 for exact wording). Frequencies of the avoidance items appear in Supplementary File 1.

Data Analysis.—Trial 2 used the same analytic approach as Trial 1. The measurement model (RMSEA=.083, CFI=.983, TLI=.997) and the mediation model (RMSEA=.067, CFI=.983, TLI=.979) both exhibited adequate fit.

Results

Trial 1 participants had a mean age of 40 years and smoked an average of nine cigarettes per day (Table 1). The mean age of Trial 2 participants was 42; they smoked an average of 11 cigarettes per day. Both trials included a substantial number of sexual minority, African American, low-education, and low-income smokers.

In Trial 1, exposure to intervention messages led to greater avoidance than the control messages (mean [SD]=2.3 [1.2] vs. 1.7 [1.0], $p<.001$). Negative affect mediated this effect (mediated effect=.21, $p<.001$; Table 3). Intervention messages increased negative affect ($\beta=.41$, $p<.001$; Figure 3), which, in turn, was associated with greater avoidance of the warnings ($\beta=.51$, $p<.001$). Intervention messages also increased message reactance ($\beta=.24$, $p<.001$) and message reactance was subsequently associated with greater avoidance ($\beta=.10$, $p<.05$). Message reactance mediated the effect of intervention messages on avoidance, but the effect was small (mediated effect=.03, $p<.001$). Finally, message avoidance was associated with more forgoing or butting out of cigarettes in the past week ($\beta=.30$, $p<.001$).

The pattern of findings in Trial 2 was nearly identical (Figure 3). Intervention messages led to greater avoidance than the control messages (mean [SD]=2.1 [1.0] vs. 1.8 [1.0], $p=.0013$). Negative affect mediated the effect of intervention messages on avoidance of the messages (mediated effect=.12, $p<.001$). Intervention messages increased negative affect ($\beta=.18$, $p<.001$), which, in turn, was associated with greater avoidance ($\beta=.68$, $p<.001$). In contrast with Trial 1, reactance explained none of the effect of intervention messages on avoidance (mediated effect=.01, $p=.65$). Intervention messages did not change message reactance ($\beta=.02$, $p=.64$), but like Trial 1, message reactance was associated with greater avoidance ($\beta=.24$, $p<.001$). As in Trial 1, message avoidance was associated with more forgoing or butting out of cigarettes in the past week ($\beta=.40$, $p<.001$).

Discussion

In two large trials with smokers, exposure to cigarette pack risk messages led to greater message avoidance compared to control messages. Most prior studies on this topic have used observational data to examine avoidance of warnings in a real-world context (Borland et al., 2009a; Fathelrahman et al., 2013; Wardle et al., 2010; Yong et al., 2013; Zacher et al., 2014; Zhang et al., 2011). These studies show that policy changes to strengthen cigarette pack warnings are followed by increases in warning avoidance (Noar et al., 2017). Experimental data from our two trials corroborate these observational studies, supporting the findings that some smokers avoid cigarette pack messages after initial exposure, and that stronger risk messages lead to greater avoidance. We also found that avoidance was associated with more forgoing or butting out of cigarettes. These behaviors predict cessation attempts in longitudinal studies in several countries (Borland et al., 2009b; Cho et al., 2016; Cho et al., 2018; Li et al., 2014; Partos et al., 2014). Our findings add to this growing body of evidence

that avoidance may be indicative of greater message effectiveness (Brewer et al., 2018b; Cho et al., 2016; Thrasher et al., 2016). However, more theoretical and empirical work is needed to understand why avoidance predicts smoking cessation.

These studies also explored the counterintuitive phenomenon that avoidance of cigarette pack risk messages is a marker for behavior change (Brewer et al., 2018b; Cho et al., 2016; Thrasher et al., 2016) by examining the psychological mechanism motivating smokers to avoid cigarette pack risk messages. In both trials, negative emotions such as anxiety, disgust, guilt, sadness, and fear were associated with message avoidance. We consider negative affect to be a sign of adaptive processing in the context of cigarette pack risk messages based on strong evidence showing that negative affect leads to health-promoting changes in attitudes and behavior (Brewer et al., 2018b; Cho et al., 2018; Emery et al., 2014; Evans et al., 2015; Hall et al., 2017a; McCaul et al., 1996; Peters et al., 2016; Peters et al., 2014; Sheeran et al., 2014). Thus, these studies show that an adaptive response (i.e., negative affect) explains most of why smokers avoid messages.

In contrast with negative affect, we found that message reactance did not have much impact on avoidance. In Trial 1, reactance explained a small part of the effect of risk messages on avoidance, and in Trial 2, it did not explain any of the effect. Prior studies have found that reactance has small associations with lower perceived message effectiveness (Hall et al., 2016), quit intentions (Hall et al., 2017a) and policy support (Hall et al., 2018); one study found null effects of reactance on behavioral intentions (Blanton et al., 2014). The undermining effects on cessation behavior are even less pronounced (Brewer et al., 2018b) or nonexistent (Cho et al., 2016; Thrasher et al., 2016). These studies build on these findings, demonstrating that reactance does not have large effects in the context of cigarette pack risk messages because reactance explained little (Trial 1) or none (Trial 2) of smokers' avoidance behaviors.

Our findings suggest that the *motivation* for avoidance might help to shed light on the consequences of avoidance. In this study, we observed two possible pathways for generating avoidance. In one pathway, smokers who engage in avoidance due to negative affect may actively try to avoid the warnings because of feeling scared or guilty about the harms of smoking (which may mean they are more likely to quit smoking). In the other pathway, smokers who report avoiding warnings because of reactance may avoid thinking about the warnings except to denigrate the warnings and reinforce their own views and biases (which may mean they are less likely to quit smoking). The specific communication context is likely to determine which of these pathways is more important and thus whether the avoidance reflects message engagement or rejection. In our studies, fear was associated with avoidance and avoidance was associated with subsequent quitting behaviors (Brewer et al., 2018b), perhaps because pack messages were ever present and tied directly to the behavior of smoking. In other contexts, reactance could drive message avoidance, perhaps when the behavior is not widely believed to be threatening or the message reflects a threat to personal freedom (e.g., "cigarette use by youth is dangerous and illegal"). Future studies should examine whether avoidance that is primarily driven by reactance is adaptive or maladaptive with respect to motivating healthier behavior. Qualitative research may shed light on

additional reasons why people avoid risk messages, as well as possible explanations for why avoidance is, at least sometimes, a marker for message effectiveness.

Finally, when countries enrich tobacco risk messages with aversive images, message avoidance tends to increase (Noar et al., 2016). This may lead policymakers and others to conclude that such warnings are ineffective because smokers are actively avoiding them, but our data suggest otherwise. After the implementation of plain packs and larger pictorial warnings in Australia in 2012, for example, smokers were more likely to place their packs face down and to conceal the new warnings using cases and other means (Zacher et al., 2014). Contrary to the conclusion that these actions render the warnings ineffective, the current study coupled with prior research suggests quite the opposite – that these actions are likely markers for the warnings having impact.

Limitations

Study strengths include the experimental design allowing us to assess the causal impact of risk messages on avoidance, the inclusion of diverse samples of smokers, and the use of a psychometrically strong measure of avoidance. However, Trial 1 examined the effect of adding pictorial warnings to cigarette packs, as well as implementing other label formatting changes required by the 2009 Tobacco Control Act, in comparison with the present text-only warnings in the U.S. While the trial aimed to compare the current warning policy to the new one in the Act, the use of this research design leaves open the possibility that the observed effects on mediators and avoidance may be due to the combination of adding pictures and other changes (e.g., location, size, and content). In Trial 2, the intervention labels were text-only on the side of the pack, and the mediators and outcome were assessed at the final study visit, making it harder to confirm the temporal ordering of effects. These differences between the trials may affect the comparability of the results. However, the similar findings between Trial 1 and Trial 2 suggest that the findings are robust to differences in the timing of measurement. The associations between the mediators and avoidance, as well as avoidance and forgoing or butting out cigarettes, were based on observational data, limiting our ability to assume causal associations between those variables. Thus, these correlational results should be interpreted with caution and future studies should attempt to replicate these results by experimentally manipulating the observed variables. Finally, these analyses were exploratory in nature; as such, we did not pre-register these hypotheses or analyses.

Conclusions

Policymakers should be encouraged by evidence showing that strong cigarette pack risk messages, especially those with images, are likely to encourage smoking cessation (Brewer et al., 2016; Cappella, 2016; Noar et al., 2016) and therefore prevent death and disease (Levy et al., 2016). However, some smokers will inevitably avoid health warnings and other cigarette pack risk messages. We argue that, in the context of cigarette pack risk messages, avoidance is not maladaptive defensive processing but instead a sign of *deeper* processing. In other words, smokers avoid warnings precisely *because* they are hard-hitting and elicit productive types of negative emotions, and avoidance is associated with more quitting-related behaviors. Future research should explore the role of avoidance of other types of

health communications – such as mass media campaigns – where risk messages are not displayed on the product itself, as well as among different populations (youth, non-smokers).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: Research reported in this publication was supported by The National Cancer Institute and FDA Center for Tobacco Products (CTP) under Award Numbers P30CA016086–38S2 and P50CA180907. T32-CA057726 from the National Cancer Institute of the National Institutes of Health supported MGH's time writing the paper. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the Food and Drug Administration.

References

- Bentler PM (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, 107, 238–246. [PubMed: 2320703]
- Blanton H, Snyder LB, Strauts E, & Larson JG (2014). Effect of graphic cigarette warnings on smoking intentions in young adults. *PloS One*, 9, e96315. [PubMed: 24806481]
- Bollen KA (1989). *Structural equations with latent variables*. New York, NY: Wiley.
- Borland R, Wilson N, Fong GT, Hammond D, Cummings KM, Yong H-H, et al. (2009a). Impact of graphic and text warnings on cigarette packs: findings from four countries over five years. *Tobacco Control*, 18, 358–364. [PubMed: 19561362]
- Borland R, Yong H, Wilson N, Fong GT, Hammond D, Cummings KM, et al. (2009b). How reactions to cigarette packet health warnings influence quitting: Findings from the ITC Four-Country survey. *Addiction*, 104, 669–675. [PubMed: 19215595]
- Brehm JW (1966). *A theory of psychological reactance*. New York, NY: Academic Press.
- Brehm JW, & Brehm SS (1981). *Psychological reactance: A theory of freedom and control*. New York, NY: Academic Press
- Brewer NT, Hall MG, Lee JG, Peebles K, Noar SM, & Ribisl KM (2015). Testing warning messages on smokers' cigarette packages: a standardised protocol. *Tobacco Control*, 25, 153–159. [PubMed: 25564282]
- Brewer NT, Hall MG, Noar SM, Parada H, Jr, Stein-Seroussi A, Bach LE, et al. (2016). Effect of pictorial cigarette pack warnings on changes in smoking behavior: A randomized clinical trial. *JAMA Intern Med*, 176, 905–912. [PubMed: 27273839]
- Brewer NT, Jeong M, Mendel JR, Hall MG, Zhang D, Parada H, Jr, et al. (2018a). Cigarette pack messages about toxic chemicals: a randomised clinical trial [published online ahead of print April 13, 2018]. *Tobacco Control*.
- Brewer NT, Parada JH, Hall MG, Boynton MH, Noar SM, & Ribisl KM (2018b). Understanding why pictorial cigarette pack warnings increase quit attempts [published online ahead of print May 29, 2018]. *Annals of Behavioral Medicine*.
- Brodar KE, Hall MG, Butler EN, Parada H, Stein-Seroussi A, Hanley S, et al. (2016). Recruiting diverse smokers: Enrollment yields and cost. *Int J Environ Res Public Health*, 13.
- Cappella JN (2016). Advancing the evidentiary base for tobacco warning labels: A commentary. *Social Science and Medicine*, 164, 130–132. [PubMed: 27492576]
- Cho YJ, Thrasher JF, Swayampakala K, Yong HH, McKeever R, Hammond D, et al. (2016). Does reactance against cigarette warning labels matter? Warning label responses and downstream smoking cessation amongst adult smokers in Australia, Canada, Mexico and the United States. *PloS One*, 11, e0159245. [PubMed: 27411100]
- Cho YJ, Thrasher JF, Yong HH, Szklo AS, O'Connor RJ, Bansal-Travers M, et al. (2018). Path analysis of warning label effects on negative emotions and quit attempts: A longitudinal study of

- smokers in Australia, Canada, Mexico, and the US. *Social Science and Medicine*, 197, 226–234. [PubMed: 29096946]
- DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, & Rossi JS (1991). The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *Journal of Consulting and Clinical Psychology*, 59, 295–304. [PubMed: 2030191]
- Emery LF, Romer D, Sheerin KM, Jamieson KH, & Peters E (2014). Affective and cognitive mediators of the impact of cigarette warning labels. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*, 16, 263–269. [PubMed: 23946325]
- Evans AT, Peters E, Strasser AA, Emery LF, Sheerin KM, & Romer D (2015). Graphic warning labels elicit affective and thoughtful responses from smokers: Results of a randomized clinical trial. *PloS One*, 10, e0142879. [PubMed: 26672982]
- Fathelrahman AI, Li L, Borland R, Yong HH, Omar M, Awang R, et al. (2013). Stronger pack warnings predict quitting more than weaker ones: finding from the ITC Malaysia and Thailand surveys. *Tobacco Induced Diseases*, 11, 20. [PubMed: 24330614]
- Hall MG, Marteau TM, Sunstein CR, Ribisl KM, Noar SM, Orlan EN, et al. (2018). Public support for pictorial warnings on cigarette packs: an experimental study of US smokers. *Journal of Behavioral Medicine*, 41, 398–405. [PubMed: 29411272]
- Hall MG, Sheeran P, Noar SM, Boynton MH, Ribisl KM, Parada H, Jr, et al. (2017a). Negative affect, message reactance and perceived risk: how do pictorial cigarette pack warnings change quit intentions? [published online ahead of print December 16 2017]. *Tobacco Control*.
- Hall MG, Sheeran P, Noar SM, Ribisl KM, Bach LE, & Brewer NT (2016). Reactance to health warnings scale: Development and validation. *Annals of Behavioral Medicine*, 50, 736–750. [PubMed: 27333895]
- Hall MG, Sheeran P, Noar SM, Ribisl KM, Boynton MH, & Brewer NT (2017b). A brief measure of reactance to health warnings. *Journal of Behavioral Medicine*, 40, 520–529. [PubMed: 28120228]
- Hammond D, Fong GT, McDonald PW, Brown KS, & Cameron R (2004). Graphic Canadian cigarette warning labels and adverse outcomes: Evidence from Canadian smokers. *American Journal of Public Health*, 94, 1442–1445. [PubMed: 15284057]
- Hammond D, Fong GT, McDonald PW, Cameron R, & Brown KS (2003). Impact of the graphic Canadian warning labels on adult smoking behaviour. *Tobacco Control*, 12, 391–395. [PubMed: 14660774]
- Hardcastle SJ, Chan DC, Caudwell KM, Sultan S, Cranwell J, Chatzisarantis NL, et al. (2015). Larger and more prominent graphic health warnings on plain-packaged tobacco products and avoidant responses in current smokers: A qualitative study. *International Journal of Behavioral Medicine*.
- Hayes AF (2009). *Beyond Baron and Kenny: Statistical mediation analysis in the new millennium*. *Commun Monogr*, 76, 408–420.
- International Tobacco Control Policy Evaluation Project. (2003). *4-Country W2 Replenishment Survey*.
- Keller PA, & Block LG (1996). Increasing the persuasiveness of fear appeals: The effect of arousal and elaboration. *Journal of Consumer Research*, 448–459.
- Kline RB (2011). *Principles and practice of structural equation modeling*. New York, NY: Guilford Press.
- Kowitz SD, Goldstein AO, Schmidt AM, Hall MG, & Brewer NT (2017). Attitudes toward FDA regulation of newly deemed tobacco products. *Tob Regul Sci*, 3, 504–515. [PubMed: 29450218]
- Levy DT, Mays D, Yuan Z, Hammond D, & Thrasher JF (2016). Public health benefits from pictorial health warnings on US cigarette packs: A SimSmoke simulation. *Tobacco Control*.
- Li L, Borland R, Fong GT, Jiang Y, Yang Y, Wang L, et al. (2014). Smoking-related thoughts and microbehaviours, and their predictive power for quitting: findings from the International Tobacco Control (ITC) China Survey. *Tobacco Control*, Advance online publication.
- Maynard OM, Attwood A, O'Brien L, Brooks S, Hedge C, Leonards U, et al. (2014). Avoidance of cigarette pack health warnings among regular cigarette smokers. *Drug and Alcohol Dependence*, 136, 170–174. [PubMed: 24485554]

- McCaul KD, Branstetter AD, Schroeder DM, & Glasgow RE (1996). What is the relationship between breast cancer risk and mammography screening? A meta-analytic review. *Health Psychology*, 15, 423–429. [PubMed: 8973921]
- McCloud RF, Okechukwu C, Sorensen G, & Viswanath K (2017). Cigarette graphic health warning labels and information avoidance among individuals from low socioeconomic position in the U.S. *Cancer Causes and Control*, 28, 351–360. [PubMed: 28255678]
- McQueen A, Caburnay C, Kaphingst K, Sanders-Thompson V, Tovar M, Waters E, et al. (2013). What are the reactions of diverse US smokers when graphic warning labels are affixed to their cigarette packs? American Society of Preventive Oncology Conference Memphis, TN.
- McQueen A, Kreuter MW, Boyum S, Thompson VS, Caburnay CA, Waters EA, et al. (2015). Reactions to FDA-proposed graphic warning labels affixed to U.S. smokers' cigarette packs. *Nicotine Tob Res*, 17, 784–795. [PubMed: 25589676]
- Morgan JC, Southwell BG, Noar SM, Ribisl KM, Golden SD, & Brewer NT (2017). Frequency and content of conversations about pictorial warnings on cigarette packs. *Nicotine Tob Res*, 20, 882–887.
- Noar SM, Francis DB, Bridges C, Sontag JM, Brewer NT, & Ribisl KM (2017). Effects of strengthening cigarette pack warnings on attention and message processing: A systematic review. *Journalism & Mass Communication Quarterly*, 94, 416–442. [PubMed: 29975497]
- Noar SM, Francis DB, Bridges C, Sontag JM, Ribisl KM, & Brewer NT (2016). The impact of strengthening cigarette pack warnings: Systematic review of longitudinal observational studies. *Social Science and Medicine*, 164, 118–129. [PubMed: 27423739]
- Nonnemaker J, Farrelly M, Kamyab K, Busey A, & Mann N (2010). Experimental study of graphic cigarette warning labels: Final results report. Research Triangle Park, NC: RTI International.
- Osman A, Thrasher JF, Yong HH, Arillo-Santillan E, & Hammond D (2017). Disparagement of health warning labels on cigarette packages and cessation attempts: results from four countries. *Health Education Research*, 32, 524–536. [PubMed: 29136169]
- Parada H, Jr., Hall MG, Boynton MH, & Brewer NT (2017). Trajectories of responses to pictorial cigarette pack warnings. *Nicotine Tob Res*.
- Partos TR, Borland R, Thrasher JF, Li L, Yong HH, O'Connor RJ, et al. (2014). The predictive utility of micro indicators of concern about smoking: findings from the International Tobacco Control Four Country study. *Addictive Behaviors*, 39, 1235–1242. [PubMed: 24813549]
- Peters C, & Enders C (2002). A primer for the estimation of structural equation models in the presence of missing data: Maximum likelihood algorithms. *Journal of Targeting, Measurement and Analysis for Marketing*, 11, 81–95.
- Peters E, Evans AT, Hemmerich N, & Berman M (2016). Emotion in the law and the lab: The case of graphic cigarette warnings. *Tob Regul Sci*, 2, 404–413. [PubMed: 29057296]
- Peters E, Romer D, & Evans A (2014). Reactive and thoughtful processing of graphic warnings: Multiple roles for affect University of North Carolina, Lineberger Comprehensive Cancer Center, Center for Regulatory Research on Tobacco Communication. Chapel Hill, NC.
- Population Assessment of Tobacco and Health Study. (2018). PATH: Population Assessment of Tobacco and Health.
- Prochaska JO, & DiClemente CC (1986). Toward a comprehensive model of change In Miller WR, & Heather N (Eds.), *Treating addictive behaviors* pp. 3–27). New York: Plenum Press.
- Sheeran P, Harris PR, & Epton T (2014). Does heightening risk appraisals change people's intentions and behavior? A meta-analysis of experimental studies. *Psychological Bulletin*, 140, 511–543. [PubMed: 23731175]
- Sidi Y, & Harel O (2018). The treatment of incomplete data: Reporting, analysis, reproducibility, and replicability. *Social Science and Medicine*.
- Steiger JH (1990). Structural model evaluation and modification: An interval estimation approach. *Multivariate Behav Res*, 25, 173–180. [PubMed: 26794479]
- Thrasher JF, Swayampakala K, Borland R, Nagelhout G, Yong HH, Hammond D, et al. (2016). Influences of self-efficacy, response efficacy, and reactance on responses to cigarette health warnings: A longitudinal study of adult smokers in australia and canada. *Health Commun*, 31, 1517–1526. [PubMed: 27135826]

- Tucker LR, & Lewis C (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38, 1–10.
- Wardle H, Pickup D, Lee L, Hall J, Pickering K, Grieg K, et al. (2010). Evaluating the impact of picture health warnings on cigarette packets. London, UK: Public Health Research Consortium.
- Watson D, Clark LA, & Tellegen A (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology*, 54, 1063. [PubMed: 3397865]
- Wegner DM (1994). Ironic processes of mental control. *Psychological Review*, 101, 34–52. [PubMed: 8121959]
- Witte K (1992). Putting the fear back into fear appeals: The extended parallel process model. *Commun Monogr*, 59, 329–349.
- World Health Organization (2012). WHO global report: Mortality attributable to tobacco. Geneva, Switzerland
- Yong HH, Borland R, Thrasher JF, Thompson ME, Nagelhout GE, Fong GT, et al. (2014). Mediation pathways of the impact of cigarette warning labels on quit attempts. *Health Psychology*, 33, 1410–1420. [PubMed: 24977309]
- Yong HH, Fong GT, Driezen P, Borland R, Quah AC, Sirirassamee B, et al. (2013). Adult smokers' reactions to pictorial health warning labels on cigarette packs in Thailand and moderating effects of type of cigarette smoked: findings from the international tobacco control southeast Asia survey. *Nicotine and Tobacco Research*, 15, 1339–1347. [PubMed: 23291637]
- Zacher M, Bayly M, Brennan E, Dono J, Miller C, Durkin S, et al. (2014). Personal tobacco pack display before and after the introduction of plain packaging with larger pictorial health warnings in Australia: an observational study of outdoor cafe strips. *Addiction*, 109, 653–662. [PubMed: 24428427]
- Zhang JY, Borland R, Coghil K, Petrovic-Lazarevic S, Young D, Yeh CH, et al. (2011). Evaluating the effect of health warnings in influencing Australian smokers' psychosocial and quitting behaviours using fuzzy causal network. *Expert Syst Appl*, 38, 6430–6438. [PubMed: 26560158]

Research highlights

- We explored why smokers avoid cigarette pack risk messages in two large trials.
- Smokers avoided risk messages because the messages made them feel negative emotions including fear.
- These negative emotions are a sign of an adaptive response to cigarette pack risk messages.
- Reactance was not the main reason why smokers avoided messages.
- Avoidance was also associated with more forgoing or butting out of cigarettes.

A. Intervention arm **B. Control arm****SURGEON GENERAL'S WARNING:**

Cigarette Smoke Contains
Carbon Monoxide.

SURGEON GENERAL'S WARNING:

Quitting Smoking Now Greatly Reduces
Serious Risks to Your Health.

SURGEON GENERAL'S WARNING:

Smoking by Pregnant Women May Result
in Fetal Injury, Premature Birth, and
Low Birth Weight.

SURGEON GENERAL'S WARNING:

Smoking Causes Lung Cancer, Heart Disease,
Emphysema, and May Complicate Pregnancy.

Figure 1.

Labels placed on smokers' cigarette packs in intervention arm (Panel A) and control arm (Panel B) in Trial 1.

Figure 2.

Labels placed on smokers' cigarette packs in intervention arm (Panel A) and control arm (Panel B) in Trial 2.

A. Trial 1 **B. Trial 2** **Figure 3.**

Structural equation models assessing the impact of intervention messages on message avoidance in Trial 1 ($n=2,149$, Panel A) and Trial 2 ($n=719$, Panel B). Values in parentheses show bivariate association between intervention messages and avoidance. $*p<.05$. $**p<.001$.

Table 1.

Participant characteristics.

	Trial 1 (n=2,149)				Trial 2 (n=719)			
	Control (n=1,078)		Intervention (n=1,071)		Control (n=359)		Intervention (n=360)	
	n	(%)	n	(%)	n	(%)	n	(%)
Age								
18–20 years	48	(4.5)	51	(4.9)	-	-	-	-
21–29 years	268	(25.2)	278	(26.5)	81	(22.6)	83	(23.1)
30–39 years	232	(21.9)	221	(21.1)	86	(24.0)	78	(21.7)
40–49 years	211	(19.9)	188	(17.9)	59	(16.4)	75	(20.8)
50–59 years	229	(21.6)	229	(21.8)	90	(25.1)	91	(25.3)
60+ years	74	(7.0)	82	(7.8)	43	(12.0)	33	(9.2)
Mean (SD)	39.7	(13.4)	39.8	(13.7)	42.8	(13.6)	42.1	(13.2)
Gender								
Male	507	(47.4)	532	(50.1)	172	(47.9)	196	(54.4)
Female	548	(51.2)	512	(48.2)	169	(47.1)	151	(41.9)
Transgender ^a	15	(1.4)	19	(1.8)	18	(5.0)	12	(3.6)
Gay, lesbian, or bisexual	173	(16.3)	195	(18.8)	88	(24.5)	93	(25.8)
Hispanic	92	(8.6)	89	(8.5)	56	(15.6)	44	(12.2)
Race								
American Indian or Alaska Native	7	(.7)	11	(1.1)	17	(4.7)	17	(4.7)
Asian	28	(2.7)	42	(4.0)	29	(8.1)	31	(8.6)
Black or African American	484	(45.8)	510	(48.9)	124	(34.5)	133	(36.9)
Native Hawaiian/other Pacific Islander	11	(1.0)	6	(.6)	11	(3.1)	12	(3.3)
Other/multiracial	134	(12.7)	117	(11.2)	42	(11.7)	35	(9.7)
White	393	(37.2)	358	(34.3)	136	(37.9)	132	(36.7)
Education								
High school graduate or less	333	(31.1)	344	(32.5)	67	(18.7)	89	(24.7)
Some college	519	(48.5)	502	(47.4)	151	(42.1)	124	(34.4)
College graduate	156	(14.6)	156	(14.7)	114	(31.8)	122	(33.9)
Graduate degree	63	(5.9)	58	(5.5)	27	(7.5)	25	(6.9)
Low income ^b	570	(53.0)	589	(55.3)	197	(54.9)	216	(60.0)
Household income, annual								
\$0-\$24,999	566	(53.3)	589	(55.8)	154	(42.9)	170	(47.2)
\$25,000-\$49,999	272	(25.6)	266	(25.2)	88	(24.5)	89	(24.7)
\$50,000-\$74,999	110	(10.4)	92	(8.7)	53	(14.8)	35	(9.7)
\$75,000+	115	(10.8)	109	(10.3)	64	(17.8)	66	(18.3)
Trial site								
California	594	(55.1)	592	(55.3)	359	(100.0)	360	(100.0)
North Carolina	484	(44.9)	479	(44.7)	-	-	-	-
Cigarettes smoked per day, mean (SD)	8.8	(6.6)	8.7	(7.3)	9.97	(12.2)	11.62	(16.9)

	Trial 1 (n=2,149)				Trial 2 (n=719)			
	<u>Control (n=1,078)</u>		<u>Intervention (n=1,071)</u>		<u>Control (n=359)</u>		<u>Intervention (n=360)</u>	
	<i>n</i>	(%)	<i>n</i>	(%)	<i>n</i>	(%)	<i>n</i>	(%)
Quit intentions, mean (SD)	2.2	(.9)	2.3	(.9)	2.4	(1.0)	2.4	(0.9)

Note. Characteristics did not differ by trial arm in either trial (all $p > .05$). - not applicable.

^aFor Trial 2, includes other gender identity

^bTrial 1 150% of federal poverty level. Trial 2 200% federal poverty level.

Table 2.Latent variables used in the measurement and structural equation models (Trial 1 $n=2,148$ /Trial 2 $n=704$)

Latent variable Trial 1/Trial 2	Indicator item wording [response scale]	Factor loading Trial 1/Trial 2
Negative affect ($\alpha=.92/.91$)	How much did the warning on your cigarette packs make you feel...	
	Anxious?	.86/.85
	Disgusted?	.88/.88
	Guilty?	.88/.87
	Sad?	.85/.86
	Scared?	.93/.87
	[not at all (coded as 1), a little (2), somewhat (3), very (4), extremely (5)]	
Message reactance ($\alpha=.75/.78$)	Say how much you agree or disagree with each statement below about the warning we put on your packs.	
	This warning annoys me.	.79/.89
	This warning is trying to manipulate me.	.79/.66
	The health effect on this warning is overblown. ^a	.77/.80
	[strongly disagree (1), somewhat disagree (2), neither agree nor disagree (3), somewhat agree (4), strongly agree (5)]	
Message avoidance ($\alpha=.90/.83$)	In the last week...	
	How often have you tried to avoid looking at the warning label on your cigarette packs?	.98/.92
	In the last week, how often have you tried to avoid thinking about the warning label on your cigarette packs?	.95/.89
	In the last week, how often have you put your cigarettes away because you didn't want others to see the warning label on the pack?	.83/.81
	[never (1), rarely (2), sometimes (3), often (4), all of the time (5)]	

Note. Analyses excluded $n=1$ Trial 1 and $n=15$ Trial 2 participants with missing data on all items in the model. Survey items in the table are from Trial 1; Trial 2 used identical items but used the word “labels” instead of “warning” or “warning label.” In Trial 1, the correlation of negative affect with message reactance was .05 ($p=.07$), negative affect with message avoidance was .54 ($p<.001$), and message reactance with message avoidance was .18 ($p<.001$). In Trial 2, the correlation of negative affect with message reactance was .01 ($p=.90$), negative affect with message avoidance was .66 ($p<.001$), and message reactance with message avoidance was .26 ($p<.001$).

^a Trial 2 wording: “The labels are overblown.”

Table 3.Multiple mediation of intervention messages' impact on message avoidance (Trial 1 $n=2,149$ /Trial 2 $n=719$)

Mediator variable	<i>a</i> pathway		<i>b</i> pathway		Mediated effect	
	βa	<i>p</i>	βb	<i>p</i>	$\beta a * \beta b$ (95% CI)	<i>p</i>
Negative affect						
Trial 1	.41	<.001	.51	<.001	.21 (.18, .24)	<.001
Trial 2	.18	<.001	.68	<.001	.12 (.07, .17)	<.001
Message reactance						
Trial 1	.24	<.001	.10	<.05	.03 (.01, .04)	<.001
Trial 2	.02	.64	.24	<.001	.01 (-.02, .03)	.65

Note. Table reports standardized path coefficients and mediated effects.

^a pathways are the associations between intervention message exposure and the mediators.

^b pathways are the associations between the mediators and message avoidance, controlling for intervention message exposure and the other mediators. Analyses excluded $n=1$ Trial 1 and $n=15$ Trial 2 participants with missing data on all items in the model.