the blood stream, or split up by hydrolysis into bodies of smaller molecular weight. The authorities whom I have consulted think that the precipitate would favour the coagulation of the blood. The clinical experience of surgeons and post-mortem examinations would help to settle this question. I venture in conclusion to suggest that surgeons should introduce zinc wire into an aneurism which they may wish to treat electrolytically rather than the wire recommended by Dr Stewart or other clinical authorities who have not tested the coagulating effects of these metals experimentally in the physical laboratory, for my experiments, which confirm those of Leduc, convince me that the passage of an electric current through an aneurism in the orthodox Stewart-Moore-Corradi method in no way aids in bringing about the coagulation of the blood: whatever coagulation is produced is due only to the introduction of the foreign body.

3. CHRONIC INFLAMMATORY OEDEMA OF THE SUB-MUCOUS TISSUES OF THE NOSE

By J. S. FRASER, M.B., CH.B., F.R.C.S.Ed.,
Assistant Surgeon, Ear and Throat Department, Royal Infirmary, Edinburgh

FORMATION AND FUNCTIONS OF THE NOSE

THE nasal septum is seldom or never absolutely straight, in fact a straight septum is an abnormally normal one; the outer wall of the nose is thrown into convolutions by the turbinals so as to expose as much secreting surface as possible in the small space available. The inspired air entering the nasal chambers thus strikes against the septum or turbinals before passing back to the naso-pharynx, and is warmed, moistened, and filtered to some extent from dust particles and bacteria before it reaches the cavities of the pharynx and larynx. The nasal mucous membrane is likely, therefore, to be a very active structure; it is well supplied with blood-vessels, and in certain parts, notably in the inferior turbinal region, these vessels take part in the formation of erectile tissue; the size of the nasal cavities can thus be regulated according to the state of the air entering the nose; the inferior turbinals swell up in cold weather or in dusty atmospheres; excitement also influences the state of engorgement of the turbinals, and the well-known phrase, "open-mouthed with astonishment," would be more accurately replaced by "shut-nosed with astonishment." The size of the inferior turbinals may also be seen to vary during the short period covered

by an examination of the anterior nares, and we are probably all familiar with the fact that if we lie on one side in bed the nostril on that side tends to become blocked.

METHOD OF CARRYING OUT THE PRESENT INVESTIGATION

The number of specimens examined was 70; of these 22 were inferior turbinals, of which 6 were normal, 6 in a condition of slight œdema, 5 markedly œdematous, and 5 in the condition known as "cauliflower hypertrophy." This state, however, is merely one of very extreme œdematous infiltration. Eighteen specimens of the middle turbinal were examined, in addition to 30 which formed the subject (1) of a previous paper. Of the middle turbinals 3 were normal, 10 showed slight œdema, and the rest marked œdematous hypertrophy, or, in other words, nasal polypus formation. In all but a few cases the tissues were removed during life, washed in water to remove blood and mucus from the surface, and placed at once in 5 per cent. nitric acid in spirit. After a week they were transferred to 10 per cent. nitric acid in spirit and changed three or four times according to the size of the specimen. The time required for decalcification varied in specimens which contained apparently an equal amount of bone. The tissues were carried through in the usual way and embedded in paraffin and cut often in serial sections. It was found by no means easy to get good sections of tissues of such varying degrees of density as the turbinals, containing bone, periosteum, loose connective tissue glands, and delicate mucous membrane. The staining method used was the ordinary one—hæmatoxylin and eosin, but Unna's orcein method was used in addition in almost all cases for elastic fibres.

SITES OF CHRONIC INFLAMMATORY ŒDEMA IN THE NOSE

(A) *The Septum Nasi*.—(1) Anteriorly in the region of the tubercle; (2) posteriorly. Swelling is often seen by posterior rhinoscopy, apparently in the region of the posterior border of the vomer high up towards the roof of the choanæ, giving rise to an appearance that may be compared to the "ace of hearts." This is probably merely an extension backwards of the œdema in the region of the tubercle.

(B) *Middle Turbinal*.—The anterior end, lower border, or outer surface are most frequently affected; these parts are the most dependent.

(c) *Inferior Turbinal*.—The anterior end is affected rather more often than the posterior end, but frequently the anterior end, inner surface, and lower border, as well as the posterior end, are all affected.

NORMAL STRUCTURE OF INFERIOR AND MIDDLE TURBINAL BODIES

Inferior Turbinal.—(Fig. 1). This consists of a bony framework with an irregular surface; within this we find so-called marrow spaces lined by delicate connective tissue and containing a central thin-walled blood-vessel. The periosteum consists of two or three layers of elongated cells, and just outside this there is a dense layer of elastic tissue from which strands of elastic fibres run out radially in all directions towards the mucous membrane (Fig. 7). Beyond

FIG. A

the periosteum we have a layer of fairly dense connective tissue containing the larger blood-vessels and some unstripped muscle-fibres, and outside this again the erectile tissue-layer made up of thin-walled blood-vessels of irregular size and shape; the erectile tissue is better marked at both ends and along the inferior border of the turbinal. Just under the superficial epithelium and its basement membrane we have the racemose mucous glands (Fig. 2), the ducts of which open through the basement membrane usually into smaller hollows on the surface. Immediately under the basement membrane there is a thin layer of elastic fibres running parallel to the surface but connected with the fibres that run out radially from the periosteum; the whole turbinal has thus a framework of elastic fibres which causes it to return to its normal size when vascular engorgement is over. The covering layer consists of ciliated columnar epithelium, beneath which are several layers of cubical or irregular cells situated upon the basement membrane.

Blood supply: The larger vessels run near the periosteum more or less parallel to the long axis of the turbinal; from these smaller vessels run radially outwards to the blood sinuses and to the mucous

CHANGES IN THE INFERIOR TURBINAL.

FIG. 1.—NORMAL INFERIOR TURBINAL: the bony core of the turbinal is over-decalcified, but the figure shows the large blood-vessels in the periosteal layer, the blood spaces best marked along the lower border and inner surface, the numerous glands just under the basement membrane, and the superficial epithelium (absent in parts). Note the small cell infiltration of superficial layers most marked on inner surface.
× 6 diam.

FIG. 2.—NORMAL MUCOUS MEMBRANE of inferior turbinal. Note superficial epithelium containing what appear to be "intra-epithelial" glands; this appearance may be due to oblique section of a gland duct; small cell infiltration (1) of superficial layers, (2) between gland acini.
× 50 diam.

Inner surface, showing
slight œdema

* Normal bone

FIG. 3.—EARLY INFLAMMATORY œDEMA of the inferior turbinal; superficial epithelium is absent; note that the glands are slightly pushed inwards towards the bony core of the turbinal. The œdema of the septal surface of the turbinal is shown when contrasted with outer surface where the tissues are almost normal.

× 6 diam.

FIG. 4.—Marked chronic inflammatory œdema of the inferior turbinal; "cauliflower hypertrophy"; the glands markedly displaced inwards; dilated ducts are seen opening into the hollows between the papillae in the lower part of the section; the blood spaces are few and small; the bony core of the turbinal is, however, quite healthy; the walls of the blood-vessels appear to be considerably thickened.

× 6 diam.

FIG. 5.—Marked chronic inflammatory œdema of posterior end of inferior turbinal. Notice œdema of superficial layers and displacement of glands and blood spaces.

× 6 diam.

In all cases the microscopic section has been made in the coronal plane.

PLATE VII.—TO ILLUSTRATE DR J. S. FRASER'S PAPER.

CHANGES IN THE MIDDLE TURBINAL.

FIG. 6.—Normal middle turbinal. Note the more delicate structure of the middle turbinal ; the glands are numerous and quite near to the surface ; the blood spaces are small and fewer than in the inferior turbinal.

× 6 diam.

FIG. 7.—Elastic tissue of the middle turbinal. Around the perosteum the fibres form a dense network from which more delicate strands radiate out into the substance of the turbinal surrounding the blood spaces and glands ; just under the basement membrane there is a delicate layer connected to the radiating fibres.

× 50 diam.

FIG. 8.—Early CHRONIC INFLAMMATORY ŒDEMA of the middle turbinal, or POLYPOID middle turbinal. Note large connective-tissue spaces containing serum along the lower border of the turbinal; the glands in this region are displaced inwards; the bone is healthy.
 × 6 diam.

FIG. 9.—Higher power view of same specimen as shown in Fig. 8. Note glands pushed inwards by œdema of superficial layers.
 × 50 diam.

In all cases the microscopic section has been made in the coronal plane.

PLATE VIII.—TO ILLUSTRATE DR J. S. FRASER'S PAPER.

CHANGES IN THE MIDDLE TURBINAL—*Continued.*

FIG. 10.—Slight chronic inflammatory oedema of the middle turbinal; slight dilatation of gland ducts; inward displacement of glands and blood spaces. Note marked small cell infiltration of superficial layers.

FIG. 11.—Edema of middle turbinal in case of ETHMOIDAL SUPPURATION. The normal tissues are crowded towards the bone; the thickened lining membrane of the cell in the middle turbinal is well seen, and the small cell infiltration of its mucous membrane. × 6 diam.

FIG. 12.—This section shows the appearance of the tissues at the base of the polypoid swelling. Note the dilatation of the lymphatic vessels and spaces. In the centre of the field a blood-vessel is seen containing organised tissue—probably a thrombus.
 × 50 diam.

FIG. 13.—Section of TYPICAL "NASAL POLYPUS," showing large connective-tissue spaces filled with serum, dilated gland ducts and small blood-vessels; the superficial epithelium is almost normal.
 × 6 diam.

In all cases the microscopic section has been made in the coronal plane.

PLATE IX.—TO ILLUSTRATE DR J. S. FRASER'S PAPER.

glands, and, finally, very small vessels ramify just under the basement membrane.

Normally in living people there is always a certain amount of small-cell infiltration under the basement membrane and around the glands and superficial blood-vessels; it is only in the unborn that this is not found (Kubo).

Middle Turbinal.—(Fig. 6.) This is much the same as that of the inferior turbinals, but with the following differences: the bone is thinner and altogether more delicate; not infrequently there is a definite cell in the middle turbinal lined with ciliated epithelium, in fact, in these cases the middle turbinal contains one of the ethmoidal cells. The erectile tissue is not so marked in the middle turbinal as in the inferior, and the whole structure is looser and more delicate. From an anatomical point of view we must remember that the middle turbinal is situated higher up in the nose than the inferior.

MICROSCOPICAL APPEARANCES IN CASES OF ŒDEMA OF THE MIDDLE AND INFERIOR TURBINALS

(a) *The Inferior Turbinal.*—After writing a paper on “The Ætiology of Nasal Polypus,” it occurred to me that further light might be thrown on this question by the examination of the inferior turbinal bodies, and I was not surprised to find that the changes leading to so-called hypertrophy of the *inferior* turbinals were exactly similar to those seen in the middle turbinals in cases of nasal polypus formation.

It is necessary in the first place to distinguish between engorgement of the inferior turbinals due to distension of the blood spaces (cavernous tissue) and œdema of the submucous tissue (so-called hypertrophy of the inferior turbinals). This can readily be done by the application of cocaine to the turbinal; vascular engorgement goes down within a few seconds, whilst œdema is little, if at all, affected. In some cases both conditions are present, and, on the application of cocaine, the turbinal shrinks to some extent, but still remains considerably enlarged. It is extremely probable, however, that chronic engorgement is the first stage of inflammation, leading to œdematous infiltration when changes have occurred in the walls of the blood-vessels. In early cases of œdema of the inferior turbinal, the process commences just under the basement membrane, which is frequently thickened (Fig. 3). Kubo considers this thickening to mean inflammation; the leucocyte infiltration is increased,

especially round the vessels and glands, and, in some cases, amounts almost to the formation of lymph nodules. The glands are slightly pushed inwards by the œdema and the ducts dilated; the blood spaces, periosteum, bone, and marrow are not affected.

As the œdema increases (Fig. 4) the glands and blood spaces are pushed further back, and the surface of the turbinal tends to become crenated or scalloped, as seen in sections—papillary, as seen on surface view. This is due to the gland ducts tacking down the mucous membrane at various spots, and can be well seen in the posterior ends of the inferior turbinals by posterior rhinoscopy in cases of so-called “mulberry hypertrophy.”

In the œdematous submucous tissue there are a large number of cells showing myxomatous degeneration, and it is due to this fact that nasal polypi were so long regarded as myxomata.

Even in the most marked conditions of œdema of the inferior turbinal—the so-called “cauliflower hypertrophy”—the surface epithelium is but little changed (Fig. 5) and the cilia are well preserved; here and there leucocytes may be seen making their way through the superficial epithelium to the surface; this fact accounts for the muco-purulent discharge present in so many of these cases.

If we examine one of the papillæ on the surface of such an inferior turbinal under a high power, we are at once struck by the close resemblance to a nasal polypus.

The changes in the inferior turbinal are more marked on the inner surface, *i.e.* the surface next the septum over which the air current passes (Fig. 3).

In advanced cases there may be some thickening of the periosteum, but the bone and marrow spaces are always normal so far as my observations go. Finally, I have noticed that in these cases the blood spaces are usually small; I have remarked before that they are pressed towards the bony centre of the turbinal by the œdema which occurs between them and the mucous membrane, and this fact probably accounts for their diminution in size.

(b) *Middle Turbinal*.—The first change noted is œdema of the connective-tissue layer just under the basement membrane (Fig. 8); the connective-tissue spaces are enlarged and contain serum, and the glands and blood spaces are gradually displaced inwards towards the bony centre of the turbinal as the œdema increases. The epithelial covering of the turbinal, along with the basement membrane, is of course pushed away from the bony framework,

and later on comes to hang down in the nasal cavity ; this is the condition we recognise clinically as " polypoid middle turbinal." The ducts of the glands in many cases become obstructed—on account of the œdematous infiltration—and in this way large cystic spaces are developed.

It is very difficult to speak with certainty as to the condition of the mucous membrane ; during the course of hardening, decalcification, embedding, or staining, it often happens that the surface epithelium gets separated and lost ; but it is also remarkable that in cases of marked nasal polypus formation the superficial epithelium is apparently quite normal.

CAUSES OF ŒDEMA

Without doubt the two most common pathological conditions affecting respectively the middle and inferior turbinals are nasal polypus and inferior turbinal hypertrophy (so called). Differences in the macro- and microscopic appearances are due to, *firstly*, difference in the structure of the two turbinals, and, *secondly*, difference in anatomical position, or, in other words, the middle turbinal is of looser and more delicate structure and situated further from the floor of the nose.

In both cases the cause of the œdema is to be found in repeated or severe attacks of acute catarrh (coryza) : in this condition the turbinals are hyperæmic and slightly œdematous, but as a rule resolution occurs and the parts return to normal : on the other hand in a few cases the inflammation becomes chronic, and the deeper layers of the mucous membrane are involved. Clinically, these chronic cases are characterised by constant nasal discharge (mucous or muco-purulent) and nasal obstruction ; the patients complain that they are never free from a " cold " in the head, and they usually state that they have suffered from nasal discharge and obstruction for several years. The tissues of the nose are thus constantly soaking in their own secretions, and a return to normal is prevented. It has been pointed out to me that an analogy may be found in the case of papilloma of the bladder : this condition is so serious because it is impossible to keep the parts dry, whereas warts on the hand tend to shrivel up and disappear when moisture is excluded.

On going over the cases of inferior turbinal enlargement which presented themselves at Dr Logan Turner's clinic during the year

1907, I found that the actual number was 242, but the great majority of these were cases of vascular engorgement. In most of the cases there was only slight enlargement, but it was noted that in 30 there was a co-existing polypoid condition of the middle turbinals, and probably this number would have been greater had cocaine been applied in all instances to the inferior turbinals so as to obtain a better view of the middle turbinal region. Of the 79 cases classified under the head of nasal polypus, there were 20 with co-existing enlargement of the inferior turbinal.

FORMATION OF POLYPI

Oedema occurs beneath the mucous membrane along the lower border or outer surface of the turbinal: the mucous membrane thus sags down in the nose, and in advanced cases reaches the floor of the nasal cavities, or even protrudes from the anterior or posterior nares.

Before this stage is reached, however, the developing nasal

FIG. B.

polypus is acted upon by the respiratory air currents, and its base of attachment is specially dragged on in the act of blowing the nose: gravity also tends to increase the formation of the oedematous swelling once it has begun. From what has been said it is not difficult to understand that in some cases the base becomes constricted so that a pear-shaped swelling results, but in my opinion the genuine pear-shaped polypus is much less frequent than is generally believed. I hold that the loop of the snare is frequently responsible for the pear shape, and that in most cases the polypus is broad based along the lower border or outer surface of the turbinal. If we think of the frequency of oedema in inflammation of such loose structures as the prepuce, scrotum, and lower eyelid, we need not wonder that it occurs in the chronic inflammation of the middle turbinal region of the nose. In the inferior turbinal the case is somewhat different; here the tissue is denser, and the lower border

of the turbinal is already almost on the floor of the nasal chamber ; consequently we do not meet with such marked œdematous swelling. At the *posterior* end of the inferior turbinal we may, however, in advanced cases, see a large papillary mass of œdematous tissue hanging down in the posterior surface of the soft palate.

Inflammation in any situation is always associated with exudation, and the looser the tissue the greater the amount of exudation. I believe that in early cases of rhinitis we have an excessive amount of vascular turgescence associated with a slight degree of œdematous infiltration ; this goes down under cocaine application. If the inflammation becomes chronic the œdema increases most probably on account of changes in the walls of the blood-vessels and blood spaces. Lymphatic and venous obstruction no doubt play a part, and I have observed swelling of the endothelial cells lining the lymphatic vessels in one or two of my specimens. Watson Williams (3) recently showed a specimen of nasal polypus in which a lymphatic vessel appeared to be blocked by inflammatory products. Hopman (4) thinks that the cause of the œdema is some impairment of the circulation in the efferent venous vessels, and the engorged state of the small veins in many of the specimens examined by me supports this theory (Fig. 12). Grünwald (5) explains the œdema by changes in the vessels of the nature of peri- and endo-arteritis, but one is bound to note that even in apparently normal turbinals the vessel walls seem to be abnormally thick. Goodale (6) says, " It is possible that the inflammatory infiltration which penetrates the substance of the turbinals may lead in some places to a stasis from compression of the veins " ; the same writer entirely disagrees with the bone disease theory of nasal polypus formation.

In advanced cases the elastic fibres are hard to find—at least in the œdematous parts ; near the periosteum they are present and normal in appearance. The striking similarity of the changes in the middle and inferior turbinals will be apparent from the illustrations, and nobody has yet said that inferior turbinal œdema or hypertrophy is due to bone disease ; we may well ask why œdema of the middle turbinal should be due to this cause. It is true that the bone is often affected in advanced cases of nasal polypus formation, associated with ethmoidal suppuration, but in this case the bone disease is *the effect*, not the cause (Fig. 11). Yonge's (7) theory as to the causation of nasal polypi was not well received at the last meeting of the British Medical Association at Exeter, and as I have dealt with it in a previous paper I need not do so again.

In regard to Grünwald's theory that nasal polypi are associated with sinus suppuration, I can only state that this appears to be true in only 50 per cent. of cases. I think we are not far wrong if we regard the character of the discharge present as diagnostic—that is to say, if nasal polypi are found associated with a clear watery discharge no sinus suppuration is present; if the discharge is muco-purulent, sinus suppuration may or may not be present; if the discharge is distinctly purulent then sinus suppuration is present.

Morell Mackenzie used to urge against the inflammatory origin of nasal polypi that while nasal catarrh is very frequent in children, polypi are extremely rare. There are two good answers to this objection: (1) the middle turbinal region is not easy to examine in children, and minor degrees of œdema may easily pass unnoticed; and (2) it is extremely probable that the vascular changes leading up to marked polypus formation take years to occur—in other words, children have ceased to be children when polypi are developed.

In conclusion, I wish to express my thanks to Dr Logan Turner, who has kindly allowed me to make use of his clinical material, to the Royal College of Physicians (Edinburgh) for permission to work in their Laboratory, and to Dr James Ritchie for much advice and assistance in carrying out the work.

REFERENCES.

- (1) *Scot. Med. and Surg. Journ.*, April 1907.
- (2) *Archiv. f. Laryngologie*, Bd. xix. S. 202.
- (3) *Proceedings of the Royal Society of Medicine*, 1908.
- (4) *Wiener med. Presse*, 1883.
- (5) "Nasal Suppuration," 1900.
- (6) "Diseases of Eye, Ear, Nose and Throat," by Posey and Wright, p. 706.
- (7) "Polypus of the Nose," 1906.

Meeting X.—July 1, 1908

Dr JAMES RITCHIE, *President, in the Chair*

I. EXHIBITION OF PATIENTS

Dr Byrom Bramwell exhibited (a case of OSTEITIS DEFORMANS (Paget's disease)). The disease is very rare in Scotland. Osler states (in his last edition, published in 1905) that only about sixty-seven typical cases have been recorded. The notes are as follows:—