The informational entropy endowed in cortical oscillations

Arturo Tozzi¹ · James F. Peters^{2,3} · Mehmet Niyazi Çankaya⁴

Received: 8 March 2018 / Revised: 31 May 2018 / Accepted: 14 June 2018 / Published online: 18 June 2018
© Springer Nature B.V. 2018

Abstract

A two-dimensional shadow may encompass more information than its corresponding three-dimensional object. Indeed, if we rotate the object, we achieve a pool of observed shadows from different angulations, gradients, shapes and variable length contours that make it possible for us to increase our available information. Starting from this simple observation, we show how informational entropies might turn out to be useful in the evaluation of scale-free dynamics in the brain. Indeed, brain activity exhibits a scale-free distribution that leads to the variations in the power law exponent typical of different functional neurophysiological states. Here we show that modifications in scaling slope are associated with variations in Rényi entropy, a generalization of Shannon informational entropy. From a three-dimensional object's perspective, by changing its orientation (standing for the cortical scale-free exponent), we detect different two-dimensional shadows from different perception angles (standing for Rényi entropy in different brain areas). We show how, starting from known values of Rényi entropy (easily detectable in brain fMRIs or EEG traces), it is feasible to calculate the scaling slope in a given moment and in a given brain area. Because changes in scale-free cortical dynamics modify brain activity, this issue points towards novel approaches to mind reading and description of the forces required for transcranial stimulation.

Keywords Rényi entropy · Power laws · Nervous system · Scale-free · Shadows · Central nervous system

Introduction

Counter-intuitively, a 3D object might encompass less information than a 2D one (we do not see the hidden side of the moon). When observed from a given standpoint, an opaque 3D object could be less informative than a series of

its 2D shadows (Fig. 1a). Here we ask: is there a way to increase the information content of the shadow, to make it possible the detection of more details about the object, let us say a toy? The answer is positive, if we rotate the toy clockwise (or anticlockwise) along its central axis (Fig. 1b, c). In this case, we achieve a pool of different shadows and shape contours with various angulations, able to increase our total available information. Therefore, 3D object's shadows (taken in its totality) are more informative, for a watching observer, whenever he moves. This “toy” analogy gives us the possibility to assess brain function in a novel way. For the purpose of this paper, the toy at rest stands for a given brain scale-free exponent, and its shadows for the corresponding Shannon entropy. In turn, the changes in location of the moving toy stand for changes in brain scale-free exponents, while the modifications in shadow's shape and its undulating contour stand for the corresponding Rényi entropy (Rényi 1961, 1966; Peters 2017; Peters and Ramanna 2016).

Using our example of a toy as a 3D object and its moving shadows, the aim of this paper is to show how it is feasible to extract information about such toy, by looking at its moving shadows: in neurophysiological terms, we

✉ Arturo Tozzi
tozziarturo@libero.it

James F. Peters
James.Peters3@umanitoba.ca

Mehmet Niyazi Çankaya
mehmet.cankaya@usak.edu.tr

¹ Computational Intelligence Laboratory, University of Manitoba, Winnipeg, MB R3T 5V6, Canada

² Department of Electrical and Computer Engineering, University of Manitoba, 75A Chancellor's Circle, Winnipeg, MB R3T 5V6, Canada

³ Department of Mathematics, Faculty of Arts and Sciences, Adiyaman University, 02040 Adiyaman, Turkey

⁴ Applied Sciences School, Department of International Trading, Department of Statistics, Faculty of Arts and Science, Usak University, Usak, Turkey

Fig. 1 A toy under a light source. In **a**, the 2D shadow offers more details of the 3D object than the view on the toy from above. **b**, **c** Show how toy rotations lead to shadows with different shapes, and, consequently, with different information content. Therefore, a complete 360° rotation of the object allows a complete evaluation of the shadows' shapes

want to assess the relationships among scale-free exponents and generalized informational entropies in the brain. Furthermore, we would consider the recent advancement in the neurophenomenology research that established a link between a nested operational architectonics hierarchy of brain functioning (indexed by the EEG field) and the phenomenal (mental) level of brain organization (Fingelkurts and Fingelkurts 2001; Fingelkurts et al. 2010). According to this line of research, the meanings, which subjectively (mentally) are experienced as thoughts or perceptions, can best be described objectively as created and carried by dynamic fields of neural activity of different sizes that are nested within the operational architectonics hierarchy that organize hundreds of millions of neurons and trillions of synapses (Fingelkurts et al. 2009, 2013a). We perform simulations in order to demonstrate how, starting from the Rényi entropy easily detectable in neurodata series (e.g.,

from brain fMRI or EEG), it is feasible to calculate the cortical scale-free exponent in a given moment, and, consequently, the corresponding mental state.

Shannon entropy on a toy

The brain activity, e.g., in our analogy the toy and its movements, can be described in terms of scale-free dynamics. Indeed, the brain activity observed at many spatiotemporal scales exhibits fluctuations with complex scaling behavior (Newman 2005; Fingelkurts et al. 2010, 2013a), including not just cortical electric oscillations, but also membrane potentials and neurotransmitter release (Linkenkaer-Hansen et al. 2001; Fox and Raichle 2007; Milstein et al. 2009). In particular, the frequency spectrum of the cerebral electric activity displays a scale-invariant behavior, characterized by a power spectrum, a frequency and an exponent that equals the negative slope (which we will term “scaling slope”) of the line in a log power versus log frequency plot (Pritchard 1992; Van de Ville et al. 2010). In simple terms, a scaling slope 1 = stands for a horizontal line parallel to the x axis: the steeper the line, the higher the scale-free behaviour. The (spatial) fractals and (temporal) power laws can be regarded as intrinsic properties of the brain and characterize a large class of neuronal processes (de Arcangelis and Herrmann 2010; He et al. 2010; Déli et al. 2017); moreover, pink noise distributions contain information about how large-scale physiological and pathological outcomes arise from the interactions of many small-scale processes (Jirsa et al. 2014).

Keeping this in mind, a given slope in a given moment of brain activity stands, in our example, for the toy at rest. Toy features can be described through their shadow, e.g., the Shannon entropy. The classical informational Shannon entropy is (Shannon 1948):

$$H(X) = - \sum_{x \in X} \Pr(x) \log_2[\Pr(x)].$$

The term X is a random variable with n possible outcomes, while $p_i = \Pr(x)$, for $i = 1, 2, 3, \dots, n$, is a probability distribution $\Pr$ on a finite set. Shannon entropy states that, under ergodic conditions, if we know the values of $\Pr(x)$, we may obtain the values of $H(X)$. In other words, $H(X)$ is a probability density function that defines a generic probability distribution $\Pr$ such that, if we modify $\Pr$, we obtain a different value of entropy on Shannon's curve. Hence, changing the probabilities usually occurs when we have prior information. In terms of the object depicted in Fig. 1a, the steady toy stands for the systems' microstates, undetectable by macroscopic observers, while the shadow stands for the lower-dimensional Shannon entropy.

Looking at the shadow (the Shannon entropy), we achieve otherwise undetectable information about opaque 3D objects, such as the toy features.

Now we need to find a brain functional counterpart for the moving toy, its changing shape and varying contours of shadows. Indeed, the cortical scale-free slopes are not constant, rather they display multiple possible exponents during various mental states. Despite a few skeptical studies (Tribukait and Eiken 2016), it is widely accepted that different functional states—spontaneous fluctuations, task-evoked, perceptual and motor activity (Buzsáki and Watson 2012), cognitive demands (Buiatti et al. 2007; Fetterhoff et al. 2014), ageing (Suckling et al. 2008)—account for variations in scale-free exponents across cortical regions (Tinker and Velazquez 2014; Wink et al. 2008). Accordingly, we may view brain activity as an ensemble of intertwined (mono)fractals, each with its own power slope and scaling range, each marking dynamical transitions between different response regimes (Papo 2014; see also review Fingelkurts et al. 2013a). In terms of our toy, the whole brain activity stands for all the possible 360° of rotation and their corresponding shadows.

Introducing Rényi entropy

The 360° changes in toy angulation that gives rise to different shadows can be described in terms of Shannon entropy generalizations. Among the available ones (e.g., Tsallis 1988), here we favour the one-parameter class of Rényi entropies (Rényi 1966), a flexible and underestimated information-theory based index (Cambell 1965). Rényi entropy (1961) is characterized by a changing parameter α and can be defined as:

$$H_{\alpha}(X) = \frac{1}{1-\alpha} \sum_{\substack{i=1, \\ x \in X}}^N p_i^{\alpha}(x),$$

where $0 \leq \alpha < \infty$ and $p_i^{\alpha}(x)$ is the probability of event x . The Rényi entropy approaches the Shannon entropy as α approaches 1. In other words, the Shannon entropy is a case of Rényi entropy in which $\alpha = 1$. Rényi entropy has applications in dynamical systems (Hentschel and Procaccia 1983), coding (Cambell 1965), information transfer (Jizba et al. 2012), theories in quantum mechanics, black holes' mutual information (Dong 2016), assessment of human heartbeat fluctuations and coding/noncoding DNA sequences analysis (Costa et al. 2005). Rényi entropy has also been used to quantify ecosystems dynamics and diversity: from land cover types (Carranza et al. 2007; De Luca et al. 2011), to coastal dunes environments (Drius et al. 2013), from urban mosaics (Carranza et al. 2007) to species diversity in large areas (Rocchini et al. 2013).

Why the Rényi entropy, instead of the Shannon's one?

This paragraph will be devoted to elucidate why Rényi entropy should be preferred, in order to assess brain activity. Shannon entropy depicts the discrete probabilities of an event in terms of a single curve. However, in many physical and biological cases, diversity cannot be reduced to a single index information, since all its aspects cannot be captured in a single statistic (Gorelick 2006). Rényi entropy, in turn, depicts the discrete probabilities of an event in terms of many curves. Therefore, unlike Shannon entropy, Rényi entropy makes it possible to describe the system's status not just at a specific moment, but also when its trend *varies with time* (Müller et al. 2000; Patil and Taillie 2001). Indeed, a complete characterization of landscape diversity can be achieved if, instead of a single index, a parametric family of indices is used, whose members have varying sensitivities to the presence of rare and abundant elements (Jost 2007). Unlike Shannon entropy, which is single-valued, Rényi entropy is useful in measuring a range of information levels such as those in A-yeh and Peters (2016), the dynamics of non-stationary processes (Shalymov and Fradkov 2016) and the measuring of the distribution of entropy changes over time (see, e.g., Jauregui et al. 2018). In other words, Rényi entropy offers a “continuum of possible diversity measures” (Ricotta and Avena 2003) at diverse spatiotemporal scales, becoming increasingly regulated by the commonest when α gets higher. The change in α exponent can be regarded as a scaling operation that takes place not in the real, but in the data space (Podani 1992): we are thus allowed to evaluate how changes of Rényi parameter influence the structure of information measures in the probability space of the scale-free dynamics. Diversity profiles are flat when the landscape is even, and steeply decrease as the landscape turns uneven (Jost 2010). Therefore, Rényi's formulation allows a continuum of possible diversity measures which differ in their sensitivity to the rare and abundant elements in a landscape.

Because of its build-in pre-disposition to account for self-similar systems, Rényi entropy is an effective tool to describe multifractal systems (Jizba and Arimitsu 2001). The generalized fractal dimensional and the Rényi exponent can be thought of as interchangeable. In technical terms, the Rényi parameter is connected via a Legendre transformation with the multifractal singularity spectrum. See Jizba and Arimitsu (2001) and Jizba and Korbel (2014) for technical details on mathematical proof. In particular, this means that Rényi entropy and scale-free exponent (and therefore the scaling slope) are strictly correlated. Therefore, changes in scaling slope lead to changes in the Rényi

Fig. 2 Rényi entropy' circle perimeter and its values (computed by the `Rényi.z` function in version 3.2.3 of R software) are superimposed, respectively, to the moving toy and its shadow. Left part: Cosine theorem for the evaluation of travel length Rényi entropies in a model of two-dimensional brain (modified from Van Essen 2005). Right

part: simulated Rényi values for each circle's perimeter values for $n = 60$ and a Rényi exponent = 4; θ is generated from a uniform distribution in $(0, 1)$ and multiplied by 360. The Rényi entropy values' bandwidth is around 4.6

parameter, and vice versa (Słomczynski et al. 2000): scale-free systems lead to different probability outcomes, based solely on increases or decreases of the scaling slope. In sum, the “probabilistic” virtues of Rényi entropy represent a novel physics-based approach to probe brain scale-free dynamics, with the potential to lead to new insights into brain systems at all space–time scales and all levels of complexity.

Once elucidated why Rényi entropy is more suitable than Shannon's, in the next paragraph we will try to answer to the crucial question: what for? How might such a scheme help, in the experimental assessment of brain pink noise? Can the variations (fluctuations) in Rényi entropy can be correlated with the values of a circle perimeter (standing for 360° angles of our toy example rotations)? Our moving toy might help to elucidate the issue. Indeed, toy rotations form an angle on a 360° circumference, so that every angle corresponds to a different shadow, i.e., to a different value of Rényi entropy (Fig. 1b, c). In neurophysiological terms, we are allowed to calculate the changes in brain scaling slopes, just by evaluating a full spectrum of the corresponding Rényi entropies. The procedure will be described in the next paragraph.

Working on randomly generated angles

Here we show the procedure in order to correlate our formulation of Rényi entropy with brain scale-free dynamics. We performed simulations of randomly generated angles (Çankaya et al. 2015) on a 2D circle. Every angle displays

a random degree, so that their total sum stands for a 360° full circle. The number of rays originating from the circle's center can be modelled through the cosine theorem, which states that:

$$per_i^2 = b_i^2 + c_i^2 - 2b_i c_i \cos(\theta), \quad i = 1, 2, \dots, n,$$

where b and c (which could be randomly generated in simulations) are the length of rays originating from the circle's central point. The parameter per is one of the three sides of the triangle lying on the circle perimeter (Fig. 2, left side). The randomly generated parameter θ stands for the angle between two rays, while n stands for the number of rays. The sum of per_i , namely $\sum_{i=1}^n per_i$, can be regarded as the circle's perimeter. Each set of perimeter values produces just one Rényi entropy value: therefore, we replicated the simulation one hundred times.

Note that the fluctuations in Rényi entropy values depend just on the different perimeter values. In terms of brain assessment, we evaluate the cortical scale-free activity and the scaling slopes instead of evaluating random numbers. In such a vein, two cerebral hemispheres can be unfolded and flattened into a two-dimensional reconstruction by computerized procedures (Van Essen 2005). In this model, the nervous electric activity occurs on a 2D brain surface with an approximated circular shape (Fig. 2, left side). Perimeter values allow us to detect the nervous activity's travel lengths on the brain surface. After the brain signal's travel lengths have been achieved, the corresponding Rényi Entropy values can be computed (“EntropyEstimation” package: <https://cran.r-project.org/web/packages/EntropyEstimation/EntropyEstimation.pdf>).

Figure 2, right side, displays simulated Rényi values obtained from the perimeter values computed through the cosine theorem. In toy's terms, this means that a larger rotation gives rise to higher levels of information detectable from the shadows, while, in brain's terms, the higher the variations in scaling slopes, the higher the detected Rényi entropy and vice versa.

Conclusions

We showed that a clear correlation does exist in the brain between cortical scaling slope and brain activity evaluated in terms of probability states. Our approach enables us to achieve unknown scale-free exponents from known entropy changes, allowing us to evaluate the macro-states of the nervous system based on a sole order parameter, although we lack a perfect knowledge of its micro-states. In our simulations, random numbers generated from different distributions are able to modify the values of Rényi entropy: this means that the brain scale-free spectrum can be extrapolated, just starting from the values of entropy experimentally detected in EEG or fMRI traces. Considering the connection between brain activity dynamics and the mental states (Fingelkurts and Fingelkurts 2001; Fingelkurts et al. 2009, 2010, 2013a), this observation paves the way to tackle the difficult issue of mind reading.

We hypothesize that, in order to optimize its functions, the brain could be equipped with intrinsic mechanisms of scaling features: in this framework, changes in scale-free exponents play a crucial role in information processing, leading to variations in entropy and probability of mental states. Empirical evidence suggests that cognitive tasks are modulated by the scale-free exponents of the brain fluctuation probability function, leading to a shrinking of multifractal spectrum and/or transitions from mono- to multifractal distributions (Popivanov et al. 2006; Fraiman and Chialvo 2012).

This approach paves the way for novel approaches to already existing neural network models (Gravier et al. 2016; Megam Ngouonkadi et al. 2016) and for innovative diagnostic and therapeutic strategies. For example, the widely-diffused method of pairwise entropy in neuroimaging techniques (Watanabe et al. 2014; Peters et al. 2017) might benefit of the Rényi entropy-based approach. It is indeed feasible to start from known parameters of probability distribution, in order to achieve the unknown values of multifractal slopes. In particular, a change in Rényi exponent, caused by variations in scaling slopes, might lead to different probability outcomes, which can be calculated if just their pairwise entropies are known.

We also conjecture that the scale-free-like brain electric activity—setting aside their supposed relationships with self-organized criticality (Bak et al. 1987; Tozzi 2015), nonequilibrium steady-state dynamics or second order phase transitions (Papo 2014; Perkins et al. 2014; Tozzi 2014; Tozzi and Peters 2016)—could be modulated through the superimposition of an external electric currents characterized by carefully chosen scale-free exponents. This may have a practical application. For example, our results suggest that transcranial electrical stimulation's techniques need to take into account not only the amplitude and frequency of the applied waveforms (Reato et al. 2013), but also their scaling slope. A further tenable field of application are the diseases that have been linked to disturbance of brain networks—Alzheimer's disease, depression, attention deficit hyperactivity disorder, schizophrenia, autism (Fox and Raichle 2007; Fingelkurts and Fingelkurts 2010), meaning that they could be ameliorated, or even treated, by appropriate exogenous electromagnetic fields—e.g., via selective application of electric/magnetic waves of specific scale-free exponent on target micro-areas—able to “recover” and restore the physiological brain functions (Sunderam et al. 2009; Fingelkurts and Fingelkurts 2010, 2015) and related subjective experiences (Fingelkurts et al. 2013b).

Acknowledgements The Authors would like to thank Andrew and Alexander Fingelkurts for commenting upon an earlier version of this manuscript.

References

- A-yeh E, Peters JF (2016) Rényi entropy in measuring information levels in Voronoï tessellation cells with application in digital image analysis. *Theory Appl Math Comput Sci* 6(16):77–95
- Bak P, Tang C, Wiesenfeld K (1987) Self-organized criticality: an explanation of the 1/f noise. *Phys Rev Lett* 59(4):381–384
- Buiatti M, Papo D, Baudonnière P-M, van Vreeswijk C (2007) Feedback modulates the temporal scale-free dynamics of brain electrical activity in a hypothesis testing task. *Neuroscience* 146:1400–1412. <https://doi.org/10.1016/j.neuroscience.2007.02.048>
- Buzsáki G, Watson BO (2012) Brain rhythms and neural syntax: implications for efficient coding of cognitive content and neuropsychiatric disease. *Dialogues Clin Neurosci* 4:345–367
- Cambell LL (1965) A coding theorem and Rényi entropy. *Inf Control* 8(4):423–429
- Çankaya MN, Bulut YM, Doğru FZ, Arslan O (2015) A bimodal extension of the generalized gamma distribution. *Revista Colombiana de Estadística* 38(2):371–378
- Carranza ML, Acosta A, Ricotta C (2007) Analyzing landscape diversity in time: the use of Rényi's generalized entropy function. *Ecol Indic* 7:505–510
- Costa M, Goldberger AL, Peng CK (2005) Multiscale entropy analysis of biological signals. *Phys Rev E Stat Nonlinear Softw Matter Phys* 71(2):021906
- de Arcangelis L, Herrmann HJ (2010) Learning as a phenomenon occurring in a critical state. *Proc Natl Acad Sci* 107:3977–3981

- De Luca E, Novelli C, Barbato F, Menegoni P, Iannetta M, Nascetti G (2011) Coastal dune systems and disturbance factors: monitoring and analysis in central Italy. *Environ Monit Assess* 183:437–450
- Déli E, Tozzi A, Peters JF (2017) Relationships between short and fast brain timescales. *Cogn Neurodyn* 11(6):539–552
- Dong X (2016) The gravity dual of Rényi entropy. *Nat Commun* 7:12472. <https://doi.org/10.1038/ncomms12472>
- Drius M, Malavasi M, Rosario Acosta AT, Ricotta C, Carranza ML (2013) Boundary-based analysis for the assessment of coastal dune landscape integrity over time. *Appl Geogr* 45:41–48
- Fetterhoff D, Opris I, Simpson SL, Deadwyler SA, Hampson RE, Kraft RA (2014) Multifractal analysis of information processing in hippocampal neural ensembles during working memory under $\Delta 9$ -tetrahydrocannabinol administration. *J Neurosci Methods*. <https://doi.org/10.1016/j.jneumeth.2014.07.013>
- Fingelkurts AA, Fingelkurts AA (2001) Operational architectonics of the human brain biopotential field: towards solving the mind-brain problem. *Brain Mind* 2:261–296
- Fingelkurts AA, Fingelkurts AA (2010) Alpha rhythm operational architectonics in the continuum of normal and pathological brain states: current state of research. *Int J Psychophysiol* 76:93–106
- Fingelkurts AA, Fingelkurts AA (2015) Operational architectonics methodology for EEG analysis: theory and results. *Neuromethods* 91:1–59. https://doi.org/10.1007/7657_2013_60
- Fingelkurts AA, Fingelkurts AA, Neves CFH (2009) Phenomenological architecture of mind and operational architectonics of the brain: the unified metastable continuum. *New Math Nat Comput* 5:221–244
- Fingelkurts AA, Fingelkurts AA, Neves CFH (2010) Natural world physical, brain operational, and mind phenomenal space-time. *Phys Life Rev* 7:195–249
- Fingelkurts AA, Fingelkurts AA, Neves CFH (2013a) Consciousness as a phenomenon in the operational architectonics of brain organization: criticality and self-organization considerations. *Chaos Solitons Fract* 55:13–31
- Fingelkurts AA, Fingelkurts AA, Neves CFH (2013b) The structure of brain electromagnetic field relates to subjective experience: Exogenous magnetic field stimulation study. Presented at Neuroscience Finland 2013 meeting: optogenetics and brain stimulation, Helsinki, 22 Mar 2013
- Fox MD, Raichle ME (2007) Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nat Rev Neurosci* 8:700–711
- Fraiman D, Chialvo DR (2012) What kind of noise is brain noise: anomalous scaling behavior of the resting brain activity fluctuations. *Front Physiol* 3:307. <https://doi.org/10.3389/fphys.2012.00307>
- Gorelick R (2006) Combining richness and abundance into a single diversity index using matrix analogues of Shannon's and Simpson's indices. *Ecography* 29:525–530
- Gravier A, Quek C, Duch W, Wahab A, Gravier-Rymaszewska J (2016) Neural network modelling of the influence of channelopathies on reflex visual attention. *Cogn Neurodyn* 10(1):49–72. <https://doi.org/10.1007/s11571-015-9365-x> (Epub 9 Nov 2015)
- He BJ, Zempel JM, Snyder AZ, Raichle ME (2010) The temporal structures and functional significance of scale-free brain activity. *Neuron* 66(3):353–369. <https://doi.org/10.1016/j.neuron.2010.04.020>
- Hentschel GHE, Proccacia I (1983) The infinite number of generalized dimensions in fractals and strange attractors. *Physica D* 8(3):435–444
- Jauregui M, Zunino L, Lenzi EK, Mendes RS, Ribeiro HV (2018) Characterization of time series via Rényi complexity-entropy curve. *Proc R Soc Lond Ser A Math Phys Eng Sci* 498:74–85
- Jirsa VK, Stacey WC, Quilichini PP, Ivanov AI, Bernard C (2014) On the nature of seizure dynamics. *Brain* 137(Pt 8):2210–2230. <https://doi.org/10.1093/brain/awu133>
- Jizba P, Arimitsu T (2001) The world according to Rényi: thermodynamics of fractal systems. *AIP Conf Proc* 597:341–348
- Jizba P, Korbel J (2014) Multifractal diffusion entropy analysis. *Physica A* 413:438–458
- Jizba P, Kleinert H, Shefaat M (2012) Rényi information transfer between financial time series. *Physica A* 391(10):2971
- Jost L (2007) Partitioning diversity into independent alpha and beta components. *Ecology* 88(10):2427–2439
- Jost L (2010) The relation between evenness and diversity. *Diversity* 2:207–232
- Linkenkaer-Hansen K, Nikouline VV, Palva JM, Ilmoniemi RJ (2001) Long-range temporal correlations and scaling behavior in human brain oscillations. *J Neurosci* 21(4):1370–1377
- Megam Ngouonkadi EB, Fotsin HB, Nono MK, Fotso PHL (2016) Noise effects on robust synchronization of a small pacemaker neuronal ensemble via nonlinear controller: electronic circuit design. *Cogn Neurodyn* 10(5):385–404. <https://doi.org/10.1007/s11571-016-9393-1> (Epub 11 Jun 2016)
- Milstein J, Mormann F, Fried I, Koch C (2009) Neuronal shot noise and Brownian $1/f^2$ behavior in the local field potential. *PLoS ONE* 4(2):4338. <https://doi.org/10.1371/journal.pone.0004338>
- Müller F, Hoffmann-Kroll R, Wiggering H (2000) Indicating ecosystem integrity e theoretical concepts and environmental requirements. *Ecol Model* 130:13–23
- Newman MEJ (2005) Power laws, Pareto distributions and Zipf's law. *Contemp Phys* 46:323–351
- Papo D (2014) Functional significance of complex fluctuations in brain activity: from resting state to cognitive neuroscience. *Front Syst Neurosci* 8:112. <https://doi.org/10.3389/fnsys.2014.00112>
- Patil GP, Taillie C (2001) A multiscale hierarchical Markov transition matrix model for generating and analyzing thematic raster maps. *Environ Ecol Stat* 8:5–20
- Perkins TJ, Foxall E, Glass L, Edwards R (2014) A scaling law for random walks on networks. *Nat Commun* 5:5121. <https://doi.org/10.1038/ncomms6121>
- Peters JF (2017) Foundations of computer vision. Computational geometry, visual image structures and object shape recognition. Springer, Berlin. <https://doi.org/10.1007/978-3-319-52483-2>
- Peters JF, Ramanna S (2016) Maximal nucleus clusters in Pawlak paintings. Nerves as approximating tools in visual arts. *Proc Fed Conf Comp Sci Info Syst* 8:199–202. <https://doi.org/10.15439/2016F004>
- Peters JF, Tozzi A, Ramanna S, İnan E (2017) The human brain from above: an increase in complexity from environmental stimuli to abstractions. *Cogn Neurodyn* 11(4):391–394
- Podani J (1992) Space series analysis: processes reconsidered. *Abstracta Botanica* 16:25–29
- Popivanov D, Stomonyakov V, Minchev Z, Jivkova S, Dojnov P et al (2006) Multifractality of decomposed EEG during imaginary and real visual-motor tracking. *Biol Cybern* 94:149–156
- Pritchard WS (1992) The brain in fractal time: $1/f$ -like power spectrum scaling of the human electroencephalogram. *Int J Neurosci* 66:119–129
- Reato D, Rahman A, Bikson M, Parra LC (2013) Effects of weak transcranial alternating current stimulation on brain activity—a review of known mechanisms from animal studies. *Front Human Neurosci* 7:687
- Rényi A (1961) On measures of entropy and information. In: *Proc fourth Berkeley symp math stat and probability*, vol I. University of California Press, Berkeley, pp 547–457. MR0132570
- Rényi A (1966) On the amount of information in a random variable concerning an event. *J Math Sci* 1:30–33

- Ricotta C, Avena G (2003) On the relationship between Pielou's evenness and landscape dominance within the context of Hill's diversity profiles. *Ecol Indic* 2:361–365
- Rocchini D, Delucchi L, Bacaro G, Cavallini P, Feilhauer H et al (2013) Calculating landscape diversity with information-theory based indices: a GRASS GIS solution. *Ecol Inf* 17:82–89
- Shalymov DS, Fradkov AL (2016) Dynamics of non-stationary processes that follow the maximum of the Rényi entropy principal. *Proc R Soc Lond Ser A Math Phys Eng Sci* 472(2185):20150324
- Shannon CE (1948) A mathematical theory of information. *Bell Syst Tech J* 27:379–423
- Słomczynski W, Kwapień J, Zyczkowski K (2000) Entropy computing via integration over fractal measures. *Chaos* 10(1):180–188
- Suckling J, Wink AM, Bernard FA, Barnes A, Bullmore E (2008) Endogenous multifractal brain dynamics are modulated by age, cholinergic blockade and cognitive performance. *J Neurosci Methods* 174(2):292–300. <https://doi.org/10.1016/j.jneumeth.2008.06.037>
- Sunderam S, Chernyy N, Peixoto N, Mason JP, Weinstein SL et al (2009) Seizure entrainment with polarizing low-frequency electric fields in a chronic animal epilepsy model. *J Neural Eng* 6(4):046009. <https://doi.org/10.1088/1741-2560/6/4/046009>
- Tinker J, Velazquez JL (2014) Power law scaling in synchronization of brain signals depends on cognitive load. *Front Syst Neurosci* 8:73. <https://doi.org/10.3389/fnsys.2014.00073>
- Tozzi A (2014) Evolution: networks and energy count. *Nature* 515:343. <https://doi.org/10.1038/515343c>
- Tozzi A (2015) How to turn an oscillation in a pink one. *J Theor Biol* 377:117–118. <https://doi.org/10.1016/j.jtbi.2015.04.018>
- Tozzi A, Peters JF (2016) Towards a fourth spatial dimension of brain activity. *Cogn Neurodyn* 10(3):189–199
- Tribukait A, Eiken O (2016) On the time course of short-term forgetting: a human experimental model for the sense of balance. *Cogn Neurodyn* 10(1):7–22
- Tsallis C (1988) Possible generalization of Boltzman-Gibbs statistics. *J Stat Phys* 52(2):479–487
- Van de Ville D, Britz J, Michel CM (2010) EEG microstate sequences in healthy humans at rest reveal scale-free dynamics. *PNAS* 107(42):18179–18184. <https://doi.org/10.1073/pnas.1007841107>
- Van Essen DC (2005) A population-average, landmark- and surface-based (PALS) atlas of human cerebral cortex. *Neuroimage* 28:635–666
- Watanabe T, Masuda N, Megumi F, Kanai R, Rees G (2014) Energy landscape and dynamics of brain activity during human bistable perception. *Nat Commun* 5:4765. <https://doi.org/10.1038/ncomms5765>
- Wink AM, Bullmore E, Barnes A, Bernard F, Suckling J (2008) Monofractal and multifractal dynamics of low frequency endogenous brain oscillations in functional MRI. *Human Brain Mapp* 29(7):791–801. <https://doi.org/10.1002/hbm.20593>