Published in final edited form as:

Curr Opin Immunol. 2018 August ; 53: 13–21. doi:10.1016/j.coi.2018.03.019.

Distinct and complementary roles of CD4 T cells in protective immunity to influenza virus

Andrea J. Sant^{*,1,2}, Katherine A. Richards¹, and Jennifer Nayak^{1,2,3}

¹David H. Smith Center for Vaccine Biology and Immunology, University of Rochester Medical Center

²Department of Microbiology and Immunology, University of Rochester Medical Center

³Department of Pediatrics, Division of Infectious Diseases, University of Rochester Medical Center

Abstract

CD4 T cells play a multiplicity of roles in protective immunity to influenza. Included in these functions are help for high affinity antibody production, enhancement of CD8 T cell expansion, function and memory, acceleration of the early innate response to infection and direct cytotoxicity. The influenza-specific CD4 T cell repertoire in humans established through exposures to infection and vaccination has been found to be highly variable in abundance, specificity and functionality. Deficits in particular subsets of CD4 T cells recruited into the response result in diminished antibody responses and protection from infection. Therefore, improved strategies for vaccination should include better methods to identify deficiencies in the circulating CD4 T cell repertoire, and vaccine constructs that increase the representation of CD4 T cells of the correct specificity and functionality.

Key issues and persistent challenges in understanding the contributions of CD4 T cells in protective immunity to influenza

Understanding the precise role of CD4 T cell immunity to influenza infection and vaccination is achievable but challenging. A key distinction between CD4 T cells and other cells in the adaptive immune response, such as B cells and CD8 T cells, is the multiplicity of functions that they contribute to protective influenza immunity. In order to potentiate CD4 T cell immunity for influenza, it is important to factor in three considerations: the diversity, specificity and abundance of influenza-specific CD4 T cells in human populations, the discreet functional activities that CD4 T cells convey in response to infection or vaccination, and finally, the CD4 T cell functions that are limiting factors in the efficacy of protection from influenza infection or responses to vaccination. In this review, we will discuss each of

*Corresponding author: Andrea J. Sant, andrea_sant@urmc.rochester.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

these issues, focusing whenever possible on the most recent studies and those derived from analyses of humans.

Abundance and specificity of human influenza-specific CD4 T cells

Recent studies have sought to quantify and characterize the repertoire of circulating CD4 T cells from human subjects with specificity towards influenza antigens [1–8]. In general, these studies have found that in most healthy adults, there are detectable influenza-specific CD4 T cells, but that the abundance is highly variable [1,3,5,7,9]. With the increased interest in the design of universal influenza vaccines [10,11], there has been particular interest in candidate epitopes that would elicit broadly cross-reactive CD4 T cells that are genetically conserved across viral strains [12–14]. In healthy donors, there is prominent reactivity in influenza CD4 T cells specific for the internal virion proteins M1, NP [5–7,9], and polymerase [1,4]. HA-reactive CD4 T cells have also been reported to be abundant in the memory compartment of many individuals [2–4,9,15], and these may be particularly important for provision of help for neutralizing antibody responses [16–20]. In humans, reactivity to HA is enriched for specificities in the more highly conserved HA2 domain [3], allowing candidate epitopes in this region to be utilized in universal vaccine efforts. CD4 T cells specific for M1, NP, polymerase proteins and highly conserved segments of HA are likely to be the major specificities elicited in response to infection with heterologous or novel potentially pandemic strains [2,21–27] and may contribute to attenuating the course of infection. The degree of cross reactivity in CD4 T cells elicited in response to the novel 2009 potentially pandemic strain is likely to have contributed to the mild course of disease observed in many subjects, despite a lack of cross-reactive neutralizing antibody.

The method used to assess CD4 T cell specificity and abundance is important to consider in interpretation of any studies that quantify influenza reactivity. First, it is essential to note whether or not specificity and dominance are determined directly *ex vivo* [1,3–5,7,8,15] or after expansion [6,9]. If after expansion, whether all potential epitopes or a subset of peptides or antigens were used is also important. Complex or uneven mixtures of antigens [6] or pre-selected epitopes based on predictive algorithms [1,7], or particular MHC types [1,6,8] will bias the generality of the conclusions. In addition, short term *in vitro* activation of T cells may lead to expansion of selective specificities or functions [28]. Direct *ex vivo* studies with unbiased and overlapping peptide sets, coupled with intracellular cytokine staining or cytokine EliSpots allows the most specificities to be quantified. These methods do have the caveat that quantification will be limited to CD4 T cells that produce known secreted mediators. Other cytokine-independent approaches, such as use of peptide-MHC multimers [8] overcome this deficiency, but this method only samples a fraction of the repertoire based on MHC restriction, and detects only those CD4 T cells with relatively high avidity for their ligand [29].

Functional contribution of CD4 T cells to protective immunity to influenza

CD4 T cell help for antibody responses is the most generally acknowledged and essential contribution of CD4 T cell responses to protective immunity induced by influenza vaccines and future infection [30]. The antigen-specific cognate interactions between CD4 T cells and

B cells promotes both the rapid extrafollicular and the later-evolving germinal center response that drives immunoglobulin affinity maturation and long lived B cell immunity (reviewed in [31,32]). Recent data has shown a close correlation between the elicited antibody response to vaccination and the emergence of cells in peripheral blood with markers reminiscent of follicular helper CD4 T cells (reviewed in [33–35]), arguing that CD4 T cell help for vaccine responses can be tracked soon after vaccination [36].

CD4 T cells also play key functions distinct from delivery of B cell help. They are important for protective immunity conveyed by CD8 T cells, enhancing priming, expansion and establishment of long-lived memory [37]. Intriguingly, a recent study [38] provided strong evidence that CD4 T cell help is critical for development of effector CD8 T cells that have cytotoxic potential, and express molecules important for homing and extravasation, all features required for protective immunity to influenza. CD4 T cell help for CD8 T cells during priming may involve sequential contact with discrete subsets of antigen presenting cells that culminates in localized delivery of help [39]. At the site of infection, CD4 T cells also promote the positioning of memory CD8 T cells in the infected respiratory tract [40]. Another discrete function of CD4 T cells that has been increasingly validated is cytotoxicity, which has the potential to directly eliminate infected cells (reviewed in [41–43]). In addition to enhancing adaptive immunity, a potentially critical function of influenza-specific CD4 T cell memory is the ability to accelerate recruitment of innate effectors to the lung. The role of the early innate response to infection is well documented (reviewed in [44]) and recent studies have shown influenza-specific CD4 T memory promotes this early response, and blunts virus infection [45,46]. A final activity of influenza-specific CD4 T cells, identified primarily in animal models, is regulatory and repair function. These activities are contributed by diverse cell types including conventional Treg, as well as IL-10- or amphiregulin-producing cells. Such CD4 T cells can diminish the damage associated with the profound inflammatory responses [47–50] that occur upon infection with highly pathogenic strains of influenza. In general, although influenza-specific CD4 T cells are enriched in IFN- γ production and thus can be considered “Th1-like”, cytokine production in humans is tremendously complex [27,51], relative to what is typically observed in mice, which have more limited frequencies of exposure to influenza antigens. Therefore, assigning functional activities of influenza-specific human CD4 T cells to single cytokines is quite difficult. Below, we detail recent studies on two of the important and distinct functional subsets of influenza-specific CD4 T cells: those with cytolytic function and those important for enhancing high affinity antibody responses.

Cytolytic CD4 T cells.

There has been increasing appreciation of a distinct subset of CD4 T cells with cytolytic function (reviewed in [41–43]). Influenza-specific CD4 T cells with cytotoxic potential, first identified after *in vitro* culture, have now been detected directly *ex vivo*. In many respects, cytotoxic CD4 T cells resemble CD8 T cells both in terms of key transcriptional regulators [52], efficiency of killing [53] and in primary cytotoxic mechanisms involving perforin and granzymes [41,43]. Both IL-2 and inflammatory signals, abundant in responses to viral infections, are thought to be central elements that promote the cytolytic potential of CD4 T cells [54,55]. Recent studies in human peripheral blood CD4 T cells have identified the

transcription factor HOBIT (“Homolog of Blimp-1 in T cells”) as a unique identifier of cells expressing granzymes, perforin and other markers linked to cytolytic function [56]. Until recently, CD4 T cells with cytotoxic function have been quantified by expression of stored cytolytic mediators, such as granzyme, requiring that cells be permeabilized prior to analyses. However, recently, cell surface markers, such as CRTAM (class I–restricted T cell–associated molecule) [52] and the natural killer cell marker NKG2C/E [57] have been defined that have the potential to quantify and permit isolation of intact cells with cytotoxic potential. The role of cytolytic CD4 T cells in influenza immunity has been supported by the prominence of this phenotype in the infected lung [41,54,57] and has been found as correlate of protection in human challenge studies [5].

One intriguing unresolved question is whether cytolytic CD4 T cells perform a unique function distinct from CD8 T cells. Because of their similarity in function and development, why does the host need the apparent redundant mechanism for cell mediated cytotoxicity? Two possible non-mutually exclusive possibilities can be envisioned. Cytotoxic CD4 T cells may serve as a complementary mechanism, eliminating infected cells at selected sites within the lung, perhaps controlled by a unique array of lung positioning molecules [58,59]. Recent studies have revealed the diversity of infected and antigen bearing cells detectable in the lung early after infection [60] that may be located at distinct sites in the respiratory tract. Alternatively or additionally, CD4 T cells may serve as a failsafe mechanism when CD8 T cell epitopes are lost from circulating influenza strains [61], which may be less likely to occur in CD4 T due to their broader epitope specificity.

Follicular helper cells.

Because of the importance of neutralizing antibody in protection against influenza [62–64], there has been increasing emphasis on the subset of CD4 T cells termed T follicular helper cells (Tfh) that promote the generation and maintenance of the germinal center reaction and the production of high affinity, class-switched antibody. Tfh are characterized by the expression of the chemokine receptor CXCR5, promoting localization to the B cell zone of secondary lymphoid organs. Here, they also express high levels of ICOS and PD-1 ([65,66] and reviewed in [35,67,68]). The transcriptional repressor Bcl-6 directs lineage commitment [69,70]. Cytokines, including IL-6 and IL-21 [71], and T cell receptor (TcR) signal strength [72–75] have all been shown to influence Tfh differentiation. Interestingly, beyond TcR signal strength and cytokine milieu, particular dendritic cell subsets may also be critical in priming of Tfh. Lung CD11b+ migratory dendritic cells (cDC2) have been implicated in priming following intranasal immunization [76], and late appearing APCs were identified in the Tfh response following influenza virus infection [77].

The relationship between circulating CXCR5+ cells and the Tfh within secondary lymphoid organs was uncertain until relatively recently. In circulation, CXCR5+ cells have lower expression of Bcl-6 as well ICOS and PD-1 [78–80]. However, despite these phenotypic differences, circulating CXCR5+ CD4 T cells demonstrate a superior capacity to help B cells upon activation [78,80,81]. Circulating CXCR5+ cells can be further distinguished by expression of CCR6, CXCR3, ICOS and PD-1, with helper activity concentrated within the CXCR3- cells that are PD1+ and/or ICOS+[36,78,81–83].

There has been much progress in understanding the human Tfh response following influenza vaccination. Increases in Tfh expressing activation markers (ICOS⁺ or ICOS⁺ PD1⁺) correlates with the magnitude [36,82–84] and avidity [85] of influenza-specific antibody. Circulating Tfh cells identified after vaccination appear to be clonally related those within the germinal center and transition to a more quiescent phenotype at memory [79,86]. While circulating influenza-specific Tfh have reactivity to multiple influenza proteins [85], they are relatively enriched within cells specific for the HA compared to the NP protein [15]. Although the mechanisms underlying this disparity are unknown, studies to characterize whether there are differences in Tfh frequency between epitopes within the more conserved and the more divergent portions of HA could help to clarify whether repeated boosting by vaccination or infection influences Tfh abundance. The role of antigen specificity in CD4 T cell help for antibody response is a critical parameter in predicting and boosting the capacity of the human host to produce protective antibody. Recent studies in both animal models of infection [18,20,87] and human vaccination studies [16,17] suggest that the most effective help for antibody responses occurs when the antigen specificity of the CD4 T cells matches that of the B cell. This obligate linkage likely reflects that nature of the antigen taken up by the immunoglobulin receptor, and the resulting MHC class II:peptide complexes that are displayed at the cell surface of the antigen specific B cells that recruit CD4 T cell help in secondary lymphoid tissue. This constraint likely limits the CD4 T cell help available for novel avian strains of influenza.

The challenge: Identification of the CD4 T cells subset(s) that are a limiting factor for human protective immunity to influenza virus.

It is clear that the presence of neutralizing antibody to influenza is the best correlate of protection. However, the ability of influenza virus to mutate and to deviate from serum-mediated protection and the resulting high burden of seasonal influenza infection indicates that current strategies to induce sterilizing immunity are inadequate. Thus, there is a need to enhance protective immunity by more effectively engaging the cellular arm of the immune response. Despite this need, a significant challenge is definition of the CD4 T cells whose representation is deficient and needs to be boosted in humans by vaccination. In studies of human immunity to influenza, one of the most challenging issues is identification of cells that will be drawn into and participate in the response to infection or vaccination. For example, the vast majority of studies that have identified correlates of influenza-specific CD4 T cells with the elicited antibody responses have done so only through examination of cells that have increased their representation after vaccination [16,17,36,82,84,88,89]. Very few correlates have been identified within the circulating CD4 T cell repertoire prior to vaccination or infection. One human influenza challenge study did correlate levels of pre-existing cytotoxic CD4 T cells with protection, but this subset was identified after in vitro expansion [5]. More sensitive detection of additional markers such as chemokine receptors expressed by influenza-specific CD4 T cells, coupled with early sampling times (e.g. 1–3 days post vaccination or infection) and human challenge studies [90–93] would likely provide insight into the circulating memory CD4 T cell subsets that are recruited into the response and whose presence in peripheral blood of humans either positively or negatively correlates with protective immunity to influenza. As knowledge is gained in these issues, we

will have more of the needed insight into the most critical deficiencies in the influenza specific CD4 T cell repertoire and with this, the needed framework for improved vaccine strategies.

Many of the *in vivo* studies performed to date that have implicated particular effector functions of CD4 T cells have utilized animal models where the contribution of CD4 T cells to immunity has been demonstrated either using adoptive transfer strategies to introduce a CD4 T cell with dominant function into a naïve host or where functionality of the CD4 T cells can be eliminated or promoted by genetic means [94–96]. Extrapolating these types studies to humans is difficult because of complex pre-existing immunity that accumulates over a lifetime of exposure. To be able to exploit the contribution of CD4 T cells toward protective immunity or more robust responses to vaccination, it is essential to quantify the functional activities of CD4 T cells that are limiting in protective immunity to influenza (See Figure 1). Do most people need more cytotoxic CD4 T cells or is more tissue resident memory needed? Is CD4 T cell help for rapid antibody responses to infection limiting for most people? Do only a small number of influenza-specific cells in humans have the capacity to be recruited to the lung early after infection and quickly mobilize the innate response? The unique and opposing transcription factor expression in cytolytic CD4 T cells and Tfh cells indicates that these subsets are reciprocally regulated [43,97] and vaccine strategies may need to focus the elicited response on one subset or the other. Understanding these issues will help direct the priorities in design and composition of more effective vaccines that most effectively establish protective immunity.

From the studies of human responses to vaccination and animal models of infection, it is clear that CD4 T cells of the correct functional subset [82–85] and epitope specificity [16,18,20,87,98] can be a limiting factor in eliciting neutralizing antibody. Evidence from animal models has supported the possibility that more abundant CD4 T cells of the correct specificity can enhance antibody production during the course of infection [18]. Such early antibody responses, even of non-neutralizing HA or NA specificities, could blunt the course of infection, through cellular and Fc-receptor-based mechanisms [98–102]. Vaccine strategies that populate the host with CD4 T cells that are poised to become Tfh cells of the desired protein specificity, matching the B cell specificity, may thus lead to antibody responses that can be lead to either sterilizing immunity and/or attenuated disease. Greater peripheral Tfh following adjuvanted vaccination [103], use of high dose influenza vaccination in the elderly [104], and strategies that target protein vaccines to cell surface proteins such as Clec9a [105], or particular dendritic cell subsets [106] suggests that it may be possible to use alternative vaccination strategies to optimize Tfh priming, which may foster stable Tfh memory that can promote future protective antibody responses.

The possibility of enhancing protection from influenza infection through amplification of T cell responses has been supported by recent vaccine and challenge studies in humans [90,91]. Use of virus vectors to expand T cells specific for M1 and NP, without elicitation of antibodies was associated with a lower infection frequency, virus shedding and diminished symptomology when a challenge virus was administered [90]. Consistent findings were observed when multi-epitope vaccines were similarly tested [91]. It is likely that these types of experimental challenge models [92,93], coupled with strategies to selectively promote

expansion of CD4 T cells with the desired functionality and specificity could provide critical new insight into the most limiting protective functions of CD4 T cells.

The roles of specific viral antigens as targets of vaccination, different innate activators, and antigen persistence are increasingly better understood [107–111]. C-lectin receptors are becoming increasingly used to both target antigens to APC and activate intracellular signaling that collectively leads to initiation of particular effector functions (reviewed in [112–114]). For example, coupling antigens to antibodies specific for Clec9a can promote establishment of Tfh cells [105], whereas an antigen conjugate directing uptake Dectin 1 can enhance recruitment of Th17 [115]. Also, because virus infection, known to induce long-term protective memory, has been shown to lead to long-lived reservoirs of viral antigen [116–120], vaccine strategies that can establish these depots may be particularly efficacious. Finally, the increasing recognition of the importance of local immunity in the respiratory tract suggests that efforts to promote resident CD4 T cells in the lung, through the use of intranasal attenuated viruses or synthetic vaccines, may promote establishment of memory cells with enhanced ability to home back to the lung after infection. Priming of CD4 T cells with lung-derived dendritic cells, either in isolation [121] or through infection or administration of attenuated vaccines [30,108] is known to enhance homing potential of CD4 T cells to the lung, support this view.

Collectively, recent advances have provided much of the insight needed to initiate cross protective immunity at the systemic level as well as the site of infection. The control of CD4 T cell fate decisions that regulate generation and expansion of discrete functional subsets of CD4 T cells are complex and multilayered. The regulatory events likely involve distinct sets of APC, T cell receptor-derived signals, cytokines and innate activators. The major challenge will be to design the appropriate vaccination and challenge studies to reveal and ultimately correct the most critical deficits in CD4 T cell functions in human populations.

Acknowledgments

This project has been funded in whole or in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, under CEIRS Contract No. HHSN272201400006C from the National Institutes of Health to Andrea Sant and National Institutes of Health 1K08AI106954 and Doris Duke Charitable Foundation Clinical Scientist Development Award 2015098 awarded to Jennifer Nayak.

Literature Cited

1. Assarsson E, Bui HH, Sidney J, Zhang Q, Glenn J, Oseroff C, Mbawuike IN, Alexander J, Newman MJ, Grey H, et al.: Immunomic analysis of the repertoire of T-cell specificities for influenza A virus in humans. *J Virol* 2008, 82:12241–12251. [PubMed: 18842709]
2. Babon JA, Cruz J, Orphin L, Pazoles P, Co MD, Ennis FA, Terajima M: Genome-wide screening of human T-cell epitopes in influenza A virus reveals a broad spectrum of CD4(+) T-cell responses to internal proteins, hemagglutinins, and neuraminidases. *Hum Immunol* 2009, 70:711–721. [PubMed: 19524006]
3. Richards KA, Nayak J, Chaves FA, DiPiazza A, Knowlton ZA, Alam S, Treanor JJ, Sant AJ: Seasonal Influenza Can Poise Hosts for CD4 T-Cell Immunity to H7N9 Avian Influenza. *J Infect Dis* 2015, 212:86–94. [PubMed: 25492919]
4. Richards KA, Topham D, Chaves FA, Sant AJ: Cutting edge: CD4 T cells generated from encounter with seasonal influenza viruses and vaccines have broad protein specificity and can directly

recognize naturally generated epitopes derived from the live pandemic H1N1 virus. *J Immunol* 2010, 185:4998–5002. [PubMed: 20889549]

5. Wilkinson TM, Li CK, Chui CS, Huang AK, Perkins M, Liebner JC, Lambkin-Williams R, Gilbert A, Oxford J, Nicholas B, et al.: Preexisting influenza-specific CD4+ T cells correlate with disease protection against influenza challenge in humans. *Nat Med* 2012, 18:274–280. [PubMed: 22286307]
6. Chen L, Zanker D, Xiao K, Wu C, Zou Q, Chen W: Immunodominant CD4+ T-cell responses to influenza A virus in healthy individuals focus on matrix 1 and nucleoprotein. *J Virol* 2014, 88:11760–11773. [PubMed: 25078703]
7. Savic M, Dembinski JL, Kim Y, Tunheim G, Cox RJ, Oftung F, Peters B, Mjaaland S: Epitope specific T-cell responses against influenza A in a healthy population. *Immunology* 2016, 147:165–177. [PubMed: 26489873]
8. Uchtenhagen H, Rims C, Blahnik G, Chow IT, Kwok WW, Buckner JH, James EA: Efficient ex vivo analysis of CD4+ T-cell responses using combinatorial HLA class II tetramer staining. *Nat Commun* 2016, 7:12614. [PubMed: 27571776]
9. Hayward AC, Wang L, Goonetilleke N, Fragaszy EB, Bermingham A, Copas A, Dukes O, Millett ER, Nazareth I, Nguyen-Van-Tam JS, et al.: Natural T Cell mediated Protection against Seasonal and Pandemic Influenza. Results of the Flu Watch Cohort Study. *Am J Respir Crit Care Med* 2015, 191:1422–1431. [PubMed: 25844934]
10. Paules CI, Marston HD, Eisinger RW, Baltimore D, Fauci AS: The Pathway to a Universal Influenza Vaccine. *Immunity* 2017, 47:599–603. [PubMed: 29045889]
11. Krammer F, Palese P: Advances in the development of influenza virus vaccines. *Nat Rev Drug Discov* 2015, 14:167–182. [PubMed: 25722244]
12. Tutykhina I, Esmagambetov I, Bagaev A, Pichugin A, Lysenko A, Shcherbinin D, Sedova E, Logunov D, Shmarov M, Ataullakhanov R, et al.: Vaccination potential of B and T epitope-enriched NP and M2 against Influenza A viruses from different clades and hosts. *PLoS One* 2018, 13:e0191574. [PubMed: 29377916]
13. Lee YT, Ko EJ, Lee Y, Kim KH, Kim MC, Lee YN, Kang SM: Intranasal vaccination with M2e5x virus-like particles induces humoral and cellular immune responses conferring cross-protection against heterosubtypic influenza viruses. *PLoS One* 2018, 13:e0190868. [PubMed: 29324805]
14. Eliasson DG, Omokanye A, Schon K, Wenzel UA, Bernasconi V, Bemark M, Kolpe A, El Bakkouri K, Ysenbaert T, Deng L, et al.: M2e-tetramer-specific memory CD4 T cells are broadly protective against influenza infection. *Mucosal Immunol* 2018, 11:273–289. [PubMed: 28295019]
15. Leddon SA, Richards KA, Treanor JJ, Sant AJ: Abundance and specificity of influenza reactive circulating memory follicular helper and nonfollicular helper CD4 T cells in healthy adults. *Immunology* 2015, 146:157–162. [PubMed: 26094691]
16. Nayak JL, Fitzgerald TF, Richards KA, Yang H, Treanor JJ, Sant AJ: CD4+ T-cell expansion predicts neutralizing antibody responses to monovalent, inactivated 2009 pandemic influenza A(H1N1) virus subtype H1N1 vaccine. *J Infect Dis* 2013, 207:297–305.
17. Nayak JL, Richards KA, Yang H, Treanor JJ, Sant AJ: Effect of influenza A(H5N1) vaccine prepandemic priming on CD4+ T-cell responses. *J Infect Dis* 2015, 211:1408–1417. [PubMed: 25378637]
- *. With [18], these experiments provide evidence that selectively amplifying the CD4 T cell memory repertoire that is specific for HA-derived epitopes enhances future HA-specific antibody responses to vaccination or infection.
18. Alam S, Knowlden ZA, Sangster MY, Sant AJ: CD4 T cell help is limiting and selective during the primary B cell response to influenza virus infection. *J Virol* 2014, 88:314–324. [PubMed: 24155379]
- *.
19. Sant AJ, McMichael A: Revealing the role of CD4(+) T cells in viral immunity. *J Exp Med* 2012, 209:1391–1395. [PubMed: 22851641]
20. Alam S, Chan C, Qiu X, Shannon I, White CL, Sant AJ, Nayak JL: Selective prepriming of HA-specific CD4 T cells restores immunological reactivity to HA on heterosubtypic influenza infection. *PLoS One* 2017, 12:e0176407. [PubMed: 28493882]

21. Richards KA, Chaves FA, Sant AJ: Infection of HLA-DR1 transgenic mice with a human isolate of influenza A virus (H1N1) primes a diverse CD4 T-cell repertoire that includes CD4 T cells with heterosubtypic cross-reactivity to avian (H5N1) influenza virus. *J Virol* 2009, 83:6566–6577. [PubMed: 19386707]
22. Duvvuri VR, Duvvuri B, Jamnik V, Gubbay JB, Wu J, Wu GE: T cell memory to evolutionarily conserved and shared hemagglutinin epitopes of H1N1 viruses: a pilot scale study. *BMC Infect Dis* 2013, 13:204. [PubMed: 23641949]
23. Duvvuri VR, Moghadas SM, Guo H, Duvvuri B, Heffernan JM, Fisman DN, Wu GE, Wu J: Highly conserved cross-reactive CD4+ T-cell HA-epitopes of seasonal and the 2009 pandemic influenza viruses. *Influenza Other Respir Viruses* 2010, 4:249–258.
24. Alam S, Sant AJ: Infection with seasonal influenza virus elicits CD4 T cells specific for genetically conserved epitopes that can be rapidly mobilized for protective immunity to pandemic H1N1 influenza virus. *J Virol* 2011, 85:13310–13321. [PubMed: 21976658]
25. Weinfurter JT, Brunner K, Capuano SV, 3rd, Li C, Broman KW, Kawaoka Y, Friedrich TC: Cross-reactive T cells are involved in rapid clearance of 2009 pandemic H1N1 influenza virus in nonhuman primates. *PLoS Pathog* 2011, 7:e1002381. [PubMed: 22102819]
26. Bethell D, Saunders D, Jongkaewwattana A, Kramyu J, Thitithayanont A, Wiboonut S, Yongvanitchit K, Limsalakpetch A, Kum-Arb U, Uthaimongkol N, et al.: Evaluation of in vitro cross-reactivity to avian H5N1 and pandemic H1N1 2009 influenza following prime boost regimens of seasonal influenza vaccination in healthy human subjects: a randomised trial. *PLoS One* 2013, 8:e59674. [PubMed: 23555741]
27. Roti M, Yang J, Berger D, Huston L, James EA, Kwok WW: Healthy human subjects have CD4+ T cells directed against H5N1 influenza virus. *J Immunol* 2008, 180:1758–1768. [PubMed: 18209073]
28. Koning D, Costa AI, Hasrat R, Grady BP, Spijkers S, Nanlohy N, Kesmir C, van Baarle D: In vitro expansion of antigen-specific CD8(+) T cells distorts the T-cell repertoire. *J Immunol Methods* 2014, 405:199–203. [PubMed: 24512815]
29. Martinez RJ, Evavold BD: Lower Affinity T Cells are Critical Components and Active Participants of the Immune Response. *Front Immunol* 2015, 6:468. [PubMed: 26441973]
30. Devarajan P, Bautista B, Vong AM, McKinstry KK, Strutt TM, Swain SL: New Insights into the Generation of CD4 Memory May Shape Future Vaccine Strategies for Influenza. *Front Immunol* 2016, 7:136. [PubMed: 27148257]
31. Victora GD, Wilson PC: Germinal center selection and the antibody response to influenza. *Cell* 2015, 163:545–548. [PubMed: 26496601]
32. Dufaud CR, McHeyzer-Williams LJ, McHeyzer-Williams MG: Deconstructing the germinal center, one cell at a time. *Curr Opin Immunol* 2017, 45:112–118. [PubMed: 28319730]
33. Linterman MA, Hill DL: Can follicular helper T cells be targeted to improve vaccine efficacy? *F1000Res* 2016, 5.
34. Ueno H: Human Circulating T Follicular Helper Cell Subsets in Health and Disease. *J Clin Immunol* 2016, 36 Suppl 1:34–39. [PubMed: 26984851]
35. Hale JS, Ahmed R: Memory T follicular helper CD4 T cells. *Front Immunol* 2015, 6:16. [PubMed: 25699040]
36. Herati RS, Reuter MA, Dolfi DV, Mansfield KD, Aung H, Badwan OZ, Kurupati RK, Kannan S, Ertl H, Schmader KE, et al.: Circulating CXCR5+PD-1+ response predicts influenza vaccine antibody responses in young adults but not elderly adults. *J Immunol* 2014, 193:3528–3537. [PubMed: 25172499]
37. Bevan MJ: Helping the CD8(+) T-cell response. *Nat Rev Immunol* 2004, 4:595–602. [PubMed: 15286726]
38. Ahrends T, Spanjaard A, Pilzecker B, Babala N, Bovens A, Xiao Y, Jacobs H, Borst J: CD4(+) T Cell Help Confers a Cytotoxic T Cell Effector Program Including Coinhibitory Receptor Downregulation and Increased Tissue Invasiveness. *Immunity* 2017, 47:848–861 e845. [PubMed: 29126798]
- *. Using a DNA “cancer vaccination” model that included a CD8 epitope with or without incorporation of a CD4 “helper” epitope, this work describes the gene expression profiles, ingenuity analyses

and in vivo functional studies indicating that many of the key functions needed by cytolytic CD8 T cells are deficient when a source of CD4 help is not included in vaccines.

39. Eickhoff S, Brewitz A, Gerner MY, Klauschen F, Komander K, Hemmi H, Garbi N, Kaisho T, Germain RN, Kastenmuller W: Robust Anti-viral Immunity Requires Multiple Distinct T Cell-Dendritic Cell Interactions. *Cell* 2015, 162:1322–1337. [PubMed: 26296422]
40. Laidlaw BJ, Zhang N, Marshall HD, Staron MM, Guan T, Hu Y, Cauley LS, Craft J, Kaech SM: CD4+ T cell help guides formation of CD103+ lung-resident memory CD8+ T cells during influenza viral infection. *Immunity* 2014, 41:633–645. [PubMed: 25308332]
41. Brown DM, Lampe AT, Workman AM: The Differentiation and Protective Function of Cytolytic CD4 T Cells in Influenza Infection. *Front Immunol* 2016, 7:93. [PubMed: 27014272]
42. Juno JA, van Bockel D, Kent SJ, Kelleher AD, Zaunders JJ, Munier CM: Cytotoxic CD4 T Cells-Friend or Foe during Viral Infection? *Front Immunol* 2017, 8:19. [PubMed: 28167943]
43. Takeuchi A, Saito T: CD4 CTL, a Cytotoxic Subset of CD4(+) T Cells, Their Differentiation and Function. *Front Immunol* 2017, 8:194. [PubMed: 28280496]
44. Iwasaki A, Pillai PS: Innate immunity to influenza virus infection. *Nat Rev Immunol* 2014, 14:315–328. [PubMed: 24762827]
45. Teijaro JR, Verhoeven D, Page CA, Turner D, Farber DL: Memory CD4 T cells direct protective responses to influenza virus in the lungs through helper-independent mechanisms. *J Virol* 2010, 84:9217–9226. [PubMed: 20592069]
46. Strutt TM, McKinstry KK, Dibble JP, Winchell C, Kuang Y, Curtis JD, Huston G, Dutton RW, Swain SL: Memory CD4+ T cells induce innate responses independently of pathogen. *Nat Med* 2010, 16:558–564, 551p following 564. [PubMed: 20436484]
47. Arpaia N, Green JA, Moltedo B, Arvey A, Hemmers S, Yuan S, Treuting PM, Rudensky AY: A Distinct Function of Regulatory T Cells in Tissue Protection. *Cell* 2015, 162:1078–1089. [PubMed: 26317471]
48. Brincks EL, Roberts AD, Cookenham T, Sell S, Kohlmeier JE, Blackman MA, Woodland DL: Antigen-specific memory regulatory CD4+Foxp3+ T cells control memory responses to influenza virus infection. *J Immunol* 2013, 190:3438–3446. [PubMed: 23467933]
49. Egarnes B, Gosselin J: Contribution of Regulatory T Cells in Nucleotide-Binding Oligomerization Domain 2 Response to Influenza Virus Infection. *Front Immunol* 2018, 9:132. [PubMed: 29445379]
50. Zaiss DMW, Gause WC, Osborne LC, Artis D: Emerging functions of amphiregulin in orchestrating immunity, inflammation, and tissue repair. *Immunity* 2015, 42:216–226. [PubMed: 25692699]
51. Mosmann TR, Naim I, Rebhahn J, Datta S, Cavenaugh JS, Weaver JM, Sharma G: SWIFT-scalable clustering for automated identification of rare cell populations in large, high-dimensional flow cytometry datasets, part 2: biological evaluation. *Cytometry A* 2014, 85:422–433. [PubMed: 24532172]
52. Takeuchi A, Badr Mel S, Miyauchi K, Ishihara C, Onishi R, Guo Z, Sasaki Y, Ike H, Takumi A, Tsuji NM, et al.: CRTAM determines the CD4+ cytotoxic T lymphocyte lineage. *J Exp Med* 2016, 213:123–138. [PubMed: 26694968]
- *. Using the cell surface protein CRTAM, previously identified on activated CD8 T cells, this study shows the transcriptional relatedness between CD8 cytotoxic cells and cytotoxic CD4 cells and that ectopic expression of CRTAM on CD4 T cells promotes their cytotoxic potential.
53. Hildemann SK, Eberlein J, Davenport B, Nguyen TT, Victorino F, Homann D: High efficiency of antiviral CD4(+) killer T cells. *PLoS One* 2013, 8:e60420. [PubMed: 23565245]
54. Workman AM, Jacobs AK, Vogel AJ, Condon S, Brown DM: Inflammation enhances IL-2 driven differentiation of cytolytic CD4 T cells. *PLoS One* 2014, 9:e89010. [PubMed: 24586481]
55. Hua L, Yao S, Pham D, Jiang L, Wright J, Sawant D, Dent AL, Braciale TJ, Kaplan MH, Sun J: Cytokine-dependent induction of CD4+ T cells with cytotoxic potential during influenza virus infection. *J Virol* 2013, 87:11884–11893. [PubMed: 23986597]
56. Oja AE, Vieira Braga FA, Remmerswaal EB, Kragten NA, Hertoghs KM, Zuo J, Moss PA, van Lier RA, van Gisbergen KP, Hombrink P: The Transcription Factor Hobit Identifies Human Cytotoxic CD4(+) T Cells. *Front Immunol* 2017, 8:325. [PubMed: 28392788]

57. Marshall NB, Vong AM, Devarajan P, Brauner MD, Kuang Y, Nayar R, Schutten EA, Castonguay CH, Berg LJ, Nutt SL, et al.: NKG2C/E Marks the Unique Cytotoxic CD4 T Cell Subset, ThCTL, Generated by Influenza Infection. *J Immunol* 2017, 198:1142–1155. [PubMed: 28031335]
- *. Identifies NKG2C/E as a marker of cytotoxic CD4 T cells in the lungs of influenza infected mice, allowing preparative sorting and detailed characterization of their cytotoxic mechanisms and transcriptional profiles.
58. Richter M, Ray SJ, Chapman TJ, Austin SJ, Rebhahn J, Mosmann TR, Gardner H, Kotelianski V, deFougerolles AR, Topham DJ: Collagen distribution and expression of collagen-binding alpha1beta1 (VLA-1) and alpha2beta1 (VLA-2) integrins on CD4 and CD8 T cells during influenza infection. *J Immunol* 2007, 178:4506–4516. [PubMed: 17372009]
59. Takamura S: Persistence in Temporary Lung Niches: A Survival Strategy of Lung-Resident Memory CD8(+) T Cells. *Viral Immunol* 2017, 30:438–450. [PubMed: 28418771]
60. DiPiazza A, Nogales A, Poulton N, Wilson PC, Martinez-Sobrido L, Sant AJ: Pandemic 2009 H1N1 Influenza Venus reporter virus reveals broad diversity of MHC class II-positive antigen-bearing cells following infection in vivo. *Sci Rep* 2017, 7:10857.
61. Voeten JT, Bestebroer TM, Nieuwkoop NJ, Fouchier RA, Osterhaus AD, Rimmelzwaan GF: Antigenic drift in the influenza A virus (H3N2) nucleoprotein and escape from recognition by cytotoxic T lymphocytes. *J Virol* 2000, 74:6800–6807. [PubMed: 10888619]
62. Treanor J, Wright PF: Immune correlates of protection against influenza in the human challenge model. *Dev Biol (Basel)* 2003, 115:97–104. [PubMed: 15088780]
63. de Jong JC, Palache AM, Beyer WE, Rimmelzwaan GF, Boon AC, Osterhaus AD: Haemagglutination-inhibiting antibody to influenza virus. *Dev Biol (Basel)* 2003, 115:63–73. [PubMed: 15088777]
64. Rimmelzwaan GF, McElhaney JE: Correlates of protection: novel generations of influenza vaccines. *Vaccine* 2008, 26 Suppl 4:D41–44. [PubMed: 19230158]
65. Schaerli P, Willmann K, Lang AB, Lipp M, Loetscher P, Moser B: CXC chemokine receptor 5 expression defines follicular homing T cells with B cell helper function. *J Exp Med* 2000, 192:1553–1562. [PubMed: 11104798]
66. Breitfeld D, Ohl L, Kremmer E, Ellwart J, Sallusto F, Lipp M, Forster R: Follicular B helper T cells express CXC chemokine receptor 5, localize to B cell follicles, and support immunoglobulin production. *J Exp Med* 2000, 192:1545–1552. [PubMed: 11104797]
67. Vinuesa CG, Linterman MA, Yu D, MacLennan IC: Follicular Helper T Cells. *Annu Rev Immunol* 2016, 34:335–368. [PubMed: 26907215]
68. Crotty S: A brief history of T cell help to B cells. *Nat Rev Immunol* 2015, 15:185–189. [PubMed: 25677493]
69. Yu D, Rao S, Tsai LM, Lee SK, He Y, Sutcliffe EL, Srivastava M, Linterman M, Zheng L, Simpson N, et al.: The transcriptional repressor Bcl-6 directs T follicular helper cell lineage commitment. *Immunity* 2009, 31:457–468. [PubMed: 19631565]
70. Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, Wang YH, Dong C: Bcl6 mediates the development of T follicular helper cells. *Science* 2009, 325:1001–1005. [PubMed: 19628815]
71. Nurieva RI, Chung Y, Hwang D, Yang XO, Kang HS, Ma L, Wang YH, Watowich SS, Jetten AM, Tian Q, et al.: Generation of T follicular helper cells is mediated by interleukin-21 but independent of T helper 1, 2, or 17 cell lineages. *Immunity* 2008, 29:138–149. [PubMed: 18599325]
72. Fazilleau N, McHeyzer-Williams LJ, Rosen H, McHeyzer-Williams MG: The function of follicular helper T cells is regulated by the strength of T cell antigen receptor binding. *Nat Immunol* 2009, 10:375–384. [PubMed: 19252493]
73. Tubo NJ, Pagan AJ, Taylor JJ, Nelson RW, Linehan JL, Ertelt JM, Huseby ES, Way SS, Jenkins MK: Single naive CD4+ T cells from a diverse repertoire produce different effector cell types during infection. *Cell* 2013, 153:785–796. [PubMed: 23663778]
74. Keck S, Schmalzer M, Ganter S, Wyss L, Oberle S, Huseby ES, Zehn D, King CG: Antigen affinity and antigen dose exert distinct influences on CD4 T-cell differentiation. *Proc Natl Acad Sci U S A* 2014, 111:14852–14857. [PubMed: 25267612]

75. Krishnamoorthy V, Kannanganat S, Maienschein-Cline M, Cook SL, Chen J, Bahroos N, Sievert E, Corse E, Chong A, Sciammas R: The IRF4 Gene Regulatory Module Functions as a Read-Write Integrator to Dynamically Coordinate T Helper Cell Fate. *Immunity* 2017, 47:481–497 e487. [PubMed: 28930660]
- *. Provides evidence that the transcription factor Irf4 serves as a rheostat of TcR signal strength during CD4 T cell activation with increased signal strength promoting “effector” CD4 T cells, at the expense of Tfh. Like [79] these studies emphasize the opposing transcriptional programs and multiplicity of factors involved in fate decisions for cytotoxic and follicular helper cells.
76. Krishnaswamy JK, Gowthaman U, Zhang B, Mattsson J, Szeponik L, Liu D, Wu R, White T, Calabro S, Xu L, et al.: Migratory CD11b(+) conventional dendritic cells induce T follicular helper cell-dependent antibody responses. *Sci Immunol* 2017, 2.
- *. This research explored the role of different subsets of DCs and linked antigen presentation by CD11b+ migratory type 2 cDC with Tfh induction. This suggests that methods of antigenic targeting could be utilized to more selectively prime a Tfh-biased CD4 T cell response.
77. Yoo JK, Fish EN, Braciale TJ: LAPCs promote follicular helper T cell differentiation of Ag-primed CD4+ T cells during respiratory virus infection. *J Exp Med* 2012, 209:1853–1867. [PubMed: 22987801]
78. Locci M, Havenar-Daughton C, Landais E, Wu J, Kroenke MA, Arlehamn CL, Su LF, Cubas R, Davis MM, Sette A, et al.: Human circulating PD-1+CXCR3-CXCR5+ memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody responses. *Immunity* 2013, 39:758–769. [PubMed: 24035365]
79. Heit A, Schmitz F, Gerdt S, Flach B, Moore MS, Perkins JA, Robins HS, Aderem A, Spearman P, Tomaras GD, et al.: Vaccination establishes clonal relatives of germinal center T cells in the blood of humans. *J Exp Med* 2017, 214:2139–2152. [PubMed: 28637884]
- *. This study specifically linked clones of germinal center and activated circulating Tfh. In addition, it suggested that following antigenic challenge, ICOS+ PD1+ Tfh will transition to a less activated, PD1 single positive phenotype and persist within the circulating memory pool.
80. Chevalier N, Jarrossay D, Ho E, Avery DT, Ma CS, Yu D, Sallusto F, Tangye SG, Mackay CR: CXCR5 expressing human central memory CD4 T cells and their relevance for humoral immune responses. *J Immunol* 2011, 186:5556–5568. [PubMed: 21471443]
81. Morita R, Schmitt N, Bentebibel SE, Ranganathan R, Bourdery L, Zurawski G, Foucat E, Dullaers M, Oh S, Sabzghabaei N, et al.: Human blood CXCR5(+)CD4(+) T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion. *Immunity* 2011, 34:108–121. [PubMed: 21215658]
82. Bentebibel SE, Lopez S, Obermoser G, Schmitt N, Mueller C, Harrod C, Flano E, Mejias A, Albrecht RA, Blankenship D, et al.: Induction of ICOS+CXCR3+CXCR5+ TH cells correlates with antibody responses to influenza vaccination. *Sci Transl Med* 2013, 5:176ra132.
83. He J, Tsai LM, Leong YA, Hu X, Ma CS, Chevalier N, Sun X, Vandenberg K, Rockman S, Ding Y, et al.: Circulating precursor CCR7(lo)PD-1(hi) CXCR5(+) CD4(+) T cells indicate Tfh cell activity and promote antibody responses upon antigen reexposure. *Immunity* 2013, 39:770–781. [PubMed: 24138884]
- *. Using both mouse and human models, this study probed the role of activated circulating Tfh cells in B cell responses and, using an adoptive transfer strategy, examined the ability of these cells to be recalled upon antigenic rechallenge.
84. Spensieri F, Siena E, Borgogni E, Zedda L, Cantisani R, Chiappini N, Schiavetti F, Rosa D, Castellino F, Montomoli E, et al.: Early Rise of Blood T Follicular Helper Cell Subsets and Baseline Immunity as Predictors of Persisting Late Functional Antibody Responses to Vaccination in Humans. *PLoS One* 2016, 11:e0157066. [PubMed: 27336786]
85. Bentebibel SE, Khurana S, Schmitt N, Kurup P, Mueller C, Obermoser G, Palucka AK, Albrecht RA, Garcia-Sastre A, Golding H, et al.: ICOS(+)PD-1(+)CXCR3(+) T follicular helper cells contribute to the generation of high-avidity antibodies following influenza vaccination. *Sci Rep* 2016, 6:26494. [PubMed: 27231124]
- *. This study documented the presence of a diverse repertoire of circulating Tfh and, additionally, found a correlate between the expansion of an activated pool of circulating Tfh and increases in affinity antibody.

86. Herati RS, Muselman A, Vella L, Bengsch B, Parkhouse K, Del Alcazar D, Kotzin J, Doyle SA, Tebas P, Hensley SE, et al.: Successive annual influenza vaccination induces a recurrent oligoclonotypic memory response in circulating T follicular helper cells. *Sci Immunol* 2017, 2.
87. Nayak JL, Alam S, Sant AJ: Cutting edge: Heterosubtypic influenza infection antagonizes elicitation of immunological reactivity to hemagglutinin. *J Immunol* 2013, 191:1001–1005. [PubMed: 23794632]
88. Spensieri F, Borgogni E, Zedda L, Bardelli M, Buricchi F, Volpini G, Fragapane E, Tavarini S, Finco O, Rappuoli R, et al.: Human circulating influenza-CD4⁺ ICOS1+IL-21⁺ T cells expand after vaccination, exert helper function, and predict antibody responses. *Proc Natl Acad Sci U S A* 2013, 110:14330–14335. [PubMed: 23940329]
89. Koutsakos M, Wheatley AK, Loh L, Clemens EB, Sant S, Nussing S, Fox A, Chung AW, Laurie KL, Hurt AC, et al.: Circulating TFH cells, serological memory, and tissue compartmentalization shape human influenza-specific B cell immunity. *Sci Transl Med* 2018, 10.
90. Lillie PJ, Berthoud TK, Powell TJ, Lambe T, Mullarkey C, Spencer AJ, Hamill M, Peng Y, Blais ME, Duncan CJ, et al.: Preliminary assessment of the efficacy of a T-cell-based influenza vaccine, MVA-NP+M1, in humans. *Clin Infect Dis* 2012, 55:19–25. [PubMed: 22441650]
91. Pleguezuelos O, Robinson S, Fernandez A, Stoloff GA, Mann A, Gilbert A, Balaratnam G, Wilkinson T, Lambkin-Williams R, Oxford J, et al.: A Synthetic Influenza Virus Vaccine Induces a Cellular Immune Response That Correlates with Reduction in Symptomatology and Virus Shedding in a Randomized Phase Ib Live-Virus Challenge in Humans. *Clin Vaccine Immunol* 2015, 22:828–835. [PubMed: 25994549]
92. Darton TC, Blohmke CJ, Moorthy VS, Altmann DM, Hayden FG, Clutterbuck EA, Levine MM, Hill AV, Pollard AJ: Design, recruitment, and microbiological considerations in human challenge studies. *Lancet Infect Dis* 2015, 15:840–851. [PubMed: 26026195]
- *. An outstanding discussion of the multitude of factors that are important to consider in the design of human challenge studies.
93. Oxford JS: Towards a universal influenza vaccine: volunteer virus challenge studies in quarantine to speed the development and subsequent licensing. *Br J Clin Pharmacol* 2013, 76:210–216. [PubMed: 23617282]
94. McKinstry KK, Strutt TM, Kuang Y, Brown DM, Sell S, Dutton RW, Swain SL: Memory CD4⁺ T cells protect against influenza through multiple synergizing mechanisms. *J Clin Invest* 2012, 122:2847–2856. [PubMed: 22820287]
95. Brown DM, Lee S, Garcia-Hernandez Mde L, Swain SL: Multifunctional CD4 cells expressing gamma interferon and perforin mediate protection against lethal influenza virus infection. *J Virol* 2012, 86:6792–6803. [PubMed: 22491469]
96. Teijaro JR, Turner D, Pham Q, Wherry EJ, Lefrancois L, Farber DL: Cutting edge: Tissue-retentive lung memory CD4 T cells mediate optimal protection to respiratory virus infection. *J Immunol* 2011, 187:5510–5514. [PubMed: 22058417]
97. Donnarumma T, Young GR, Merckenschlager J, Eksmond U, Bongard N, Nutt SL, Boyer C, Dittmer U, Le-Trilling VT, Trilling M, et al.: Opposing Development of Cytotoxic and Follicular Helper CD4 T Cells Controlled by the TCF-1-Bcl6 Nexus. *Cell Rep* 2016, 17:1571–1583. [PubMed: 27806296]
- *. Describes that global transcriptional profiles at the single cell level of Tfh and CD4 CTL illustrating the key roles of BCL-6 and Tcf7. Use of two viruses that have distinct tendencies to elicit Tfh vs. ThCTL, and perturbation of the function of inhibitory receptors LAG-3 and PD-1 point out the multiple layers of control in these opposing CD4 T cell differentiation programs. These studies as well earlier studies on the role of signal strength, emphasize that different pathogens and/or APC used for priming may play a critical role T cell fate decisions.
98. He W, Chen CJ, Mullarkey CE, Hamilton JR, Wong CK, Leon PE, Uccellini MB, Chromikova V, Henry C, Hoffman KW, et al.: Alveolar macrophages are critical for broadly-reactive antibody-mediated protection against influenza A virus in mice. *Nat Commun* 2017, 8:846. [PubMed: 29018261]
99. Wohlbold TJ, Podolsky KA, Chromikova V, Kirkpatrick E, Falconieri V, Meade P, Amanat F, Tan J, tenOever BR, Tan GS, et al.: Broadly protective murine monoclonal antibodies against influenza

B virus target highly conserved neuraminidase epitopes. *Nat Microbiol* 2017, 2:1415–1424. [PubMed: 28827718]

100. Tan GS, Leon PE, Albrecht RA, Margine I, Hirsh A, Bahl J, Krammer F: Broadly-Reactive Neutralizing and Non-neutralizing Antibodies Directed against the H7 Influenza Virus Hemagglutinin Reveal Divergent Mechanisms of Protection. *PLoS Pathog* 2016, 12:e1005578. [PubMed: 27081859]
101. DiLillo DJ, Palese P, Wilson PC, Ravetch JV: Broadly neutralizing antiinfluenza antibodies require Fc receptor engagement for in vivo protection. *J Clin Invest* 2016, 126:605–610. [PubMed: 26731473]
- *. These studies describe the importance of immunoglobulin Fc receptor interactions that are employed for protective effects induced by non-neutralizing antibodies to influenza.
102. Wohlbold TJ, Nachbagauer R, Xu H, Tan GS, Hirsh A, Brokstad KA, Cox RJ, Palese P, Krammer F: Vaccination with adjuvanted recombinant neuraminidase induces broad heterologous, but not heterosubtypic, cross-protection against influenza virus infection in mice. *MBio* 2015, 6:e02556. [PubMed: 25759506]
- *. With reference 99, these studies provide evidence that neuraminidase, used as a vaccine antigen, can elicit broadly reactive antibodies that can be protective. Although unable to prevent infection, these antibodies diminish the morbidity caused by influenza infection” These results suggest that CD4 T cells specific for NA may also be critical in enhancing protective antibody responses.
103. Mastelic Gavillet B, Eberhardt CS, Auderset F, Castellino F, Seubert A, Tregoning JS, Lambert PH, de Gregorio E, Del Giudice G, Siegrist CA: MF59 Mediates Its B Cell Adjuvanticity by Promoting T Follicular Helper Cells and Thus Germinal Center Responses in Adult and Early Life. *J Immunol* 2015, 194:4836–4845. [PubMed: 25870238]
104. Pilkinton MA, Nicholas KJ, Warren CM, Smith RM, Yoder SM, Talbot HK, Kalams SA: Greater activation of peripheral T follicular helper cells following high dose influenza vaccine in older adults forecasts seroconversion. *Vaccine* 2017, 35:329–336. [PubMed: 27919633]
- *. This suggests that high dose vaccination in the elderly results in improved expansion of activated Tfh, leading to greater seroconversion in this high risk population.
105. Kato Y, Zaid A, Davey GM, Mueller SN, Nutt SL, Zotos D, Tarlinton DM, Shortman K, Lahoud MH, Heath WR, et al.: Targeting Antigen to Clec9A Primes Follicular Th Cell Memory Responses Capable of Robust Recall. *J Immunol* 2015, 195:1006–1014. [PubMed: 26101322]
106. Yamasaki S, Shimizu K, Kometani K, Sakurai M, Kawamura M, Fujii SI: In vivo dendritic cell targeting cellular vaccine induces CD4(+) Tfh cell-dependent antibody against influenza virus. *Sci Rep* 2016, 6:35173. [PubMed: 27739478]
107. Allie SR, Randall TD: Pulmonary immunity to viruses. *Clin Sci (Lond)* 2017, 131:1737–1762. [PubMed: 28667071]
108. Zens KD, Chen JK, Farber DL: Vaccine-generated lung tissue-resident memory T cells provide heterosubtypic protection to influenza infection. *JCI Insight* 2016, 1.
109. Nish SA, Zens KD, Kratchmarov R, Lin WW, Adams WC, Chen YH, Yen B, Rothman NJ, Bhandoola A, Xue HH, et al.: CD4+ T cell effector commitment coupled to self-renewal by asymmetric cell divisions. *J Exp Med* 2017, 214:39–47. [PubMed: 27923906]
110. Strutt TM, Dhume K, Finn CM, Hwang JH, Castonguay C, Swain SL, McKinstry KK: IL-15 supports the generation of protective lung-resident memory CD4 T cells. *Mucosal Immunol* 2017.
111. Bautista BL, Devarajan P, McKinstry KK, Strutt TM, Vong AM, Jones MC, Kuang Y, Mott D, Swain SL: Short-Lived Antigen Recognition but Not Viral Infection at a Defined Checkpoint Programs Effector CD4 T Cells To Become Protective Memory. *J Immunol* 2016, 197:3936–3949. [PubMed: 27798159]
- *. This study suggests that second “hits” of antigen-specific CD4 T cells promotes memory, raising the possibility that persistent and tissue-localized antigen may be important feature of the most protective influenza vaccines.
112. van Dinther D, Stolk DA, van de Ven R, van Kooyk Y, de Gruijl TD, den Haan JMM: Targeting C-type lectin receptors: a high-carbohydrate diet for dendritic cells to improve cancer vaccines. *J Leukoc Biol* 2017, 102:1017–1034. [PubMed: 28729358]

113. Park CG: Vaccine strategies utilizing C-type lectin receptors on dendritic cells in vivo. *Clin Exp Vaccine Res* 2014, 3:149–154. [PubMed: 25003088]
114. Hoving JC, Wilson GJ, Brown GD: Signalling C-type lectin receptors, microbial recognition and immunity. *Cell Microbiol* 2014, 16:185–194. [PubMed: 24330199]
115. Duluc D, Joo H, Ni L, Yin W, Upchurch K, Li D, Xue Y, Klucar P, Zurawski S, Zurawski G, et al.: Induction and activation of human Th17 by targeting antigens to dendritic cells via dectin-1. *J Immunol* 2014, 192:5776–5788. [PubMed: 24835401]
116. Leon B, Ballesteros-Tato A, Randall TD, Lund FE: Prolonged antigen presentation by immune complex-binding dendritic cells programs the proliferative capacity of memory CD8 T cells. *J Exp Med* 2014, 211:1637–1655. [PubMed: 25002751]
117. Kim TS, Hufford MM, Sun J, Fu YX, Braciale TJ: Antigen persistence and the control of local T cell memory by migrant respiratory dendritic cells after acute virus infection. *J Exp Med* 2010, 207:1161–1172. [PubMed: 20513748]
118. Jelley-Gibbs DM, Brown DM, Dibble JP, Haynes L, Eaton SM, Swain SL: Unexpected prolonged presentation of influenza antigens promotes CD4 T cell memory generation. *J Exp Med* 2005, 202:697–706. [PubMed: 16147980]
119. Misumi I, Alirezai M, Eam B, Su MA, Whitton JL, Whitmire JK: Differential T cell responses to residual viral antigen prolong CD4+ T cell contraction following the resolution of infection. *J Immunol* 2013, 191:5655–5668. [PubMed: 24146043]
120. Turner DL, Cauley LS, Khanna KM, Lefrancois L: Persistent antigen presentation after acute vesicular stomatitis virus infection. *J Virol* 2007, 81:2039–2046. [PubMed: 17151119]
121. Mikhak Z, Strassner JP, Luster AD: Lung dendritic cells imprint T cell lung homing and promote lung immunity through the chemokine receptor CCR4. *J Exp Med* 2013, 210:1855–1869. [PubMed: 23960189]

Highlights

Humans have a diverse repertoire of circulating CD4 T cells specific for influenza viral proteins

Critical CD4 T cell functions include help for B cell and CD8 T cell responses and cytotoxicity

Although follicular helper cells and cytotoxic CD4 T cells are critical, their differentiation is reciprocally regulated

Universal influenza vaccines should be designed overcome deficits in CD4 T cell function and specificity

Figure 1. The range of abundance and functional capacity of influenza-specific human CD4 T cells within different subjects.

Existing data from PBMC from healthy donors indicates a broad range in frequency of influenza-specific CD4 T cells. Shown here for illustration are the differences depicted as bar heights, reflecting the total abundance of influenza specific CD4 T cells. The frequency of each functionally distinct subset of cells are indicated by different colored segments within each bar. This theoretical representation reflects the extreme view that there are non-overlapping subsets of CD4 T cells that convey each of indicated functions. Thus far, only the strict dichotomy between cytolytic cells and Tfh cells is supported by their unique and opposing transcription factors [97]. In the example shown, subject 2 and subject 4 have similar frequencies of influenza-specific CD4 T cells, but may be poised differently for protective immunity. Subject 2, having more abundant cytolytic cells might be best protected from infection by eliminating infected cells, while subject 4, with high frequency of Tfh cells, would exhibit the more robust antibody response. The right pie diagrams illustrate that in provision of help for B cell responses, in Subject 4, the antigen specificity of the CD4 T cells is critically important in providing help [16–18]. The neutralizing antibody response to vaccination would likely vary, depending on the abundance of HA-specific Tfh cells drawn into the response, with different outcomes depending on whether Subject 4 had high (top) or low (bottom) levels of HA-specific CD4 T cells. It is not yet known whether other functions of CD4 T cells, such as recruitment of innate effectors to the lung, or cytotoxicity, tracks with their antigen specificity.