Review articles

Scuba diving and otology: a systematic review with recommendations on diagnosis, treatment and post-operative care

Devon M Livingstone, Kristine A Smith, Beth Lange

Division of Otolaryngology – Head and Neck Surgery, Department of Surgery, University of Calgary, Alberta, Canada

Corresponding author: Devon Livingstone MD, Division of Otolaryngology – Head and Neck Surgery, Department of Surgery, University of Calgary, 7th floor, 4448 Front Street SE, Calgary, Alberta, Canada, T3M 1M4
dmliving@ucalgary.ca

Abstract

(Livingstone DM, Smith KA, Lange B. Scuba diving and otology: a systematic review with recommendations on diagnosis, treatment and post-operative care. *Diving and Hyperbaric Medicine*. 2017 June;47(2):97-109. doi: 10.28920/dhm47.2.97-109)

Scuba diving is a popular recreational and professional activity with inherent risks. Complications related to barotrauma and decompression illness can pose significant morbidity to a diver's hearing and balance systems. The majority of dive-related injuries affect the head and neck, particularly the outer, middle and inner ear. Given the high incidence of otologic complications from diving, an evidence-based approach to the diagnosis and treatment of otic pathology is a necessity. We performed a systematic and comprehensive literature review including the pathophysiology, diagnosis, and treatment of otologic pathology related to diving. This included inner, middle, and outer ear anatomic subsites, as well as facial nerve complications, *mal de débarquement* syndrome, sea sickness and fitness to dive recommendations following otologic surgery. Sixty-two papers on diving and otologic pathology were included in the final analysis. We created a set of succinct evidence-based recommendations on each topic that should inform clinical decisions by otolaryngologists, dive medicine specialists and primary care providers when faced with diving-related patient pathology.

Key words

Head and neck; ENT; Injuries; Medical conditions and problems; Barotrauma; Decompression sickness; Review article

Introduction

Over the past half century there has been a dramatic increase in the number of recreational scuba divers, with over nine million certified in the United States in 2015 and approximately 100,000 new divers per year.¹ While scuba diving is commonly viewed as a safe recreational activity, it exposes the participant to real risks of injury or even death. More than 80% of all diving complications occur in the head and neck.² Of these, approximately 65% are outer, middle or inner ear disorders.³ It is essential that clinicians understand the physiology and physics of scuba diving as well as the diagnosis, treatment and prevention of diving-related complications. However, there are currently few evidence-based recommendations or systematic reviews on this topic. The purpose of this report is to systematically review the current literature evaluating scuba diving physiology and complications pertinent to otology, and to provide a comprehensive resource with evidence-based recommendations where possible.

Two important scuba diving injuries are barotrauma and decompression sickness (DCS). To understand their pathophysiology, clinicians must understand the physics laws governing these injuries: Boyle's Law and Henry's Law.

BAROTRAUMA AND BOYLE'S LAW

Barotrauma is a pressure-mediated injury to tissue governed by Boyle's law. As a diver descends and the pressure increases, the volume of the gas compresses. This can result in a relative negative pressure in rigid- or semi-rigid-walled air-containing spaces in the body such as the middle ear or paranasal sinuses. Negative pressure can cause mucosal oedema, haemorrhage and even perforation if the space cannot equalize. On ascent, the volume of gas increases as the ambient pressure decreases. If an air-containing space cannot equalize with the surrounding pressure, the expanding volume of the gas may result in a variety of head and neck pathologies, including middle ear perforation or pneumocephalus from sinus barotrauma.⁴

DECOMPRESSION SICKNESS AND HENRY'S LAW

As a diver descends and ambient pressure increases, progressively higher pressure gas is delivered to the lung and more inert gas dissolves in the blood stream. The amount of inert gas that dissolves in a given tissue is proportional to the maximum depth and bottom time, as well as the perfusion and diffusion characteristics of that tissue. As a diver ascends, the additional inert gas load comes out of

solution at the level of the alveolus and is exhaled. If the rate of ascent exceeds the rate of alveolar gas exchange, inert gas will dissolve inside the diver, forming bubbles within the circulation and in tissues. The severity and nature of the DCS injury vary from mild systemic, musculoskeletal and cutaneous manifestations to severe, life-threatening central nervous and cardiorespiratory symptoms.

Literature search

A systematic search of the literature was performed using the following databases: Ovid/Medline, PubMed, EMBASE, UpToDate, Rubicon Repository and Cochrane Review Database up to January 2017. A screening literature search was used to identify all literature discussing scuba diving and any otolaryngology topics. Search terms included: “*scuba*” and/or “*diving*”, and “*head and neck*”, “*otolaryngolog**”, “*otolog**” “*rhinolog**”, “*sinus surgery*” or “*laryngolog**”. Reference lists of identified publications were reviewed to ensure no relevant studies in this field were missed. Grey literature, including the Diver’s Alert Network online resources, was also queried for completeness. Inclusion criteria included any paper discussing scuba diving and otology at any level of evidence (LOE). Exclusion criteria included papers that were not available in English or in an English translation. Given the limited amount of literature available, all studies meeting these inclusion criteria were included for completeness.

The combined search resulted in identification of 398 abstracts to be reviewed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.⁵ Screening resulted in 285 abstracts being excluded owing to duplications, leaving 113 abstracts to be reviewed. In assessing eligibility, 19 abstracts were excluded as they were not available in full text or in English, two because the topic did not include scuba diving, 20 as they did not discuss otology as it relates to diving and 23 as they solely discussed rhinology and oromaxillofacial topics. This left a total of 49 articles that met the criteria of including both scuba diving and otology topics. The works cited section of these articles were reviewed and 13 additional studies were also identified for review.

Included studies were evaluated and their LOE was noted based on a reported research methodology utilized by the Oxford Centre for Evidence Based Medicine (CEBM).⁶ After quality evaluation for each study, a summary was produced that included the aggregate grade of evidence and relevant recommendations. When there was only a single study available, an aggregate grade of evidence was not provided as grades are derived from the findings of multiple studies. Two authors (DML and KAS) reviewed the literature and produced the initial manuscript. A subsequent author (BL) was asked to review and critically appraise the recommendations based on the literature. Recommendations incorporate both the quality of evidence and the balance of benefit versus harm.

After additional review and synthesis of all information, a total of 74 articles were included in the study. There were no systematic reviews or meta-analyses found. The highest levels of evidence came from randomized controlled trials (Level 1b). The Appendix contains a summary of the 44 relevant papers that constituted primary literature related to otology and scuba diving. Eighteen papers were relevant but were either reviews, grey literature or individual case reports, and thus not included in the Appendix. The references section also contains 12 papers that did not specifically mention diving, but were relevant to the method used or contextualizing the relevant literature.

External ear

OTITIS EXTERNA

Otitis externa (OE) is the most common otologic disorder among divers, afflicting nearly half of all active divers on at least one occasion.⁷ *Pseudomonas aeruginosa* is the most common micro-organism responsible.⁸ Treatment is no different among divers versus non-divers and should consist of dry ear precautions, topical antimicrobial therapy (e.g., Ciprodex: three drops to affected ear BID x 7 days) and serial debridement of the external auditory canal. Divers should be counselled on the prevalence of this condition and may consider bringing antimicrobial drops on remote diving trips. Ethanol or acetic acid otic drops may be useful to prevent OE by drying the canal post-dive, and can be considered in divers prone to this condition.⁴

Recommendations (LOE)

- Divers should be counselled on the high prevalence of OE (5);
- Suggest bringing topical therapy on remote trips (5);
- Treatment similar to non-diving related OE (5).

EXOSTOSES

Exostoses are bony outgrowths of the external auditory canal (EAC) that typically occur bilaterally in scuba divers.⁹ They are very common, with a prevalence of up to 40% among professional divers,⁸ in comparison to about 6% among the general population living in coastal regions.¹⁰ Exostoses can lead to recurrent otitis externa, recurrent otorrhoea, cerumen impaction and conductive hearing loss from canal obstruction.^{8–10} If the canal is completely occluded, an asymmetrical caloric stimulus with vertigo can occur upon submersion.⁴ Divers with symptomatic exostoses should seek evaluation by an experienced otolaryngologist for consideration of canalplasty.⁹ Surgical indications for removal are identical to non-divers, with the caveat that they are very likely to experience progression of their exostoses due to further water exposure, and are at higher risk of OE.^{7,9} Exostoses should not affect fitness to dive unless they are occluding the canal or causing recurrent infections.¹¹

Recommendations (LOE)

- Exostoses can lead to recurrent otitis externa, recurrent otorrhoea, cerumen impaction and conductive hearing loss (4);
- Divers with symptomatic exostoses should seek evaluation by an experienced otolaryngologist for consideration of canalplasty (5);
- Exostoses should not affect fitness to dive unless they are symptomatic (5).

EXTERNAL AUDITORY CANAL BAROTRAUMA

External auditory canal barotrauma can also occur in the setting of occlusion due to cerumen impaction, foreign body, or a tight fitting wetsuit or drysuit hood. Severe exostoses can also cause cerumen impaction and lead to an isolated air space within the external auditory canal. Regardless of the aetiology of the predisposing canal occlusion, a relative vacuum is created on descent causing oedema and haemorrhagic vesiculation of the canal.¹² Treatment of EAC barotrauma consists of a short course of topical analgesic or steroid ear drops, and is similar to the treatment of OE.¹³ The aetiology of canal barotrauma should be understood by the clinician and diver to prevent the occurrence of subsequent episodes.

Recommendations (LOE)

- Treat with debridement and a short course of topical steroid/antibiotic (5);
- Refrain from diving until resolved (5);
- Treatment similar to OE (5);
- Identify and mitigate inciting aetiology of EAC barotrauma (5).

Middle ear**MIDDLE EAR BAROTRAUMA AND IMPAIRED EQUALIZATION**

Middle ear barotrauma (MEBt) occurs when there is dysequilibrium between the middle ear and ambient pressure, and accounts for up to 46% of patient presentations for diving-related head and neck pathology.¹⁴ During descent, middle ear pressure becomes progressively less than ambient pressure and the Eustachian tube (ET) must open to equilibrate these pressures. If the diver is unable to equalize and continues to descend beyond 1.4 metres depth, a pressure differential of greater than 90 mmHg is created, irreversibly blocking the ET.¹² The persistent negative pressure in the middle ear space can lead to extravasation of fluid and haemorrhage into the middle ear and tympanic membrane (TM) perforation. The resulting middle ear transudation and tubal oedema can impair middle ear ventilation on ascent, leading to a progression of barotrauma.¹⁵ MEBt with TM rupture can also occur during uncontrolled ascent, and is

associated with pulmonary barotrauma if a diver ascends with a closed glottis.¹⁶

MEBt is not always associated with inner ear sequelae. In a cohort of 67 professional divers, recurrent MEBt was not associated with sensorineural hearing loss.¹⁷ Risk factors for MEBt include poor ET function and poor mastoid pneumatization.^{18,19} In patients with tubal tonsillar hypertrophy and inability to equalize on descent, there may be a role for tuboplasty operations, including laser tuboplasty. In a series of nine divers, improved middle ear equalization was demonstrated postoperatively with seven being able to return to diving long-term.²⁰

Equalization of the middle ear space through the ET is active on descent, and passive on ascent. Importantly, equalization manoeuvre effectiveness is impacted by the diver's head position relative to their body. Clearing capacity is maximized in head-up positioning.²¹ Head-down position impairs passive equalization and should be avoided during descent.²² Multiple manoeuvres exist to assist pressure equalization of the middle ear space, including Valsalva, Frenzel, Toynbee, Lowry, Edmonds and voluntary tubal opening (French: *béance tubaire volontaire* – BTV) techniques. A forced Valsalva can be injurious to the inner ear, and the BTV manoeuvre is considered the least traumatic method.¹² A patient who can perform BTV is able to voluntarily contract their tensor veli palatine muscle and can often maintain ET patency on command. The Fédération Française d'Études et de Sports Sous-Marins has created a series of exercises to facilitate learning BTV. Divers with difficulty with equalization should descend at a slow rate and attempt to equalize pressure with every breath.

Importantly, diving at shallow depth is not useful to prevent MEBt as the greatest change in volume occurs near the water's surface; TM implosion can occur at depths as shallow as 1.2 m.¹² Rupture of the TM can lead to caloric stimulation of the vestibular apparatus, vertigo and disorientation underwater. This can be a highly dangerous scenario if it leads to a diver panicking.

Recommendations (LOE)

- Avoid head down position on ascent/descent (3b);
- Do not descend if unable to equalize, do not utilise the forced Valsalva (5);
- Ideally, utilize the BTV technique for equalisation (5);
- Optimise ET function and equalization technique (2b);
- Formal testing of ET function can be considered (5);
- Tuboplasty techniques may be helpful in specific ET dysfunction cases (4).

ALTERNOBARIC VERTIGO

Alternobaric vertigo (AV) is thought to arise from asymmetrical equalization of middle ear pressure transmitted

Figure 1

Implosive mechanism of inner ear barotrauma and perilymphatic fistula: a successful forced Valsalva manoeuvre (a) communicates excessive pressure to the middle ear space, distending the tympanic membrane, causing implosion of the round window (b) and subluxation of the stapes footplate (c)

Figure 2

Explosive mechanism of inner ear barotrauma and perilymphatic fistula: a failed forced Valsalva manoeuvre due to a blocked Eustachian tube (a) in the setting of a relative vacuum within the middle ear space (b) causes elevation of CSF pressure (c); this increased pressure is communicated to the inner ear, resulting in round window rupture (d) and intracochlear hemorrhage.

via the oval and round window membranes. AV is the most common cause of transient vertigo while diving,¹⁷ typically during ascent.⁴ Vertigo can last for several minutes and is often accompanied by nausea, vomiting and disorientation.²³ Caloric effects on the vestibular system can also cause transient vertigo and are pathophysiologically distinct from the dysequilibrium associated with nitrogen narcosis. ET dysfunction is a significant risk factor for the development of AV, and is considered an independent risk factor for experiencing an adverse event or accident while diving.²⁴ However, divers with AV were not at increased risk of experiencing a life-threatening or critical event while diving among a cohort of 63 recreational divers or 64 professional divers.^{23,24}

Risk factors for AV other than ET dysfunction include previous barotrauma, noise exposure during diving, cold water diving, a history of otitis media, a history of previous episodes of AV and female gender.^{23,25,26} AV is approximately four times more prevalent in females than in males.²³ Patients should be counselled on the aetiology and nature of AV as well as the potential risks of this condition, including aspiration and death. If a diver can tolerate the transient vertigo and pause their controlled ascent or descent until their vertigo resolves, the risk to the diver is minimized. Transient vertigo of any aetiology during technical or commercial diving portends a significantly higher risk of disorientation, accidental regulator dislodgement, scuba apparatus damage and asphyxiation.

Recommendations (LOE)

- Divers must understand the risks of disorientation, accidental regulator dislodgment, scuba apparatus damage and asphyxiation due to attacks of AV (2b);

- Recommend ET function optimization to minimize AV symptoms (5).

Inner ear

INNER EAR BAROTRAUMA

Inner ear barotrauma (IEBt) occurs when pressure changes within the middle ear are transmitted to the cochlea through the round and oval windows, and can occur through either implosive or explosive mechanisms (Figures 1 and 2). The oval window is somewhat more protected from pressure changes compared to the round window due to the added stability of the stapes footplate and associated ligamentous attachments. The stapes footplate can, however, be forcefully displaced by divers that employ forced Valsalva methods for middle ear equalization during descent. High pressure air is forced into the middle ear space, causing rapid outward displacement of the stapes and inward displacement and implosion of the delicate round window membrane (Figure 1). The pressure wave generated by this manoeuvre can cause cochlear haemorrhage with disruption of Reissner's membrane and the basilar membrane. Perilymphatic fistula (PLF) can also occur due to tearing of the round window membrane via an explosive mechanism of IEBt. Rapid fluctuations in the CSF pressure can be communicated to the perilymphatic space via the cochlear aqueduct or the lamina cribosa. A failed Valsalva attempt in the setting of ET dysfunction may elevate CSF pressure causing round window membrane rupture (Figure 2). Early studies and grey literature on diving noted an association between hearing loss and diving, even in the absence of IEBt. Importantly, high frequency sensorineural hearing loss is common among commercial divers, but is likely a noise-induced phenomenon from the use of underwater machinery.²⁷ High activity sport

divers do not experience any increased risk of SNHL.^{28,29}

Certain anatomic risk factors may predispose divers to IEBt. Patients may have an enlarged cochlear aqueduct orifice that can more readily transmit elevated CSF pressures to the delicate structures of the membranous labyrinth.³⁰ This may also be a consideration among paediatric divers who have a shorter cochlear aqueduct that can more easily transmit pressure fluctuations within the subarachnoid space.³¹ The approximate length of the cochlear aqueduct in a newborn is 3.5 mm and undergoes postnatal lengthening to 10 mm in an adult.³² The additional pressure may also cause rupture of Reissner's or the basilar membrane, leading to admixture of perilymph and endolymph.³³ Other anatomic risk factors for PLF are the potential communication between the inner and middle ear through the *fissula ante fenestrum* and enlarged vestibular aqueducts.^{30,34}

Diagnosis of IEBt can be difficult, with vague and variable symptom onset. Vestibular dysfunction and/or hearing impairment as a consequence of IEBt can occur on descent owing to impaired equalization, on ascent from expanding pneumolabyrinth, or even on non-diving days while straining or lifting.³⁵ The symptomatology and severity are dependent on the specific anatomical subsite affected; isolated basilar membrane tears can present with sensorineural hearing loss (SNHL) as the only presenting complaint.³⁶ Typically, patients present with a combination of vestibular and hearing deficits, and may complain of aural fullness, tinnitus and hyperacusis.³³ IEBt must be distinguished from inner-ear DCS, though these diagnoses are not mutually exclusive. A dangerous or atypical dive profile, diving using mixed gases or other symptoms of DCS elevates the likelihood of inner-ear DCS. Importantly, IEBt can occur in the absence of otoscopic findings of barotrauma, with a normal tympanic membrane and EAC.^{37,38} Clinical suspicion must remain high in these circumstances; delay between dive and symptom onset does not exclude the diagnosis of IEBt.³⁷ The dive profile must be thoughtfully interpreted and can be used to inform clinical suspicion of IEBt and other otologic pathologies.^{37, 39}

Treatment recommendations for IEBt include high-dose steroids (250 mg prednisolone for three days) with a taper for a total duration of therapy of 18 days, a recommendation based on expert opinion.² Surgical exploration is recommended when clinical suspicion of perilymphatic fistula is high, or the patient's hearing deteriorates despite appropriate conservative therapy. If a tympanotomy is undertaken, both the oval and round windows should be patched with fascia and absorbable surgical packing such as Surgifoam, even in the absence of active intraoperative perilymphatic leakage.³³ Intraoperative identification of a PLF can be facilitated using Trendelenburg positioning, or elevation of intrathoracic pressure in an intubated patient.³⁵ Conservative therapy for PLF should include bed rest, elevation of the head of the bed to 45°, the use of stool

softeners and avoidance of straining manoeuvres.¹³ A high-resolution CT scan of the temporal bone should also be performed in all patients that experience IEBt to rule out any anatomic factors that may predispose the patient to future episodes.³⁰

Patients should be counselled on the presence of anatomic risk factors that may influence their fitness to dive. Divers who suffer IEBt are often counselled to avoid diving. This recommendation may, however, be unnecessarily restrictive. A cohort of 21 patients who suffered IEBt and continued to dive against medical advice were counselled on middle-ear equalization techniques and methods to improve ET function. No further deterioration of inner-ear function was noted among these divers over a 1–12 year follow-up.⁴⁰ A recent comprehensive review identified five criteria for returning to diving following IEBt including stable hearing loss in a narrow frequency band, absence of vertigo/dysequilibrium, mitigation of risk factors for MEBt, no further anatomical risk factors present and no further surgical intervention required.³⁵

Recommendations (LOE)

- Patients with IEBt should undergo initial observation and medical therapy (3b) including corticosteroids (5);
- Exploratory tympanotomy and round/oval window patching should be performed if there is clinical deterioration, or high suspicion for a PLF (3b);
- A high-resolution CT scan of the temporal bones should be done to rule out anatomic risk factors (4);
- ET dysfunction and risk factors for MEBt should be mitigated (5);
- A fitness-to-dive assessment and consultation should be performed by an otolaryngologist or dive medicine specialist before more diving is undertaken (5).

INNER EAR DECOMPRESSION SICKNESS

Inner-ear DCS occurs according to the principles outlined earlier with regard to Henry's Law. The vestibular portion of the labyrinth appears to be more prone to damage than the cochlea, a phenomenon that is likely related to tissue perfusion and washout of inert gas. The vestibular apparatus has a higher tissue volume to blood supply ratio in comparison to the cochlea, leading to higher rates of local supersaturation and arterial microbubble load.⁴¹ Inner-ear DCS manifests with vertigo occurring within about two hours (h) of surfacing, with up to 40% of patients also experiencing some degree of hearing loss.⁴¹ Haemorrhage and protein deposition in the membranous labyrinth can eventually develop into fibro-osseous labyrinthitis.⁴² Inner-ear DCS can occur in isolation, or in combination with other CNS manifestations of DCS. Deep technical diving using He-O₂ and trimix breathing gases appears to confer a higher risk for inner-ear DCS.³³

Treatment of inner-ear DCS includes recompression with hyperbaric oxygen treatment (HBOT) as soon as possible; if the latency to HBOT for inner-ear DCS exceeds 5 h, approximately 90% of divers can expect some degree of permanent inner ear damage.⁴³ An animal model study has shown that precipitated material within the perilymph can appear as early as 1.5 h in squirrel monkeys with experimentally induced inner-ear DCS.⁴² Unfortunately the median time to treatment is often well in excess of these critical windows.² While waiting for HBOT, 100% oxygen should be administered during transport.⁴⁴ Adjunctive medical therapy can include steroids to reduce inflammatory oedema and low molecular weight dextran to improve microcirculation.⁴⁴ Doppler sonography should also be performed in these patients to rule out a cardiac or pulmonary right-to-left (R/L) shunt. Prevalence of a R/L shunt can be as high as 82% among inner-ear DCS patients compared with 25% in the normal population.⁴⁵ Patients with isolated R/L shunts may dive in accordance with 'low bubble diving' recommendations established by the Swiss Undersea Medical Society.⁴⁶ If the patient has suffered concurrent IEBt, emergent bilateral middle ear paracentesis or tympanostomy tube insertion followed by HBOT is recommended to prevent a worsening of symptoms.^{2,47}

Recommendations (LOE)

- Patients with inner-ear DCS should receive emergent HBOT as soon as possible (3b);
- Medical adjuncts including corticosteroids can be considered (5);
- If there is concern regarding concurrent IEBt, bilateral myringotomies with equalization tubes should be performed (4);
- Once the inner-ear DCS is adequately treated, Doppler ultrasonography to rule out a R/L shunt should be performed (3b);
- The aetiology of the DCS should be considered and the patient must be counselled on preventative measures (5).

Facial nerve pathology

Facial nerve paralysis as a complication of scuba diving is a rare condition that is hypothesized to be a consequence of reduced axonal capillary blood flow in the setting of a defect in the bony covering of the tympanic segment of the facial nerve. Initially this process is caused by negative middle ear pressure due to the relative vacuum created by inadequate equalization.⁴⁸ Secondary post-ischaemic intraneural swelling from transudate or blood may perpetuate axonal compression and ischaemia.¹⁵ Direct compression of the nerve can also occur due to trapped haemotympanum.⁴⁸ A bony defect of the tegmen, the bony separation between the cranial cavity and middle ear, can also result in pneumocephalus and associated intracranial complications.⁴⁹ Compressive neuropraxia may occur during ascent from expanding trapped gas within the

middle ear space. Air can also enter a normal facial nerve canal and cause compression by tracking along the chorda tympani nerve.⁵⁰ The degree of facial nerve injury should be documented and followed closely for recovery. Oral corticosteroid therapy may be useful to minimize further neuropraxia secondary to intraneural swelling. Facial nerve decompression is generally not indicated although, similar to surgical decompression following temporal bone trauma, there is great institutional variability whether or not this procedure is performed. A simple myringotomy can be performed to evacuate any haemotympanum and relieve the middle ear over-pressure and consequent facial nerve compression.⁴⁸

Recommendations (LOE)

- Corticosteroid therapy is recommended to minimize neuropraxia secondary to direct nerve compression (5);
- There is no consensus on the role of facial nerve decompression (5);
- Perform myringotomy to relieve middle ear over pressurization (4).

Sea sickness

Sea sickness, a type of motion sickness, occurs when there is a mismatch between vestibular, proprioceptive and visual inputs.⁴ In diving, this typically occurs on a boat heading to a dive site, while a diver is attached to a shot line during decompression, or during a prolonged stay on a liveaboard diving vessel.^{4,51} Divers are typically less susceptible to sea sickness once underwater, which can promote hasty and poorly executed water entries among those affected.⁴ Almost everyone is susceptible to sea sickness, but tolerance can develop from episodic exposure and after two to three days of continuous open-ocean exposure.⁴ Symptoms include a non-vertiginous sense of dysequilibrium, nausea, increased salivation, flushing, diaphoresis and general malaise.⁵² Risk factors include female gender,⁵³ altered vestibular or visual sensory cues,⁵⁴ history of migraine⁵⁵ and hormonal effects of pregnancy and oral contraception.^{52,56} Psychosocial factors also play a role; naval cadets were at lower risk of developing sea sickness when told they were unlikely to experience it.⁵⁷ Specific environmental modifications are useful for treatment and prevention of sea sickness. Lying supine while inside a ship may decrease susceptibility to motion sickness.⁵¹ Standing on deck and staring at the horizon will lessen the degree of vestibular-visual mismatch and improves symptoms of sea sickness. Medications for treatment and prevention of sea sickness are typically antihistamines or anticholinergics. Ondansetron, a 5-HT antagonist, and droperidol, a dopamine antagonist, are typically less effective for motion sickness.⁴ Scopolamine, an anticholinergic medication given trans-dermally every 72 h, has been the subject of three randomized, prospective double blind studies on motion sickness. It has been shown to be more effective in preventing motion sickness than

promethazine, meclizine, and lorazepam and has shown superiority over cinnarizine for sea sickness prevention in naval crew.^{58,59} Caffeine may also be of benefit when combined with other anti-nauseants for motion sickness; caffeine plus promethazine was shown to be superior to scopolamine in preventing air sickness among helicopter passengers.⁶⁰ Importantly, scopolamine has not been shown to affect cognitive performance or manual dexterity in hyperbaric chamber dives.⁶¹ Dimenhydrinate, however, has been shown to have significant cognitive side effects at therapeutic doses necessary to treat experimental motion sickness, and thus is likely unsuitable for diving.⁶² In animal studies neither scopolamine nor cinnarizine increased the risk of CNS oxygen toxicity.^{63,64}

Recommendations (LOE)

- Environmental modifications such as lying supine or visual fixation on the horizon should improve symptoms of sea sickness (4);
- Transdermal scopolamine q.72 h is superior to cinnarizine for prevention of sea sickness (1b), and superior to other anti-nauseant medications for motion sickness prevention (1b);
- Caffeine may be a useful adjunct for motion sickness medications; promethazine plus caffeine is more effective than scopolamine alone (1b);
- Caution should be exercised when diving while under the influence of any of these medications, and sedating medications should be avoided (5);
- Divers should not dive while nauseated due to the risk of vomiting underwater and subsequent asphyxiation (5).

Mal de débarquement

Mal de débarquement (MdDS) is the sensation of swaying or rocking movement with dysequilibrium that occurs in individuals upon returning to land after an extended period of time in a boat. Most patients have resolution of symptoms within minutes to days. Of 236 sailors, 73% experienced MdDS symptoms for up to 24 h (mean 2.6 h).⁶⁵ In some people, symptoms are persistent for months to years and are associated with significant fatigue, anxiety and cognitive impairment. Risk factors include female gender and previous history of seasickness.⁴⁷ Human and primate studies have implicated central maladaptation of the vestibulo-ocular reflex as a primary cause of MdDS.^{66,67} Treatment for MdDS is generally considered to be ineffective, though recent work focusing on readaptation of the vestibulo-ocular reflex (VOR) has shown some promise. In one study, 24 subjects with persistent MdDS (mean duration 19.1 months) were treated by rolling the head side-to-side while being exposed to a full-field optokinetic stimulus in an attempt to readapt the VOR. Head roll frequency was matched with the subjective oscillations felt by the subjects, with direction of optic kinetic rotatory stimulus based on the Fukada stepping test or a patient's subjective sense of rotation. Seventeen

subjects showed cure or substantial improvement at one-year follow-up. Other proposed therapies include vestibular rehabilitation, and use of long-acting benzodiazepines and tricyclic antidepressants.⁶⁸

Recommendations (LOE)

- Novel techniques to alter a maladapted VOR in MdDS may prove to be useful (2b);
- Recommend trial of medical therapy, and vestibular physiotherapy as adjuncts to refractory cases (5).

Post-operative recommendations

TYMPANOSTOMY TUBES

When diving with tympanostomy tubes at sufficient depth or pressure, the surface tension of water at the tube orifice will be exceeded and water will enter the middle ear space. This can generate an asymmetrical caloric effect, leading to transient vertigo and dysequilibrium. There is also an increased risk of otitis media from water freely entering the middle ear leading to middle-ear sequelae and early tube extrusion. Patients should be counselled on these serious risks if they are considering diving with tympanostomy tubes, and may be considered unfit to dive. Some authors advocate use of one-way membrane (Castelli-type) tympanostomy tubes to prevent these sequelae.¹² A scuba diving mask that keeps water out of both ears, such as the ProEar 2000, may also be useful in this patient population.¹² The ProEar mask is built with a tube connecting the ear covers to the facial portion of the mask. This ensures that the air space within the EAC remains equal to ambient pressure, preventing EAC barotrauma.

TYMPANOPLASTY

The integrity of the TM while diving is closely related to its compliance. Sclerotic, immobile membranes may impair middle ear equalization, while a monomeric, hypermobile TM may be at higher risk for rupture during a forced Valsalva. Patients who have previously undergone tympanoplasty may also have impaired ET function at baseline and be at higher risk for associated barotrauma. Safety to return to diving should be considered on a case-by-case basis, and a trial of diving at depth in a controlled environment such as a swimming pool should be considered.⁶⁹

STAPEDECTOMY

Stapes surgery was thought to increase the risk of IEBt, owing to an iatrogenic predisposition for perilymphatic fistula formation. However, no increased risk of otologic insult was found among large cohorts of divers who had undergone stapedectomy.^{70,71} Postoperative recommendations include dry-ear precautions for the first three weeks, and allow diving one month postoperatively, assuming normal ET function.

Appendix

Primary literature on otology and scuba diving; AV – alternobaric vertigo; CRS – Chronic rhinos sinusitis; ENG – electronystagmography; ETD – ET dysfunction; ET – Eustacian tube; HBOT – hyperbaric oxygen treatment; IEBt – inner ear barotrauma; IEDCS – inner-ear decompression sickness; MEBt – middle ear barotrauma; NIHL – noise-induced hearing loss;

OM – otitis media; PLF – perilymphatic fistula; PTA – pure tone audiometry; RCT – randomised controlled trial; SCC – semicircular canal;

SHA – sinusoidal harmonic acceleration testing; SNHL – sensorineural hearing loss; TMJ – temporo-mandibular joint; VOR – vestibulo-ocular reflex

First Author	LOE	n	Study design	Methods	Results	Conclusions/comment
External ear						
DiBartolomeo ¹⁰	4	70	Retrospective cohort review	Chart review for exostoses	Incidence 6.36/1,000	Epidemiology and risk factors for exostoses identified
Middle ear						
Antonelli ⁷²	5	11	Retrospective post-mortem	18 temporal bones from 11 divers: 8 rapid ascent; 3 drowning, no rapid ascent	Middle ear haemorrhage in all rapid ascent bones; 5/11 rapid ascent TMs perforated	General commentary on histology of temporal bone injury due to rapid ascent
Ivarsson ²¹	4	3	Case series	ET clearance in various head positions	Supine positioning impaired equalization on ascent and descent	Assuming an upright body posture during exposures to pressure changes is ideal
Jumah ²⁰	4	9	Prospective, non-consecutive case series	Divers with ET laser tuboplasty; pre- and post-op ET function/opening pressure	All subjects restored ability for pressure equalization; 7/9 able to resume diving	Divers with ETD may benefit from minimally invasive laser tuboplasty
Money ¹⁶	4	3	Case series	Postmortem histology of temporal bones from recently deceased divers	MEBt found in pulmonary barotrauma patient; IEDCS case had neo-ossification in SCCs	Breath holding causes MEBt; maladaptive neo-ossification occurs following IEDCS
Ornhagen ²²	3b	7	Case control series	Dry and wet pressure chambers; measured passive ET opening vs head position	Horizontal position and prone position equivalent; head-down position worse	Avoid head down position on ascent
Roydhouse ³	3b	650	Retrospective cohort study	Review of patients with diving-related ENT pathology	64.6% ear, 23.9% teeth/TMJ, 3.1% nose, 6.6% paranasal sinuses	Most common preventable cause of diving pathology is reversible nasal congestion
Uzun ¹⁸	2b	24	Prospective cohort study	Survey, CT scan temporal bones of mastoid pneumatization	MEBt occurred in 15 ears of 11 divers; pneumatization related to risk of MEBt	Inverse relationship between degree of pneumatization and risk of barotrauma
Uzun ¹⁹	3b	31	Retrospective cohort study	History, mastoid pneumatization, ET function	MEBt in 19 ears of 14 divers; ETD higher in divers	ETD measured by 9-step test and small mastoid are risk factors for MEBt
Vertigo						
Caruso ⁷³	4	14	Retrospective, consecutive case series	Audiometry, vestibular testing and DCS questionnaire	11/14 abnormal otologic, neuro-otologic findings	Expanded classification for vertigo related to diving
Kitajima ⁷⁴	3b	64	Non-consecutive cohort study	Sonotubometry and impedance tests	Diving incident group had significantly worse ETD	Divers with ETD prone to AV; check ET function pre dive
Klingmann ²³	2b	63	Retrospective cohort	Questionnaire, otoscopy, audiometry, caloric testing, ABR and MRI, etc	Higher prevalence of AV and ETD in females	Female divers 4x more likely to suffer AV; no increased risk for death if you experience AV
Molvaer ²⁵	3b	194	Retrospective cohort	Audiometry, interview of professional divers	39% had vertigo; 33% from AV; risk factors: previous barotrauma, noise exposure, cold water diving	Manage risk factors for AV while diving; AV did not cause any critical situations; theoretical risk remains

Appendix (cont.)

First Author	LOE	n	Study design	Methods	Results	Conclusions/comment
Uzun ²⁶	4	29	Retrospective case series	Survey, otoscopy, 9-step inflation/deflation tympanometry, ET function	Previous OM present in ¾ of AV divers; 14% of divers experienced AV	Risk factors for AV include previous history of OM and impaired ET function
Inner ear						
Cantaïs ⁴⁵	4	101	Case control, DCS/non-DCS divers	Transcranial Doppler	R/L shunt associated with increased incidence of cerebral DCI	Paradoxical emboli may be potential mechanism of cochleovestibular symptoms/ cerebral DCI
Edmonds ³⁷	4	50	Consecutive case series IEBt	Clinical manifestations, audiometry and treatment	17/50 Teed grade 0	Absence of TM haemorrhage does not exclude diagnosis of IEBt
Harill ⁷¹	5	231	Survey	Survey of otologists re: activity restrictions post stapes surgery	>50% recommend no diving; 32% had experience with postop Bt	No correlation between frequency of barotrauma reported and activity restriction
House ⁷⁰	5	22	Survey	22 divers post stapedectomy	3 otalgia on descent; 1 tinnitus; 1 transient vertigo; 1 PLF; no complaints related to labyrinthine injury	Stapedectomy does not increase the risk of IEBt in divers; dry ear precautions for 3/52, diving after 1/12 postop; ensure no ETD
Klingmann ²	4	46	Retrospective case series	Review of divers treated for inner ear DCS/barotrauma	Median latency to HBOT 10 hrs; 83% DCS patients had a R/L shunt	Divers with inner ear DCS associated with R/L shunting must undergo sonography
Klingmann ⁴¹	3b	34	Retrospective non-consecutive cohort	Divers with IEDCS were analysed re: symptomatology	All had vertigo; 40% had hearing loss; symptoms within 120 min of ascent; 73% had a R/L shunt	Vestibular symptoms predominate in IEDCS, likely due to the decreased tissue washout relative to the cochlea
Parell ⁴⁰	3b	20	Retrospective cohort, previous IEBt	Serial audiometry	No divers suffered a secondary bout of IEBt despite regular diving	Divers who suffer IEBt may continue to dive
Shupak ³³	3b	9	Retrospective case series	Cases of IEDCS/IEBt reviewed over 2 years	IEBt 5/9 divers; exploratory tympanotomy in 2/5; IEDCS 4/9, 7 underwent recompression	Consider IEDCS even when diving within no-stop limits; initial observation in IEBt, surgery if deterioration
Shupak ³⁰	4	2	Retrospective case series	Cases of IEBt were reviewed	PLF repair was successful; patients had anatomic risk factors on CT scan	CT temporal bone scans should be performed in IEBt to rule out anatomic risk factors
Shupak ³³	3b	25	Case-control series	PTA, ENG and VOR and SHA	PTA hearing threshold higher and decreased VOR phase values in divers	Slightly lower VOR values may represent adaptive underwater optokinetic changes
Shupak ⁴³	4	11	Retrospective consecutive case series	Audiometry, ENG, posturography, rotatory chair	10 IEDCS and 4 IEBt had cochleovestibular deficits	IEDCS has high risk for inner-ear sequelae even with HBOT; deficits common even if asymptomatic
Tal ⁴⁷	4	3	Retrospective case series	Review of IEDCS cases	Patients with IEDCS require HBOT	If doubt over diagnosis of IEDCS, recompression should still be undertaken
Wong ³⁹	4	8	Non-consecutive case series	Report on complaints and management of inner-ear symptoms in divers	See individual aetiologies highlighted in case series	Careful history and physical examination will help distinguish between IEBt and IEDCS

Appendix (cont.)

First Author	LOE	n	Study design	Methods	Results	Conclusions/comment
General articles						
Klingmann ¹⁴	2b	306	Retrospective cohort analysis	Chart review of anatomical sub-site/symptoms	8% external ear; 46% middle ear; 18% inner ear; 17% nose, paranasal sinuses; 8% DCS	Female divers significantly more affected by ETD; CRS associated with higher number of dives
Beckett ⁷	5	770	Internet survey of divers	Risk behaviours and safety practices	DCS symptoms in 53%; OE in approx. 50% divers	Injury more common in non-certified divers
Hearing						
Goplen ¹⁷	4	67	Prospective cohort study	Audiometry, ENG, posturography; 3/6 year follow up	Occupational SNHL in commercial divers; no change in ENG or posturography	No contribution of diving frequency to high frequency hearing loss; no evidence of permanent vestibular loss
Klingmann ²⁹	3b	123	Cross-sectional control comparison	PTA, tympanometry and otoscopy of sport. vs professional divers	No differences detected	No hearing loss among rec divers; professional divers risk occupational NIHL
Molvaer ²⁷	2b	116	Prospective cohort	Audiometry of professional divers over 6 years	Diver's hearing deteriorated faster than non-divers at comparable age	Noise exposure during commercial diving likely cause of hearing loss
Taylor ²⁸	3b	16	Retrospective cohort	Audiometry	No significant differences in audiological testing	Recreational divers not at increased risk of hearing loss
Sea sickness						
Cooper ⁵³	2b	1350	Prospective cohort on ocean liner	Epidemiologic study of sea sickness	Odds ratio for seasickness: female sex 2.95; young age 0.99; non crew 19.87	Sea sickness more common in women, may share common pathophysiology with migraine
Eden ⁵⁷	1b	25	RCT of naval cadets	Experimental group given positive self-efficacy/verbal placebo' during 5-day cruise	Experimental cadets reported less seasickness; rated as better performers	Seasickness may be a self-fulfilling prophecy, and may be reduced with 'verbal placebo'
Gahlinger ⁵¹	2b	260	Prospective cohort on cruise	Epidemiologic data and cabin location in ship recorded	Risk of motion sickness associated with age and sex	Location of cabin not associated with likelihood of motion sickness
Gil ⁵⁹	1b	76	RCT with crossover	Navy crew given transdermal scopolamine and cinnarizine, follow up questionnaires	Scopolamine more effective, less drowsiness, preferred agent, 41 vs 12%	Scopolamine patch should be considered drug of choice for treatment of seasickness
Williams ⁶¹	1b	24	RCT	Dexterity, arithmetic, sentence comprehension tested, varying depth and scopolamine dose	Dexterity and comprehension impaired at depth; no effects from scopolamine	Transdermal scopolamine during dive operations may be suitable; perform field testing
MdDS						
Dal ⁶⁶	5	5	Prospective	Videonystagmography and roll chair	Adaptation of the angular VOR generated through velocity storage	Central maladaptation of the VOR implicated in MdDS
Gordon ⁶⁵	2b	234	Retrospective cohort	Survey of MdDS among healthy crew members	73% had MdDS symptoms; <6 h in 93%	Transient MdDS common and expected among crew members

Appendix (cont.)

First Author	LOE	n	Study design	Methods	Results	Conclusions/comment
Animal studies						
Antonelli ¹⁵	5	16	Prospective case control animal model	Guinea pig stapedectomy model; consecutive hyperbaric dives, electrocochleography and hair cell counts	Middle ear barotrauma in 8 stapedectomy ears vs. 5 controls; no difference in hair cell counts	No predisposition to cochlear sequelae post stapedectomy in guinea pig model of barotrauma
Arieli ⁶⁴	5	26	Prospective case control animal model	Rats given cinnarizine/control O ₂ ; exposure protocols to explore CNS toxicity	Latency to first electrical discharge on EEG increased at high O ₂ pressures	Cinnarizine does not increase risk of CNS O ₂ toxicity
Bitterman ⁶³	5	36	Prospective case control animal model	EEGs and HR measured to test interaction of scopolamine with HBO	No difference in latent period between control and scopolamine rats	Scopolamine does not alter hyperoxic seizures
Landolt ⁴²	5	100	Case control animal model	Histology of squirrel monkey temporal bones	Histologic vestibular apparatus deficits post decompression exposure	Neo-ossification within otic fluid spaces post DCS could cause permanent deficit to diver

Conclusions

Scuba diving has a significant potential for complications in the external, middle and inner ear. Otolaryngologists and clinicians with an interest in dive medicine should have a keen understanding of the pathophysiology, treatment and fitness-to-dive implications of diving-related disorders of the head and neck. The recommendations within this review are intended to supplement good clinical judgment, and should be applied within the context of each individual patient's circumstance. Overall, there is a need for more high-quality research on diving-related head and neck pathology. Research in this area can lend insight into the pathophysiology of barotrauma and DCS while improving patient care and decreasing dive-related morbidity.

References

- Buzzacott P TB, Caruso JL, Nelson C, Denoble PJ, Nord DA, Chimiak J, et al. *DAN Annual Diving Report 2012–2015*. 2015 Edition. Durham NC: Divers Alert Network; 2015. p. 127.
- Klingmann C, Praetorius M, Baumann I, Plinkert PK. Barotrauma and decompression illness of the inner ear: 46 cases during treatment and follow-up. *Otol Neurotol*. 2007;28:447-54. doi: 10.1097/MAO.0b013e318030d356. PubMed PMID: 17417111.
- Roydhouse N. 1001 disorders of the ear, nose and sinuses in scuba divers. *Can J Appl Sport Sci*. 1985;10:99-103. PubMed PMID: 4017159.
- Edmonds CB, Lippmann, Mitchell S, editors. *Diving and subaquatic medicine*. Boca Raton, FL: CRC Press; 2015.
- Foster RL. Reporting guidelines: CONSORT, PRISMA, and SQUIRE. *J Spec Pediatr Nurs*. 2012;17(1):1-2. doi: 10.1111/j.1744-6155.2011.00319.x. PubMed PMID: 22188266.
- Howick J, Glasziou P, Greenhalgh T, Heneghan C, Liberati A, Moschetti I, et al. *The Oxford Levels of Evidence 2*. Oxford: Centre for Evidence-Based Medicine; 2011. [cited 2016 January]. Available from: <http://www.cebm.net/index.aspx?o=5653>.
- Beckett A, Kordick MF. Risk factors for dive injury: a survey study. *Res Sports Med*. 2007;15:201-11. doi: 10.1080/15438620701526779. PubMed PMID: 17987508.
- Azizi MH. Ear disorders in scuba divers. *Int J Occup Environ Med*. 2011;2:20-6. PubMed PMID: 23022815.
- House JW, Wilkinson EP. External auditory exostoses: evaluation and treatment. *Otolaryngol Head Neck Surg*. 2008;138:672-8. doi: 10.1016/j.otohns.2008.01.023. PubMed PMID: 18439477.
- DiBartolomeo J. Exostoses of the external auditory canal. *Ann Otol Rhinol Laryngol*. 1979;88(Suppl 61):2-20.
- Diver's Alert Network. *Surfer's Ear*. [cited 2017 January 07]. Available from: <https://www.diversalertnetwork.org/health/ears/surfers-ear>.
- Becker GD, Parell GJ. Barotrauma of the ears and sinuses after scuba diving. *Eur Arch Otorhinolaryngol*. 2001;258:159-63. PubMed PMID: 11407445.
- Lacey JP, Amedee RG. The otologic manifestations of barotrauma. *J La State Med Soc*. 2000;152:107-11. PubMed PMID: 10851823.
- Klingmann C, Praetorius M, Baumann I, Plinkert PK. Otorhinolaryngologic disorders and diving accidents:

- an analysis of 306 divers. *Eur Arch Otorhinolaryngol*. 2007;264:1243-51. doi: 10.1007/s00405-007-0353-6. PubMed PMID: 17639445.
- 15 Antonelli PJ, Adameczyk M, Appleton CM, Parell GJ. Inner ear barotrauma after stapedectomy in the guinea pig. *Laryngoscope*. 1999;109:1991-5. doi: 10.1097/00005537-199912000-00018. PubMed PMID: 10591361.
 - 16 Money KE, Buckingham IP, Calder IM, Johnson WH, King JD, Landolt JP, et al. Damage to the middle ear and the inner ear in underwater divers. *Undersea Biomed Res*. 1985;12:77-84. PubMed PMID: 4035820.
 - 17 Goplen FK, Gronning M, Aasen T, Nordahl SH. Vestibular effects of diving - a 6-year prospective study. *Occup Med (Lond)*. 2010;60:43-8. doi: 10.1093/occmed/kqp148. PubMed PMID: 19854795.
 - 18 Uzun C, Adali MK, Korten M, Yagiz R, Aydin S, Cakir B, et al. Relationship between mastoid pneumatization and middle ear barotrauma in divers. *Laryngoscope*. 2002;112:287-91. doi: 10.1097/00005537-200202000-00016. PubMed PMID: 11889385.
 - 19 Uzun C. Evaluation of pre-dive parameters related to ET dysfunction for symptomatic middle ear barotrauma in divers. *Otol Neurotol*. 2005;26:59-64. PubMed PMID: 15699720.
 - 20 Jumah MJ, Jumah MD, Sedlmaier B. Restoring middle ear pressure equalization in divers by laser Eustachian tuboplasty. *Undersea Hyperb Med*. 2013;40:299-306. PubMed PMID: 23789565.
 - 21 Ivarsson A, Lundgren CE, Tjernstrom O, Uddman R, Ornhaugen H. Eustachian tubal function during immersion. *Undersea Biomed Res*. 1977;4:9-18. PubMed PMID: 857360.
 - 22 Ornhaugen HC, Tallberg P. Pressure equilibration of the middle ear during ascent. *Undersea Biomed Res*. 1981;8:219-27. PubMed PMID: 7324254.
 - 23 Klingmann C, Knauth M, Praetorius M, Plinkert PK. Alternobaric vertigo – really a hazard? *Otol Neurotol*. 2006;27:1120-5. doi: 10.1097/01.mao.0000235373.78116.a8. PubMed PMID: 17130801.
 - 24 Kitajima N, Sugita-Kitajima A, Kitajima S. Altered ET function in SCUBA divers with alternobaric vertigo. *Otol Neurotol*. 2014;35:850-6. doi: 10.1097/MAO.0000000000000329. PubMed PMID: 24751737.
 - 25 Molvaer OI, Albrektsen G. Alternobaric vertigo in professional divers. *Undersea Biomed Res*. 1988;15:271-82. PubMed PMID: 3212844.
 - 26 Uzun C, Yagiz R, Tas A, Adali MK, Inan N, Korten M, et al. Alternobaric vertigo in sport SCUBA divers and the risk factors. *J Laryngol Otol*. 2003;117:854-60. doi: 10.1258/002221503322542845. PubMed PMID: 14670144.
 - 27 Molvaer OI, Albrektsen G. Hearing deterioration in professional divers: an epidemiologic study. *Undersea Biomed Res*. 1990;17:231-46. PubMed PMID: 2356593.
 - 28 Taylor DM, Lippmann J, Smith D. The absence of hearing loss in otologically asymptomatic recreational scuba divers. *Undersea Hyperb Med*. 2006;33:135-41. PubMed PMID: 16716064.
 - 29 Klingmann C, Knauth M, Ries S, Tasman AJ. Hearing threshold in sport divers: is diving really a hazard for inner ear function? *Arch Otolaryngol Head Neck Surg*. 2004;130:221-5. doi: 10.1001/archotol.130.2.221. PubMed PMID: 14967755.
 - 30 Shupak A. Recurrent diving-related inner ear barotrauma. *Otol Neurotol*. 2006;27:1193-6. doi: 10.1097/01.mao.0000231499.69404.22. PubMed PMID: 16983314.
 - 31 Marchese-Ragona R, Marioni G, Ottaviano G, Gaio E, Staffieri C, de Filippis C. Pediatric sudden sensorineural hearing loss after diving. *Auris Nasus Larynx*. 2007;34:361-4. doi: 10.1016/j.anl.2006.11.004. PubMed PMID: 17475428.
 - 32 De la Cruz A, Linthicum FH, Jr., Luxford WM. Congenital atresia of the external auditory canal. *Laryngoscope*. 1985;95:421-7. PubMed PMID: 3884941.
 - 33 Shupak A, Doweck I, Greenberg E, Gordon CR, Spitzer O, Melamed Y, et al. Diving-related inner ear injuries. *Laryngoscope*. 1991;101:173-9. doi: 10.1288/00005537-199102000-00013. PubMed PMID: 1992269.
 - 34 Kohut RI, Hinojosa R, Ryu JH. A temporal bone study method to evaluate clinical diagnostic criteria: permeability of patencies of the fissula ante fenestram. *Otolaryngol Head Neck Surg*. 1991;104:143-4. PubMed PMID: 1900619.
 - 35 Elliott EJ, Smart DR. The assessment and management of inner ear barotrauma in divers and recommendations for returning to diving. *Diving Hyperb Med*. 2014;44:208-22. PubMed PMID: 25596834.
 - 36 Love JT Jr, Waguespack RW. Perilymphatic fistulas. *Laryngoscope*. 1981;91:1118-28. PubMed PMID: 7242203.
 - 37 Edmonds C. Inner ear barotrauma: a retrospective clinical series of 50 cases. *SPUMS Journal*. 2004;34:11-4.
 - 38 Freeman P, Edmonds C. Inner ear barotrauma. *Arch Otolaryngol*. 1972;95:556-63. PubMed PMID: 4666425.
 - 39 Wong RW, Walker M. Diagnostic dilemmas in inner ear decompression sickness. *SPUMS Journal*. 2004;34:5-10.
 - 40 Parell GJ, Becker GD. Inner ear barotrauma in scuba divers. A long-term follow-up after continued diving. *Arch Otolaryngol Head Neck Surg*. 1993;119:455-7. PubMed PMID: 8457309.
 - 41 Klingmann C. Inner ear decompression sickness in compressed-air diving. *Undersea Hyperb Med*. 2012;39:589-94. PubMed PMID: 22400449.
 - 42 Landolt JP, Money KE, Topliff ED, Ackles KN, Johnson WH. Induced vestibular dysfunction in squirrel monkeys during rapid decompression. *Acta Otolaryngol*. 1980;90:125-9. PubMed PMID: 6969520.
 - 43 Shupak A, Gil A, Nachum Z, Miller S, Gordon CR, Tal D. Inner ear decompression sickness and inner ear barotrauma in recreational divers: a long-term follow-up. *Laryngoscope*. 2003;113:2141-7. doi: 10.1097/00005537-200312000-00017. PubMed PMID: 14660917.
 - 44 Talmi YP, Finkelstein Y, Zohar Y. Decompression sickness induced hearing loss. A review. *Scand Audiol*. 1991;20:25-8. PubMed PMID: 1842265.
 - 45 Cantais E, Louge P, Suppini A, Foster PP, Palmier B. Right-to-left shunt and risk of decompression illness with cochleovestibular and cerebral symptoms in divers: case control study in 101 consecutive dive accidents. *Crit Care Med*. 2003;31:84-8. doi: 10.1097/01.CCM.0000038040.42972.81. PubMed PMID: 12544998.
 - 46 Swiss Undersea Medical Society. *Low bubble diving*; 2015. [cited 2017 January 08]. Available from: https://www.aqua-med.eu/en/files/pdf/Low-Bubble-Diving_GB.pdf.
 - 47 Tal D, Domachevsky L, Bar R, Adir Y, Shupak A. Inner ear decompression sickness and mal de débarquement. *Otol Neurotol*. 2005;26:1204-7. PubMed PMID: 16272943.
 - 48 Carmichael ML, Boyev KP. Middle ear barotrauma causing transient facial nerve paralysis after scuba diving. *Diving Hyperb Med*. 2016;46:260-1. PubMed PMID: 27966206.
 - 49 Kamide D, Matsunobu T, Shiotani A. Facial baroparesis caused by scuba diving. *Case Rep Otolaryngol*. 2012;2012:329536. doi: 10.1155/2012/329536. PubMed PMID: 22953110.
 - 50 Brandt MT. Oral and maxillofacial aspects of diving medicine.

- Mil Med.* 2004;169:137-41. PubMed PMID: 15040636.
- 51 Gahlinger PM. Cabin location and the likelihood of motion sickness in cruise ship passengers. *J Travel Med.* 2000;7:120-4. PubMed PMID: 11179940.
 - 52 Gahlinger PM. Motion sickness. How to help your patients avoid travel travail. *Postgrad Med.* 1999;106:177-84. doi: 10.3810/pgm.1999.10.1.719. PubMed PMID: 10533517.
 - 53 Cooper C, Dunbar N, Mira M. Sex and seasickness on the Coral Sea. *Lancet.* 1997;350:892. doi: 10.1016/S0140-6736(05)62083-1. PubMed PMID: 9310639.
 - 54 Barabas G, Matthews WS, Ferrari M. Childhood migraine and motion sickness. *Pediatrics.* 1983;72:188-90. PubMed PMID: 6866603.
 - 55 Kayan A, Hood JD. Neuro-otological manifestations of migraine. *Brain.* 1984;107:1123-42. PubMed PMID: 6334543.
 - 56 Grunfeld E, Gresty MA. Relationship between motion sickness, migraine and menstruation in crew members of a "round the world" yacht race. *Brain Res Bull.* 1998;47:433-6. PubMed PMID: 10052571.
 - 57 Eden D, Zuk Y. Seasickness as a self-fulfilling prophecy: raising self-efficacy to boost performance at sea. *J Appl Psychol.* 1995;80:628-35. PubMed PMID: 7592232.
 - 58 Dornhoffer J, Chelonis JJ, Blake D. Stimulation of the semicircular canals via the rotary chair as a means to test pharmacologic countermeasures for space motion sickness. *Otol Neurotol.* 2004;25:740-5. PubMed PMID: 15354005.
 - 59 Gil A, Nachum Z, Tal D, Shupak A. A comparison of cinnarizine and transdermal scopolamine for the prevention of seasickness in naval crew: a double-blind, randomized, crossover study. *Clin Neuropharmacol.* 2012;35:37-9. doi: 10.1097/WNF.0b013e31823dc125. PubMed PMID: 22139622.
 - 60 Estrada A, LeDuc PA, Curry IP, Phelps SE, Fuller DR. Airsickness prevention in helicopter passengers. *Aviat Space Environ Med.* 2007;78:408-13. PubMed PMID: 17484344.
 - 61 Williams TH, Wilkinson AR, Davis FM, Frampton CM. Effects of transcutaneous scopolamine and depth on diver performance. *Undersea Biomed Res.* 1988;15:89-98. PubMed PMID: 3363755.
 - 62 Paule MG, Chelonis JJ, Blake DJ, Dornhoffer JL. Effects of drug countermeasures for space motion sickness on working memory in humans. *Neurotoxicol Teratol.* 2004;26:825-37. doi: 10.1016/j.ntt.2004.07.002. PubMed PMID: 15451046.
 - 63 Bitterman N, Eilender E, Melamed Y. Hyperbaric oxygen and scopolamine. *Undersea Biomed Res.* 1991;18:167-74. PubMed PMID: 1853467.
 - 64 Arieli R, Shupak A, Shachal B, Shenedrey A, Ertracht O, Rashkovan G. Effect of the anti-motion-sickness medication cinnarizine on central nervous system oxygen toxicity. *Undersea Hyperb Med.* 1999;26:105-9. PubMed PMID: 10372430.
 - 65 Gordon CR, Spitzer O, Shupak A, Doweck I. Survey of mal de débarquement. *BMJ.* 1992;304(6826):544. PubMed PMID: 1559057.
 - 66 Dai M, Raphan T, Cohen B. Adaptation of the angular vestibulo-ocular reflex to head movements in rotating frames of reference. *Exp Brain Res.* 2009;195:553-67. doi: 10.1007/s00221-009-1825-2. PubMed PMID: 19458941.
 - 67 Guedry FE Jr, Graybiel A. Compensatory nystagmus conditioned during adaptation to living in a rotating room. *J Appl Physiol.* 1962;17:398-404. PubMed PMID: 13902616.
 - 68 Hain TC, Hanna PA, Rheinberger MA. Mal de débarquement. *Arch Otolaryngol Head Neck Surg.* 1999;125:615-20. PubMed PMID: 10367916.
 - 69 Sim RJ, Youngs RP. Otolaryngological requirements for recreational self-contained underwater breathing apparatus (SCUBA) diving. *J Laryngol Otol.* 2007;121:306-11. doi: 10.1017/S0022215106001976. PubMed PMID: 17040582.
 - 70 House JW, Toh EH, Perez A. Diving after stapedectomy: clinical experience and recommendations. *Otolaryngol Head Neck Surg.* 2001;125:356-60. doi: 10.1067/mhn.2001.118183. PubMed PMID: 11593171.
 - 71 Harrill WC, Jenkins HA, Coker NJ. Barotrauma after stapes surgery: a survey of recommended restrictions and clinical experiences. *Am J Otol.* 1996;17:835-45; discussion 45-6. PubMed PMID: 8915410.
 - 72 Antonelli PJ, Adamczyk M, Appleton CM, Parell GJ. Inner ear barotrauma after stapedectomy in the guinea pig. *Laryngoscope.* 1999;109:1991-5. doi: 10.1097/00005537-199912000-00018. PubMed PMID: 10591361.
 - 73 Caruso VG, Winkelmann PE, Correia MJ, Miltenberger GE, Love JT. Otologic and otoneurologic injuries in divers: clinical studies on nine commercial and two sport divers. *Laryngoscope.* 1977;87(4 Pt 1):508-21. doi: 10.1288/00005537-197704000-00005. PubMed PMID: 300136.
 - 74 Kitajima N, Sugita-Kitajima A, Kitajima S. Altered eustachian tube function in SCUBA divers with alternobaric vertigo. *Otol Neurotol.* 2014;35:850-6. doi: 10.1097/MAO.0000000000000329. PubMed PMID: 24751737.

Funding and potential conflict(s) of interest: nil

Submitted: 11 July 2016; revised 16 January and April 14 2017

Accepted: 20 April 2017