Analysis of cervical smears in a Muslim population

To the Editor: In the article "Analysis of cervical smears in a Muslim population" in the May-June issue of the Annals, the authors report on abnormal smears in a Muslim population. In about 21 000 cases, a good number for evaluation, the prevalence was 0.5%, which they concluded was low compared with Western populations. In my opinion, the higher rate of abnormal smears in Western Society is due to several social factors, but this theory requires more study. Early sex (before marriage), even at an age as low as 13 years, is common in the West. This is rare in a Muslim population, but may rise with adaptation to Western culture. The use of oral contraceptives at very early ages and sometimes even spermicidal condoms occurs in Western populations. Having more than one sex partner at a time is more common. Uncircumcised husbands and sex partners in non-Muslim societies also increases the prevalence of cervical carcinoma, as proved by work in India and the Western world. Circumcision has been recommended in Islamic societies for many years as a part of Muslim teachings, but not in Western societies, although many adults in the West now get circumcised. Women of lower socioeconomic profile have a lower prevalence of abnormal cervical smears than middle and upper strata women because they do not believe in family planning, bottle feeding, and other social evils that prevail in the so-called modern societies. Finally, Islam allows men to marry four wives, but not women to marry four husbands.

Dr. Syed Inseram Ali
Technical Director
Zain Clinical Laboratory
Medina Munawara,
Saudi Arabia

Performance and accuracy of an immunodiagnostic antigen detection test in diagnosing *Plasmodium falciparum* among Yemeni patients

To the Editor: I read with interest the paper of Dr. A. A. Azazy, "Performance and accuracy of an immunodiagnostic test in diagnosing *Plasmodium falciparum* among Yemeni patients" in the January-February 2004 issue (page 50). I have two issues:

1. As the sensitivity is 94%, and means are not made for light microscopic examination to detect malignant malaria, don't you risk to kill 6% of patients if you don't mention that the dipstick rapid test OptiMAL cannot detect all cases and that it is up to the doctor to use his clinical judgment?

2. Malaria is prevalent in poor counties and these sophisticated methods are generally expensive.

I agree with the author that "further investigation to evaluate the test in a large population and in various endemic areas is still needed."

Khalil Shihab, MD, DTM&Hm MPH
Previous Malaria Director
Baghdad, Iraq

Partial recovery of the visual field scotoma upon early retinal photocoagulation and the medical therapy in a preeclamptic patient

To the Editor: Preeclampsia is a major cause of maternal and perinatal morbidity and mortality. It still accounts for 200 000 maternal deaths worldwide per year.¹ We report a case of preeclampsia with branch retinal vein occlusion with corresponding visual field scotoma which partially recovered with the early laser photocoagulation. To the best of our knowledge, the present case is the first to report the combination of these findings.

A 37-year-old, gravida 3 para 2, 31-week pregnant woman admitted to the emergency room with the complaint of headache, epigastric-right upper quadrant abdominal pain and vomiting. She had persistently high systolic and diastolic blood pressures. On physical examination, significant edema was detected in her legs over the tibia. She had no history of systemic or ocular disease, operation or trauma. She was hospitalized with the diagnosis of preeclampsia. Antihypertensive therapy with methyldopa 250 mg po bid was started. Maintenance of strict fluid balance, selective haemodynamic monitoring, and selective plasma volume expansion with close maternal and fetal monitoring were provided. In the second week, she complained of acute visual disturbances such as blurred vision and a visual acuity loss in her left eye. Her ophthalmological consultation revealed that she had a best-corrected visual acuity of 20/200 and 20/20 with normal color vision tests, in the left and right eyes, respectively. Both eyes revealed normal anterior segment findings with no rubeosis iridis. Fundus examination revealed segmental intraretinal hemorrhage, dilated and the tortuous retinal veins in both eyes. Moreover, the left eye had retinal neovascularisations (1/2 disc diameter) in the upper temporal sector with slight macular edema consistent with the diagnosis of the upper temporal retinal vein occlusion. Intraocular pressure readings were both found to be within normal limits. Orthoptic examination of the eyes revealed no deviation. Refractive values of the both eyes were "− 0.25 D". There were no anisocoria. Visual field tests performed with Humphrey Field Analyzer II using central 30-2, SITA Standard strategy showed a normal test in the right eye while absolute scotoma in the lower nasal quadrant in the left eye was noted. She refused fundus fluorescein angiography and we informed her of the possible risks to the fetus. She consulted with an internist for a probable underlying coagulopathy

and vasculopathy, namely lupus erythematosus. Overall, she was diagnosed as preeclampsic.

After obtaining written informed consent, scatter retinal photocoagulation was applied in the upper temporal region of the left fundus using 250 argon green laser bursts with a spot diameter of 500 micron, and with a duration of 0.1 seconds. We analysed her visual field testings three days and one week after the laser therapy. There was an apparent partial improvement in the scotoma region with an increase in the mean sensitivity and a decrease in the mean deviation of the visual field by the end of the first week. All of the visual field tests, before and after the laser treatment, were reliable. Visual acuity of the left eye improved to 40/200 at the first week examination, with no further improvement at the first month examination.

A variety of occlusive vascular disorders may occur during pregnancy. These disorders include a Purtscher-like retinopathy, or ocular changes associated with disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, and amniotic fluid embolism.² The most common retinal abnormality seen in preeclampsia is focal arteriolar spasm and narrowing.³ In the second week of admission, she developed acute branch retinal vein occlusion. Following retinal laser photocoagulation treatment, which was performed on the same day of the diagnoses, she was found to partially recover the absolute scotoma seen in computerised visual field analysis. Apaydin et al showed that mean sensitivity of the central visual field has improved significantly 3 months after panretinal laser photocoagulation treatment.⁴ Akar et al found that significant changes seen in optic rim parameters has lasted more than 4 months after laser therapy.⁵

Palacz and Sylwestrzak showed that early laser photocoagulation treatment is beneficial in the treatment of the retinal vein thrombosis and besides that, pharmacological treatment coupled

with laser photocoagulation gives much better results than the pharmacotherapy alone.⁶ On the other hand, it is a well-known fact that there is a learning curve with automated perimetry.⁷ The improvement noted in our case could be due to the improvement in the performance of the test. Preeclampsia, as involving multiple organ systems, still remains an important cause of perinatal and maternal mortality and morbidity in all countries of the world.⁸ It should be screened through multidisciplinary approach. It seems crucial to have comprehensive baseline dilated eye examination before conception and regular follow up ophthalmologic examinations during pregnancy.

Acknowledgment

This study was performed with the support of Akdeniz University scientific research unit.

Munire Erman Akar, MD

Cemil Apaydin, MD

Yusuf Akar, MD

Akdeniz Universitesi

Tip Fakultesi

Kadin Hastaliklari ve Dogum AD

H1 Blok Kat: 1 Tup Bebek Unitesi

07070 Antalya, Turkey

E-mail: mnirea@yahoo.com

References

1. Mackay AP, Berg CJ, Atrash H. Pregnancy related mortality from preeclampsia and eclampsia. *Obstet Gynecol.* 2001; 97:533-538.
2. Schreyer P, Tzadok J, Sherman DJ, et al. Fluorescein angiography in hypertensive pregnancies. *Int J Gynecol Obstet.* 1990;34:127-132.
3. Jaffe G, Schatz H. Ocular manifestations of preeclampsia. *Am J Ophthalmol.* 1987;103:309-315.
4. Apaydin KC, Akar Y, Metinsoy M. The effect of panretinal laser photocoagulation therapy on different visual field regions of the diabetic patients. *Retina-Vitreous.* 2003;11:243-249.
5. Akar Y, Apaydin KC, Ozel A. Assessment of the effect of panretinal laser photocoagulation therapy on optic nerve head topography. *Retina-Vitreous.* 2003;11: 273-79.
6. Palacz O, Sylwestrzak Z. Effectiveness of laser therapy in retinal vein thrombosis. *Klin Oczna.* 1990;92: 22-4.
7. Heijl A, Lindgren G, Olsson J. The effect of perimetric experience in normal subjects. *Arch Ophthalmol.* 1989;107:81-86.
8. Duley L. Maternal mortality associated with hypertensive disorders of pregnancy in Africa, Asia, Latin America and the Caribbean. *Br J Obstet Gynaecol.* 1992; 99:547-553.

Subdural hematoma following obstetric analgesia, causal or coincidence?

To the Editor: Epidural analgesia is an effective technique to relieve pain during labor.¹ Intracranial subdural hematoma (ISH) can present as a rare neurological complication associated with epidural analgesia following dural puncture.²⁻⁵ We describe an intracranial subdural hematoma following the epidural injection with no dural perforation in a patient undergoing vaginal delivery. A 25-year-old pregnant woman was admitted to the hospital for delivery. She was primiparous, at 39-weeks gestation, with an uneventful pregnancy course and a normal hematologic screen. There was no significant history of illness. After she provided informed consent and with her approval, an epidural needle was inserted at L3-L4. There were no complications during the administration of the epidural catheter. After the test dose of 2 mL 2% lidocaine, 5 mL 0.5 % bupivacaine plus 5 mL normal saline were administered initially and as a top-up dose in 90 minutes. Following injections the patient was hemodynamically stable based on arterial blood pressures and heart rates. During delivery, 2.5 mL of 0.5% bupivacaine and an equal volume of normal saline were injected epidurally. The patient had adequate motor power and the second stage of labor lasted 30 minutes. The first and fifth minute Apgar scores of the newborn were 8 and 10, respectively. The patient was discharged on the postpartum day. One week after the delivery, the patient was admitted to the hospital with recurrent paresthesia in her right arm (7 to 8 times/day) lasting for 10 to 15 minutes. Her history elicited pain, which was spreading from right wrist to the forearm and occasionally to the face, beginning two days after the delivery. She had temporary, intermittent lethargy and frontal bilateral headache that was postural in nature,

exacerbated with leaning forward and decreased with the horizontal position. On physical examination, she had no abnormal findings except superficial hypoalgesia in her right forearm. Laboratory investigations showed only a high erythrocyte sedimentation rate. Vertebral artery and carotid colored Doppler blood flow measures and heart functions with echocardiography were all in normal ranges. Computerized brain tomography of the skull showed sulcal effacement in the left parietal region, isodense subdural hematoma in the supratentorial region and displacement in grey substance-Luschke interphase. Subdural hematoma in the left parietal region was confirmed with magnetic resonance imaging. She had been hospitalized and medically treated for cerebral contusion six years earlier because of head trauma. She also had suffered from occasional headaches. She was medically discharged on the seventh day without neurologic complication. Contrasted computerized tomography performed 3 weeks later showed that the hematoma was completely resolved.

Epidural blocks are widely used in obstetrics for the pain relief of labor and delivery.^{1,2} After accidental dural puncture the late neurologic complications such as epidural or subdural hematoma, subdural cyst, arachnoiditis, subdural hygroma were reported.^{3,5,6} Besides dural injury and coagulation defects, preeclampsia, or large volume (=25-30ml) and rapid injection of a solution into the epidural space may lead also increases intracerebral hemorragia risk.^{5,6} In our case, etiological factors such as preeclampsia, coagulation abnormalities, and dural injury were excluded and less than 20 mL volume was administered very slowly for epidural analgesia.⁷⁻⁹ The history of head trauma may be an etiological factor for subdural hematoma. It was also suggested that the pressure changes during vaginal delivery might increase the risk of subdural hematoma.¹⁰ The

Figure 1. The first computerized brain tomography of the skull shows sulcal effacement in left parietal region.

Figure 2. Contrasted computerized brain tomography which was performed 3 weeks later shows that the haematoma was completely resolved.

alternative diagnosis was spontaneous tear of intracranial blood vessels weakened by past head injury and caused by the valsalva maneuver during the second stage of labor. A previous history of head injury should be inquired in a patient with headaches following epidural anesthesia.

In conclusion we suggest that attacks of paresthesia in the upper body following epidural anesthesia even without any dural injury should be a reminder of intracerebral subdural hematoma which may be life threatening if not properly treated.

Balcioglu Okan, MD

Anis Aribogan, MD

Sule Akin, MD

Department of Anesthesiology

Faculty of Medicine

Cukurova University

Turkey

References

1. Banhamou D. Effect of epidural analgesia on obstetrical mechanics. *Can Anaesthesiol.* 1994; 42: 261-264.
2. Glosten B. Epidural and spinal analgesia/anesthesia. In: Chestnut DH, ed. *Obstetric anesthesia principles and practice*. Second edition. St. Louis; Mosby, Inc: 1999;360-382.

3. Palot M, Visseaux H, Botmans C, Pire JC. Epidemiology of complications obstetrical epidural analgesia. *Can Anesthesiol*. 1994;42:229-233.
4. Crawford JS. Some maternal complications of epidural for labour. *Anesthesia*. 1985; 40:1219-1225.
5. Cousins MJ, Verring BT. Epidural Neural Blockade. In: Cousins MJ, Bridenbaugh PO, eds. *Neural Blockade. In Clinical Anesthesia and Management of Pain*. 1998; 287-288,306.
6. Jack TM. Postpartum intracranial subdural hematoma. A possible complication of epidural analgesia. *Anaesthesia*. 1979; 34:176-180.
7. Giannina G, Smith D, Belfort MA, Moise KJ Jr. Automatic subdural hematoma associated with pre-eclampsia. *J Matern Fetal Med*. 1997; 6: 93-95.
8. Hubert P. Acute subdural hematoma of the convexity caused by rupture of an aneurysm which is the anteriorcommunicating artery. Approach of a case in a pregnant woman. *Neurochirurgie*. 1994; 363-368.
8. Marshal S, Hanman FS. Subdural hematoma following administration of urea for hypertension. *JAMA*. 1962; 182:813-814.
9. Eerola L, Kaunkinen L, Kaunkinen S. Fatal brain lesion following spinal anesthesia, Report of a case. *Acta Anesthesiol Scand*. 1981; 25:115-116.
10. Thons M, Neveling D, Hatzmann W. Intracerebral subdural hematoma after delivery with peridural catheter anesthesia. *Z Geburtshilfe Perinatol*. 1993; 197:235-237.