Published in final edited form as:

J Adolesc Health. 2018 September ; 63(3): 335–341. doi:10.1016/j.jadohealth.2018.03.022.

Differential Risk Factors for Unhealthy Weight Control Behaviors by Sex and Weight Status Among U.S. Adolescents

Jason M. Nagata, MD, MSc¹, Andrea K. Garber, PhD, RD¹, Jennifer L. Tabler, PhD², Stuart B. Murray, PhD³, and Kirsten Bibbins-Domingo, MD, PhD, MAS^{4,5}

¹Division of Adolescent and Young Adult Medicine, University of California, San Francisco, San Francisco, CA

²Department of Sociology and Anthropology, University of Texas Rio Grande Valley, Edinburg, TX

³Department of Psychiatry, University of California, San Francisco, San Francisco, CA

⁴Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA

⁵Department of Medicine, University of California, San Francisco, San Francisco, CA

Abstract

Purpose—To determine if previously reported risk factors for the development of unhealthy weight control behaviors differ by sex and weight status using a nationally representative longitudinal sample of adolescents followed through young adulthood.

Methods—We used nationally representative longitudinal cohort data collected from baseline (11–18 years old, 1994–1995, Wave I) and seven-year follow-up (18–24 years old, 2001–2002, Wave III) of the National Longitudinal Study of Adolescent to Adult Health (Add Health). We examined adverse childhood events and adolescent family, school, body image, and mental health factors for development of unhealthy weight control behaviors including vomiting, fasting/skipping meals, or laxative/diuretic use to lose weight at seven-year follow-up in young adulthood.

Results—Of the 14,322 included subjects, 11% reported unhealthy weight control behavior at follow-up in young adulthood, with the highest proportion (23.7%) among overweight/obese females and the lowest proportion (3.7%) among underweight/normal weight males (12.9%). All adolescent family factors were significantly associated with unhealthy weight control behaviors in underweight/normal weight females whereas none were significantly associated in overweight/obese males. Similar trends were noted for adverse childhood events and adolescent school and

Corresponding Author: Jason M. Nagata, 3333 California Street, Suite 245, San Francisco, California 94118, Telephone: +1 (626) 551-1932, jasonmnagata@gmail.com.

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflict of Interest: The authors have no conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

community factors. Adolescent self-perception of being overweight was associated with young adult unhealthy weight control behaviors among all subgroups.

Conclusions—Risk factors for unhealthy weight control behaviors may differ based on sex and weight status. Screening, prevention, and treatment interventions for unhealthy weight control behaviors in adolescents and young adults may need to be tailored based on sex and weight status.

Keywords

Obesity; overweight; eating behavior; eating disorders; disordered eating behaviors; adolescents; young adults; family functioning; body image

Implications and Contribution

With a nationally representative sample, this study shows that family and school factors are associated with unhealthy weight control behaviors in under/normal weight females but not in overweight/obese adolescents or in males. Prevention, screening, and treatment interventions for unhealthy weight control behaviors may need to be tailored based on sex and weight status.

Introduction

Unhealthy weight control behaviors including vomiting, fasting/skipping meals, or laxative/diuretic use to lose weight are common among adolescents and young adults [1] and are associated with increased risk for eating disorders [2, 3], alcohol and tobacco use [4], mental health problems [5], and poor nutritional intake and quality [5]. Given these potentially serious medical and psychosocial consequences, unhealthy weight control behaviors represent significant public health challenges. Although traditionally thought to be limited to underweight females, unhealthy weight control behaviors are increasingly recognized among those who are overweight or obese [6] and among males [7, 8].

Previous studies have identified various socioenvironmental risk factors for the development of unhealthy weight control behaviors such as family dysfunction and disconnectedness [6, 9], school disconnectedness [8, 10], and adverse childhood events [11]. Prior regional samples have reported that higher family functioning and parent connection were associated with lower odds of engaging in unhealthy weight control behaviors [6, 9, 12]. Furthermore, childhood physical [11] and sexual abuse [13] has been shown to be associated with the development of unhealthy weight control behaviors. Although some of these studies did examine risk by sex, they did not also disaggregate by weight status. The presentations of unhealthy weight control behaviors may differ by both sex [7] and by weight status, particularly for those who are overweight or obese [14]; however, little is known about how risk for unhealthy weight control behaviors may differ when disaggregated by both sex and weight status.

Therefore, the objective of this study was to determine differences in risk during adolescence by sex and weight status for the development of unhealthy weight control behaviors in young adulthood using a nationally representative longitudinal sample (Figure 1).

Methods

Study design and sample

The National Longitudinal Study of Adolescent to Adult Health (Add Health) has followed a nationally representative cohort of youth in the US from adolescence through adulthood [15]. The baseline sample used systematic sampling methods and implicit stratification to ensure that the high schools (n=80) and middle schools (n=52) selected were representative of US schools with respect to region of country, urbanicity, size, type, and ethnicity. For this particular study, we used the restricted-use baseline sample (Wave I), which was collected from 1994–1995 when subjects were 11–18 years old and seven-year follow-up data (Wave III), which was collected in 2001–2002 when subjects were 18–24 years old. We included subjects in the nationally representative weighted sample who had data at both baseline and seven-year follow-up (N=14,322). Further details about the Add Health study design, coordinated by the Carolina Population Center, can be found elsewhere [15]. The University of North Carolina Institutional Review Board approved all Add Health study procedures, and the University of California, San Francisco Institutional Review Board deemed this specific project exempt.

Procedures

At baseline and seven-year follow-up, an interviewer traveled to the home or another suitable location for the potential participant. Written consent was obtained from the parent if the participant was under age 18, or from the participant if 18 or older. Interviews lasted approximately 90 minutes and were conducted in as private an area as possible. Audio computer-assisted self-interview (baseline) and computer-assisted self-interview (follow-up) were used by participants to answer potentially sensitive questions.

Measures

Baseline measures—Demographic characteristics; socioenvironmental variables including family, school, and community factors; adverse childhood events; body image and weight factors; and mental health questions were collected during an in-home interview. Family factors included questions about family functioning and family connectedness. School and community factors included questions about how much adolescents felt cared for by friends, teachers, and other adults. Adverse childhood events included reports of childhood physical or sexual abuse or neglect. Body image and weight factors included weight perception (“how do you think of yourself in terms of weight?”) and if adolescents were currently trying to lose weight. Depression score was a modified version of the Center for Disease Epidemiology Depression Scale (CESD-20) [16]. A full list of measures is listed in online Appendix A.

Self-reported weight (pounds) and height (inches) were converted to kilograms and meters to calculate body mass index (BMI) using the standard formula weight (kilograms) divided by height (meters) squared ($BMI = \text{weight}/\text{height}^2$). BMI was then converted into sex- and age-specific percentiles and then classified as underweight (BMI <5th percentile), normal weight (5th percentile to less than the 85th percentile), overweight (85th to less than the 95th

percentile), or obese (95th percentile or greater) in accordance with guidelines from the US Centers for Disease Control and Prevention (CDC) [17].

Seven-year follow-up measures—Unhealthy weight control behaviors including vomiting, fasting/skipping meals, or laxative/diuretic use to lose weight in the past seven days were self-reported at seven-year follow-up during an in-home interview as the primary outcome variable [18].

Statistical analysis

Data analysis was performed in 2017 using STATA 15.0. We used Add Health's pre-constructed sample weights to provide a nationally representative sample [19, 20]. We tested unadjusted differences between adolescent risk factors and unhealthy weight control behaviors in young adulthood using Chi-square tests, disaggregated by sex and weight status. We used multiple logistic regression to identify associations between unhealthy weight control behaviors and family, school, community, adverse childhood events, and body image factors, adjusting for age, race/ethnicity, and household income [6, 9, 21]. These analyses are disaggregated by sex and weight status to identify potential differences in associations based on sex or weight category. Alpha was set at 95% confidence, and $p < 0.05$ is considered significant.

Results

Of the 18,924 adolescents in the nationally representative weighted baseline sample, 14,322 (75.7%) had seven-year follow-up data. Baseline demographic and anthropometric characteristics of the sample are reported in Table 1, disaggregated by sex and weight status. Overall, 11% reported unhealthy weight control behavior at seven-year follow-up, with the highest proportion (23.7%) among overweight or obese females and the lowest proportion (3.7%) among underweight or normal weight males (12.9%).

In unadjusted comparisons, underweight or normal weight females who reported unhealthy weight control behaviors reported lower satisfaction with family factors compared to those who did not report unhealthy weight control behaviors (Table 2). For instance, underweight or normal weight females with versus without unhealthy weight control behaviors reported lower satisfaction with their relationship with their mother (81.7% vs 86.2%, $p = 0.013$) and father (72.1% vs 81.0%, $p = 0.001$), respectively. All family factors were significantly associated with unhealthy weight control behaviors in underweight or normal weight females whereas none were significantly associated in males across all weight strata. Similar trends were noted for school and community factors (Table 2). Body image and weight concerns were associated with unhealthy weight control behaviors among all subgroups. Depressive symptoms were associated with increased odds of unhealthy weight control behaviors in females but not males.

Table 3 reports odds (adjusted odds ratios) of unhealthy weight control behaviors based on family, school and community, adverse childhood, body image, and mental health factors, while adjusting for age, race/ethnicity, and household income. Multiple logistic regression models are disaggregated by sex and weight status. Similar to unadjusted models, family,

school, and community factors are most consistently associated with unhealthy weight control behaviors in the underweight or normal weight female subgroup, but not other subgroups. For instance, female underweight or normal weight subjects reporting that their mother cared very much compared to not caring very much have 36% lower odds of development of unhealthy weight control behaviors at follow-up. Perception of being slightly or very overweight is associated with higher odds of unhealthy weight control behavior at follow-up in all subgroups ($p<0.05$).

Discussion

In this large, nationally representative longitudinal sample, we find that family, school, and community factors are associated with unhealthy weight control behaviors only in underweight or normal weight females. These same factors are not consistently associated with unhealthy weight control behaviors among males or overweight/obese adolescents. This finding is important since prevention, screening, and treatment interventions for unhealthy weight control behaviors may need to be tailored based on sex and weight status.

Previous studies have similarly found that a positive family environment is associated with lower odds of engaging in unhealthy weight control behaviors, but these studies did not disaggregate results by baseline weight status and sex [6, 9, 12]. Two of these studies did, however, find shared risk and protective factors (including body dissatisfaction and weight concerns) for unhealthy weight control behaviors and overweight/obesity as separate outcomes.[6, 9] This research contributes to the literature by disaggregating the odds by baseline sex and weight status, finding that these associations were only present in underweight or normal weight females.

Our findings that adverse childhood events are associated with increased odds of unhealthy weight control behaviors is consistent with previous literature [11, 13], and disaggregation by sex and weight status provides additional insights. Childhood physical abuse has previously been shown to be associated with unhealthy weight control behaviors in women [11]. When disaggregating by weight and sex we find that childhood physical abuse is associated with unhealthy weight control behaviors in underweight or normal weight females and overweight or obese males in adjusted models. Prior studies that did not disaggregate by sex and weight status did not find a significant association between childhood neglect and unhealthy weight control behaviors [11]; however, when we disaggregated these groups we found that childhood neglect was associated with increased odds of unhealthy weight control behaviors in females but not in males across both weight categories.

Our study adds to this previous literature by showing that previously reported family-related associations with unhealthy weight control behaviors may be most applicable to the subset of underweight or normal weight females. In our sample and in others, overweight or obese adolescents reported higher levels of unhealthy weight control behaviors than underweight or normal weight adolescents [6, 14]. It is possible that the risk factors and etiology for unhealthy weight control behaviors among overweight or obese adolescents and in males may be different than for underweight or normal weight females.

Unhealthy weight control behaviors and eating disorders in males are typically under-recognized and therefore represent an important area of research [7, 8]. Crucially, the present results underscore the notion that results documenting the risk and maintenance of disordered eating practices in largely female populations cannot be seamlessly extrapolated to male populations, since males with unhealthy weight control behaviors may present fundamentally differently than female counterparts [7, 8, 22–25]. In this study, depressive symptoms and childhood neglect were associated with unhealthy weight control behaviors in females but not males. The only family factor protective of unhealthy weight control behavior in underweight or normal weight males was the perception that one's father cared very much. Given that most family and school factors were not significantly associated with unhealthy weight control behaviors in males across weight categories, there may be different risk factors for unhealthy weight control behaviors in this subgroup. However, it is important to note that the questions designed to assess disordered eating practices (i.e., laxative use, purging) may have been more salient for females, given that males may engage in unhealthy weight control behaviors connected to muscularity-oriented concerns, rather than thinness oriented concerns [7, 22]. Males with eating disorders may present with a greater array of psychiatric comorbidities such as substance use and psychotic symptoms than females [7, 26]. Body image concerns, including weight perception, were significantly associated with an increased odds ratio for unhealthy weight control behaviors in males. Future research may examine other specific risk factors for unhealthy weight control behaviors in underweight or normal weight males, such as certain sports with weight cutoffs (like wrestling or rowing) [22, 27, 28], sexual orientation [29, 30], substance use, and other psychiatric comorbidity [7, 26].

Limitations of this study include the use of self-reported data at baseline (including height and weight) and follow-up, a method that may be subject to reporting bias. However, self-reported height and weight have been shown to be highly correlated with objectively measured height and weight ($r=0.99$; $p<0.001$) in community samples [31]. In addition, a seven-day timeframe was used for the recall of unhealthy weight control behaviors in Add Health, which is shorter than other measures of unhealthy weight control behaviors such as the Eating Disorder Examination Questionnaire which uses a timeframe of the past 28 days [32]. This may have underestimated the rate of unhealthy weight control behaviors compared to other measures. Strengths include the use of nationally-representative longitudinal data with seven-year follow-up in a large community sample size of adolescents followed into young adulthood.

Conclusion

Risk factors for unhealthy weight control behaviors differ based on sex and weight status, with family, school, and community factors most consistently associated with unhealthy weight control behaviors in underweight or normal weight females. Although some adverse childhood events and body image concerns were associated with unhealthy weight control behaviors in underweight or normal weight males and overweight or obese females, future research may better delineate risk factors for unhealthy weight control behaviors in overweight or obese adolescents and in males. Screening, prevention, and treatment interventions for unhealthy weight control behaviors in adolescents and young adults may

need to be tailored based on sex and weight status to address these major public health challenges.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Thanks to Nicole Capdarest-Arest for help with literature searches.

Funding: J.M.N. is a fellow in the Pediatric Scientist Development Program (K12HD00085033), funded by the American Academy of Pediatrics and the American Pediatric Society, and was supported by the Norman Schlossberger Research Fund from the University of California, San Francisco. A.K.G. was supported by NIH 5R01HD082166-02. S.B.M was supported by K23 MH115184.

References

1. Ackard DM, Neumark-Sztainer D, Story M, et al. Overeating among adolescents: prevalence and associations with weight-related characteristics and psychological health. *Pediatrics*. 2003; 111:67–74. [PubMed: 12509556]
2. Striegel-Moore RH, Bulik CM. Risk factors for eating disorders. *Am Psychol*. 2007; 62:181–198. [PubMed: 17469897]
3. McKnight Investigators. Risk factors for the onset of eating disorders in adolescent girls: results of the McKnight longitudinal risk factor study. *Am J Psychiatry*. 2003; 160:248–254. [PubMed: 12562570]
4. French SA, Story M, Downes B, et al. Frequent dieting among adolescents: psychosocial and health behavior correlates. *Am J Public Health*. 1995; 85:695–701. [PubMed: 7733431]
5. Neumark-Sztainer D, Wall M, Larson NI, et al. Dieting and disordered eating behaviors from adolescence to young adulthood: findings from a 10-year longitudinal study. *J Am Diet Assoc*. 2011; 111:1004–1011. [PubMed: 21703378]
6. Neumark-Sztainer DR, Wall MM, Haines JI, et al. Shared risk and protective factors for overweight and disordered eating in adolescents. *Am J Prev Med*. 2007; 33:359–369. [PubMed: 17950400]
7. Murray SB, Nagata JM, Griffiths S, et al. The enigma of male eating disorders: A critical review and synthesis. *Clin Psychol Rev*. 2017; 57:1–11. [PubMed: 28800416]
8. Lampard AM, Maclehose RF, Eisenberg ME, et al. Adolescents who engage exclusively in healthy weight control behaviors: Who are they? *Int J Behav Nutr Phys Act*. 2016; 13 5-016-0328-3.
9. Haines J, Kleinman KP, Rifas-Shiman SL, et al. Examination of shared risk and protective factors for overweight and disordered eating among adolescents. *Arch Pediatr Adolesc Med*. 2010; 164:336–343. [PubMed: 20368486]
10. Jones DC, Thorbjorg HV, Lee Y. Body Image and the Appearance Culture Among Adolescent Girls and Boys: An Examination of Friend Conversations, Peer Criticism, Appearance Magazines, and the Internalization of Appearance Ideals. *J Adolesc Res*. 2004; 19:323–339.
11. Fuemmeler BF, Dedert E, McClernon FJ, et al. Adverse childhood events are associated with obesity and disordered eating: results from a U.S. population-based survey of young adults. *J Trauma Stress*. 2009; 22:329–333. [PubMed: 19588510]
12. Berge JM, Wall M, Larson N, et al. The unique and additive associations of family functioning and parenting practices with disordered eating behaviors in diverse adolescents. *J Behav Med*. 2014; 37:205–217. [PubMed: 23196919]
13. Gustafson TB, Sarwer DB. Childhood sexual abuse and obesity. *Obes Rev*. 2004; 5:129–135. [PubMed: 15245381]
14. Loth K, Wall M, Larson N, et al. Disordered eating and psychological well-being in overweight and nonoverweight adolescents: secular trends from 1999 to 2010. *Int J Eat Disord*. 2015; 48:323–327. [PubMed: 25641764]

15. Harris KM, Halpern CT, Witsel E, et al. The National Longitudinal Study of Adolescent to Adult Health: Research Design. Jul.2017
16. Radloff LS. The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement*. 1977; 1:385–401.
17. Centers for Disease Control (CDC). Defining Adult Overweight and Obesity. Apr 27.2012
18. Stephen EM, Rose JS, Kenney L, et al. Prevalence and correlates of unhealthy weight control behaviors: findings from the national longitudinal study of adolescent health. *J Eat Disord*. 2014; 216-2974-2-16. eCollection 2014.
19. Harris KM. The Add Health Study: Design and Accomplishments. Carolina Population Center, University of North Carolina; Chapel Hill: 2013.
20. Chen P. Appropriate Analysis in Add Health: Correcting for Design Effects & Selecting Weights. Carolina Population Center, University of North Carolina; Chapel Hill: 2014.
21. Tabler J, Utz RL. The influence of adolescent eating disorders or disordered eating behaviors on socioeconomic achievement in early adulthood. *Int J Eat Disord*. 2015; 48:622–632. [PubMed: 25808740]
22. Murray SB, Accurso EC, Griffiths S, et al. Boys, Biceps, and Bradycardia: The Hidden Dangers of Muscularity-Oriented Disordered Eating. *J Adolesc Health*. 2018; 62:352–355. [PubMed: 29241987]
23. Nagata JM, Golden NH, Peebles R, et al. Assessment of Sex Differences in Body Composition Among Adolescents With Anorexia Nervosa. *J Adolesc Health*. 2017; 60:455–459. [PubMed: 28087266]
24. Nagata JM, Golden NH, Peebles R, et al. Assessment of sex differences in bone deficits among adolescents with anorexia nervosa. *International Journal of Eating Disorders*. 2017; 50:352–58. [PubMed: 27611361]
25. Nagata JM, Golden NH, Leonard MB, et al. Assessment of Sex Differences in Fracture Risk Among Patients With Anorexia Nervosa: A Population-Based Cohort Study Using The Health Improvement Network. *J Bone Miner Res*. 2017; 32:1082–1089. [PubMed: 28019700]
26. Striegel-Moore RH, Garvin V, Dohm FA, et al. Psychiatric comorbidity of eating disorders in men: a national study of hospitalized veterans. *Int J Eat Disord*. 1999; 25:399–404. [PubMed: 10202650]
27. Kiningham RB, Gorenflo DW. Weight loss methods of high school wrestlers. *Med Sci Sports Exerc*. 2001; 33:810–813. [PubMed: 11323553]
28. Nagata JM, Carlson JL, Kao JM, et al. Characterization and correlates of exercise among adolescents with anorexia nervosa and bulimia nervosa. *Int J Eat Disord*. 2017; 50:1394–1403. [PubMed: 29112280]
29. Hadland SE, Austin SB, Goodenow CS, et al. Weight misperception and unhealthy weight control behaviors among sexual minorities in the general adolescent population. *J Adolesc Health*. 2014; 54:296–303. [PubMed: 24182939]
30. Watson RJ, Adjei J, Saewyc E, et al. Trends and disparities in disordered eating among heterosexual and sexual minority adolescents. *Int J Eat Disord*. 2017; 50:22–31. [PubMed: 27425253]
31. Pursey K, Burrows TL, Stanwell P, et al. How accurate is web-based self-reported height, weight, and body mass index in young adults? *J Med Internet Res*. 2014; 16:e4. [PubMed: 24398335]
32. Fairburn CG, Beglin S. Eating disorder examination questionnaire. In: Fairburn CG, editor *Cognitive Behavior Therapy and Eating Disorders*. New York: Guilford Press; 2008. 309–313.

Figure 1.
Conceptual framework for how adverse childhood events and adolescent socio-environmental and psychological factors may predict young adult unhealthy weight control behaviors differently by sex and weight status.

Baseline demographic and anthropometric characteristics and outcome at seven-year follow up, by sex and baseline weight status

Table 1

	n ^a	Total	Underweight or normal weight	Overweight or obese	
			Female	Male	
N					
	14,322	14,322	5,598	4,798	1,961
Total		Mean ± SE / % ^b	Mean ± SE / % ^b	Mean ± SE / % ^b	Mean ± SE / % ^b
Demographic characteristics					
Age	14,314	15.94 ± 0.12	15.89 ± 0.12	16.08 ± 0.12	15.88 ± 0.14
Race/ethnicity	14,312				
White (non-Hispanic)		67.9%	72.2%	68.6%	64.6%
Black/African American (non-Hispanic)		15.5%	12.8%	14.8%	16.0%
Hispanic/Latino		11.8%	10.3%	11.5%	13.9%
Asian/Pacific Islander (non-Hispanic)		3.4%	3.8%	4.0%	3.1%
American Indian/Native American		0.5%	0.4%	0.3%	1.5%
Other		0.8%	0.7%	0.9%	1.0%
Household income, (thousands of dollars, parent report)	10,830	45.74 ± 1.68	49.23 ± 2.18	47.50 ± 2.05	38.92 ± 1.23
Baseline anthropometric characteristics					
Body mass index (BMI), kg/m ²	13,942	22.47 ± 0.11	20.35 ± 0.06	20.53 ± 0.09	28.21 ± 0.19
Outcome variable					
Unhealthy weight control behavior	14,322	11.0%	12.9%	3.7%	23.7%

^aGroup n may not sum to total N because of missing data^bAll means and percentages are calculated with weighted data to reflect the representative proportion in the target U.S. population

Unadjusted adolescent risk factors for unhealthy weight control behaviors (UWCB) in young adulthood, by sex and weight status

Table 2

	Unhealthy weight control behaviors						Overweight or obese at baseline					
	Underweight or normal weight at baseline											
	Female			Male			Female			Male		
	No UWCB	UWCB	p ^a	No UWCB	UWCB	p ^a	No UWCB	UWCB	p ^a	No UWCB	UWCB	p ^a
Family factors												
Mother cares (very much)	89.5 %	86.0 %	0.029	91.2 %	91.5 %	0.945	87.0 %	84.0 %	0.222	89.7 %	89.1 %	0.853
Mother communication (satisfied)	76.5 %	69.7 %	0.005	83.1 %	83.5 %	0.925	77.2 %	73.7 %	0.264	86.4 %	87.0 %	0.877
Satisfied with relationship with mother (satisfied)	86.2 %	81.7 %	0.013	91.7 %	90.9 %	0.917	84.4 %	82.4 %	0.450	93.4 %	92.6 %	0.749
Father cares (very much)	83.7 %	78.2 %	0.020	83.3 %	77.7 %	0.198	80.9 %	83.4 %	0.413	81.6 %	85.5 %	0.456
Father communication (satisfied)	73.8 %	67.0 %	0.022	79.5 %	76.5 %	0.550	72.3 %	66.8 %	0.183	80.8 %	78.2 %	0.573
Satisfied with relationship with father (satisfied)	81.0 %	72.1 %	0.001	86.1 %	81.1 %	0.240	77.3 %	72.1 %	0.111	86.8 %	84.2 %	0.511
Parents care (very much)	86.9 %	80.6 %	<0.001	84.6 %	85.2 %	0.886	83.0 %	81.8 %	0.654	86.4 %	84.5 %	0.508
Family understands (quite a bit / very much)	55.0 %	45.5 %	<0.001	56.9 %	56.5 %	0.915	49.8 %	48.5 %	0.710	61.6 %	59.0 %	0.608
Family pays attention (quite a bit / very much)	71.9 %	57.5 %	<0.001	72.9 %	68.9 %	0.380	66.2 %	59.2 %	0.023	73.3 %	66.9 %	0.142
Family has fun (quite a bit / very much)	62.7 %	51.6 %	<0.001	61.5 %	63.3 %	0.689	61.4 %	57.5 %	0.246	67.0 %	66.2 %	0.854
Want to leave home (very much)	5.7 %	9.3 %	0.003	6.0 %	8.2 %	0.344	9.1 %	9.7 %	0.794	4.5 %	5.6 %	0.592
School and community factors												
Adults care (very much)	61.5 %	53.4 %	0.002	51.0 %	44.9 %	0.236	58.4 %	55.1 %	0.353	52.3 %	59.7 %	0.146
Teachers care (very much)	19.4 %	13.4 %	0.004	17.1 %	16.8 %	0.928	19.2 %	16.7 %	0.368	17.9 %	19.5 %	0.689
Friends care (very much)	54.8 %	48.8 %	0.049	36.8 %	28.5 %	0.052	49.2 %	45.3 %	0.240	34.0 %	35.4 %	0.730
Happy at school (agree / strongly agree)	66.9 %	59.4 %	0.003	67.4 %	64.2 %	0.513	63.0 %	60.8 %	0.532	69.2 %	68.7 %	0.896
Adverse childhood events												
Any physical abuse	24.4 %	36.3 %	<0.001	29.0 %	41.4 %	0.017	29.0 %	32.6 %	0.279	29.8 %	37.3 %	0.133
Any sexual abuse	4.4 %	4.7 %	0.799	3.9 %	6.5 %	0.201	4.4 %	6.9 %	0.110	5.7 %	5.2 %	0.842
Any neglect	35.8 %	45.2 %	<0.001	43.8 %	51.0 %	0.213	37.7 %	48.1 %	0.006	41.3 %	41.5 %	0.967
Basic needs (food, clothing, keeping clean) not met	7.9 %	9.4 %	0.341	13.7 %	18.5 %	0.149	9.2 %	15.0 %	0.004	15.6 %	19.7 %	0.265
Body image and weight concerns												

	Unhealthy weight control behaviors							
	Underweight or normal weight at baseline				Overweight or obese at baseline			
	Female		Male		Female		Male	
	No UWCB	UWCB	p ^a	No UWCB	UWCB	p ^a	No UWCB	UWCB
Weight perception (slightly / very overweight)	24.8 %	40.1 %	<0.001	7.5 %	19.8 %	<0.001	60.8 %	70.7 %
Trying to lose weight	34.3 %	57.3 %	<0.001	8.3 %	25.2 %	<0.001	48.7 %	58.9 %
Mental health								
Depressive symptoms	26.3 %	38.2 %	<0.001	19.1 %	24.8 %	0.178	19.8 %	21.5 %
								0.686

Bold indicates p-value <0.05

^aPearson's chi square test

Author Manuscript

Author Manuscript

	Unhealthy weight control behaviors												
	Underweight or normal weight at baseline												
	Female				Male							Male	
	OR				OR							OR	p
Trying to lose weight	2.47				<0.001							0.012	0.133
Mental health													
Depressive symptoms	1.65				<0.001							0.006	0.482

Bold indicates p-value <0.05

^aOR = odds ratio, adjusted for age, race/ethnicity, and household income