Published in final edited form as:

Psychiatry Res. 2017 October ; 256: 444–447. doi:10.1016/j.psychres.2017.07.019.

Blood cadmium and depressive symptoms: Confounded by cigarette smoking

Danielle E. Kostrubiak^{a,*}, Caterina Vacchi-Suzzi^b, Dylan M. Smith^b, and Jaymie R. Meliker^{b,c}

^aStony Brook University School of Medicine, Stony Brook, NY, USA

^bProgram in Public Health, Department of Family, Population and Preventive Medicine, Stony Brook University, Stony Brook, NY, USA

^cConsortium for Inter-Disciplinary Environmental Research, USA

Abstract

Our aim was to explore the association between blood cadmium (BCd) and depressive symptoms, adjusting for pack years and blood cotinine, and also stratifying by smoking status. Using data from the US National Health and Nutrition Examination Survey (NHANES) 2005–2012, we categorized depressive symptoms using the PHQ-9 (Patient Health Questionnaire-9) survey and modeled depressive symptoms in relation to BCd adjusted for blood cotinine, pack years of smoking, and other covariates. We also stratified by self-reported smoking status (current, former, never). There were 11,209 subjects from 2005 to 2012, age ≥ 18 with PHQ-9, smoking, and blood cadmium data available. 876 (7.8%) met criteria for depressive symptoms. Depressive symptoms were associated with BCd levels in a crude model and with adjustment for pack years and cotinine. The association disappeared when analyzed among current, former, or never smokers. Consistent with the literature, we found an association between BCd and depressive symptoms; however, that association disappears in analyses stratified by smoking status. This suggests residual confounding may be present. It is important to stratify by smoking status when investigating health outcomes associated with BCd.

Keywords

Tobacco; Depression; Heavy metals

1. Introduction

Cadmium, like other heavy metals, is a known neurotoxicant, likely through mechanisms of oxidative stress, alterations in neurotransmitter release, damage to the blood brain barrier, and induction of neuron apoptosis (Mendez-Armenta and Rios, 2007). It has been associated with various psychiatric disorders including schizophrenia, bipolar disorder, and major depressive disorder (Mendez-Armenta and Rios, 2007; Olabanji, 2011; Orisakwe, 2014). In

*Correspondence to: Stony Brook University School of Medicine, MS-IV, Health Sciences Center, Stony Brook University, Level 3, Room 071, Stony Brook, NY, USA., Danielle.kostrubiak@stonybrookmedicine.edu (D.E. Kostrubiak).

Conflict of interest
None.

particular, previous studies have shown a positive association between elevated blood cadmium (BCd) levels and depressive symptoms (Berk et al., 2014; Han et al., 2016; Scinicariello and Buser, 2015). The link between urine cadmium (UCd) levels and depressive symptoms is less clear (Shiue, 2015)-possibly due to the fact that BCd is more indicative of short term mixed with long-term exposure and UCd is more representative of lifetime exposure (Jarup et al., 1998).

Cigarette smoking is one of the major sources of cadmium exposure in the US. Tobacco plants absorb cadmium from the soil, and persons who smoke 20 cigarettes per day absorb about 1 µg of cadmium through their lungs from smoking each day (Jarup and Akesson, 2009). In current, never and former smokers, a dose-dependent relationship has been observed in BCd- with highest blood levels in current heavy smokers (Ellingsen et al., 1997). Furthermore, smoking has a known association with depression and other mental health disorders, and smoking cessation is associated with a decreased risk of depression (Bakhshaie et al., 2015). Therefore, it is important to adequately control for cigarette smoking when investigating the association between cadmium and depressive symptoms.

We hypothesize that BCd is associated with depressive symptoms in unadjusted models, and that residual confounding may still be present after controlling for serum cotinine and pack years. Stratifying by cigarette smoking (current, former, never) may be necessary to remove the confounding influence of cigarette smoking.

2. Methods

Data from the US National Health and Nutrition Examination Survey (NHANES) (NCHS, 2016b) was used to examine the cross-sectional relationships between blood cadmium and depressive symptoms. Depressive symptoms were characterized using the PHQ-9, which was administered as part of NHANES starting in 2005 as a screening test. This is a validated Patient Health Questionnaire that has 9 questions pertaining to defining features of depression from the diagnostic statistic manual (DSM) (Kroenke, 2002). The questionnaire is scored to identify patients with depressive symptoms; scores greater than or equal to 10 were accepted as positive results for depressive symptoms (Kroenke, 2002; Milette et al., 2010).

Blood cadmium levels were assessed using whole blood specimen collection, which is discussed in further detail in the NHANES Laboratory/Medical Technologists Procedures Manual (LPM) (NCHS, 2016b). The samples are sent to the National Center for Environmental Health, where inductively coupled plasma mass spectrometry is used to determine cadmium concentrations, measured in units of micrograms per liter (NCHS, 2016b). The limit of detection for 2005–2010 was 0.14 µg/L, and 0.11 µg/L for 2011–2012 (NCHS, 2016b). Serum cotinine is also measured via whole blood specimens, following the same protocol as delineated in the LPM. Isotope dilution-high performance liquid chromatography/ atmospheric pressure chemical ionization tandem mass spectrometry is used to determine serum cotinine levels (NCHS, 2016b). The limit of detection for 2005–2012 was 0.11 ng/ml (NCHS, 2016b).

Relevant covariate information, including demographics, smoking and drinking habits, and health history was collected via home interview and available in NHANES. Poverty income ratio (PIR) is a ratio of self-reported income to the local poverty level (NCHS, 2016b). Smoking was assessed from self-report including the current use of cigarettes, past use of cigarettes, cigarettes per day, and years of smoking. Pack-years were calculated from reported cigarettes per day divided by 20 cigarettes per pack, multiplied by years of smoking. Self-reported average number of alcoholic drinks/day in the past 12 months was also downloaded from NHANES.

SAS was used for statistical analysis, by extracting data directly from the NHANES website. Chi-square testing was used to assess for categorical demographic differences (sex, race, smoking status) and t-testing was used to assess for continuous variable differences (PIR, age, blood Cd, pack-years, blood cotinine) between depressed and non-depressed subjects. Subjects ≥ 18 were included in the data analysis, and the NHANES weighting system was used to account for sampling and survey design (NCHS, 2016a). Using data from 2005 to 2012, PHQ-9 ≥ 10 was defined as positive for depressive symptoms, and modeled in a crude logistic regression for its association with blood cadmium, adjusted for age, race, alcohol use, and poverty income ratio (PIR), blood cotinine, and pack years smoking. These variables were chosen as they could potentially act as confounders due to associations with both BCd and depressive symptoms. We also modeled the logistic regression without statistical adjustments and present both sets of results. We then stratified by self-reported smoking status (current, former, never) to tease out the role of cadmium independent of variation in smoking. As a sensitivity analysis we repeated all analyses without the NHANES sampling weights and results were similar. We also modeled categorical BCd defined by quartiles, a quadratic regression, and a log transformed logistic regression, and results were similar. We ran t-tests for BCd levels between depressed and non-depressed subjects, stratified by smoking status, to aid in interpreting the results.

3. Results

The cohort included 11,209 subjects from 2005 to 2012 with PHQ-9, smoking, and blood cadmium data available, age greater than or equal to 18. Of these subjects, 876 (7.8%) met criteria for depressive symptoms as defined by PHQ-9 ≥ 10 . Patients in the depressive symptom group were predominately female ($p < 0.01$), and race distribution showed a greater percentage of Black and non-Mexican Hispanic subjects in the depressed population (Table 1). The mean age was significantly lower in the depressed group but only by two years ($p = 0.01$) and mean PIR was lower in the depressed group, ($p < 0.01$). Mean blood cadmium was higher in the depressed group ($p < 0.01$). The depressed group was also more likely to be current smokers. Additionally, both mean pack-years and blood cotinine were higher in the depressed group ($p < 0.01$).

In an unadjusted logistic regression model, depressive symptoms were associated with BCd levels (OR = 1.53, 95% CI: 1.37–1.71). In a model with adjustment for age, sex, race, PIR and alcohol use, depressive symptoms were still associated with BCd levels, although less strongly (OR = 1.37 95% CI: 1.24–1.51). This association was mitigated by further adjustment for pack-years and serum cotinine level but still significant (OR = 1.16 95% CI:

1.04–1.30; interpreted as a 16% increase in risk for depressive symptoms associated with each 1 µg/L increase in BCd) (Table 2). To rule out the possibility of residual confounding by smoking, we stratified by smoking status; however, this resulted in no association between BCd and depressive symptoms in current, former, or never smokers (Table 2), suggesting that residual confounding was present. T-tests of BCd levels between depressed and non-depressed subjects stratified by smoking status (Table 3) show a significant difference in the group as a whole and current smokers, but not in former or never smokers.

4. Discussion

Consistent with the literature, we found a positive association between BCd and depressive symptoms in this cohort of NHANES subjects, age 18, from 2005 to 2012. Mean blood cadmium was higher in the 876 subjects who met the criteria for depressive symptoms ($p < 0.01$), and they were more likely to be current smokers, with higher blood cotinine and mean pack-year levels ($p < 0.01$) (Table 1). Since smoking is strongly associated with cadmium (Ellingsen et al., 1997), smoking may confound the association. The association between BCd and depressive symptoms was mitigated by adjusting for pack years and serum cotinine, and disappeared when stratifying by current, former and never smokers, suggesting that residual confounding was present prior to stratification (Table 2). This is supported by the raw BCd data which show a significant difference in BCd between depressed subjects and non-depressed subjects in the group as a whole and in current smokers, but not in former or never smokers (Table 3). The significant difference seen in current smokers in the raw data (Table 3), but not seen in the adjusted regression model (Table 2), may be due to no controls for pack years or blood cotinine levels in the raw analysis (Table 3).

Although recent studies that have shown a link between BCd and depressive symptoms did adjust for smoking (Berk et al., 2014; Han et al., 2016; Scinicariello and Buser, 2015), their models may still have been susceptible to residual confounding. The Berk et al. (2014) study using NHANES 2005–2010- the same population as our study - found an association between BCd and depressive symptoms, defined like our study as PHQ-9 score ≥ 10 , for the upper quartile of BCd (OR 1.48 95% CI 1.16–1.90). This association remained after adjustment for serum cotinine levels; however, they did not adjust for pack-years nor did they stratify by smoking status. Similarly, a study of young adults aged 20–39 from NHANES 2007–2010 showed an association for the upper quartile of BCd (OR 2.79, 95% CI 1.84–4.25) (Scinicariello and Buser, 2015). They did not adjust for pack-years, and while they stratified by current vs. non-current smokers, they did not differentiate between former and never smokers. With their stratification, non-current smokers had a non-significant trend ($p = 0.28$), although the upper quartile of BCd still was significantly associated with depressive symptoms (OR = 2.91 95% CI: 1.12–7.58) (Scinicariello and Buser, 2015). Scinicariello and Buser (2015) included serum cotinine to verify or correct smoking status, but did not also include it in the model. Since serum cotinine is also indicative of second-hand smoke exposure (Bramer and Kallungal, 2003), we felt it useful to include it in our model. In support of this decision, mean blood cotinine level in non-smokers is 10.93 ng/ml in our study, suggesting that it is present in nonsmokers and is important to include in the model. Another recent study of 395 elderly (age > 60) people living in Seoul, South Korea, also found a positive association between BCd and depressive symptoms, and while

they did attempt to stratify by smoking status, they had very few current (23 people) and former (24) smokers in their sample (Han et al., 2016). Taken as a whole, these studies generally controlled for urine or serum cotinine, a sensitive and specific biomarker of current tobacco exposure (Bramer and Kallungal, 2003) but they did not control for pack-years, an indicator of long-term smoking exposure. In addition, they did not fully stratify by smoking status, which our results suggest is critical to address residual confounding. As a sensitivity analysis we also examined quartiles of BCD and similar to the results presented in Table 2, the association between upper quartile of BCD with depressive symptoms disappeared after stratification by smoking status and adjustment for pack-years and cotinine.

Our study is limited by several factors, foremost the cross-sectional nature of the NHANES dataset. It is, however, a large data set with weighting parameters to offset potential sampling bias and make the data generalizable to the US population. The questionnaire data in NHANES is limited as well by self-report, although questions for depressive symptoms and smoking have been validated (Kroenke, 2002; Yeager and Krosnick, 2010) and serum cotinine is also used to supplement answers to smoking questions. Future studies might include prospective data collection, tracking smoking history across time and collecting baseline BCD measurements, and measures of subsequent depressive symptoms to better understand the associations between BCD, smoking, and depression.

In studies of health outcomes that are also associated with smoking, BCD may be confounded by smoking. Stratifying by smoking status and adjusting for pack-years and serum cotinine should be used to fully control for confounding by smoking in epidemiologic studies of BCD.

Acknowledgments

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Bakhshaie J, Zvolensky MJ, Goodwin RD, 2015 Cigarette smoking and the onset and persistence of depression among adults in the United States: 1994–2005. *Compr.Psychiatry* 60, 142–148. [PubMed: 25882595]
- Berk M, Williams LJ, Andreazza AC, Pasco JA, Dodd S, Jacka FN, Moylan S, Reiner EJ, Magalhaes PV, 2014 Pop, heavy metal and the blues: secondary analysis of persistent organic pollutants (POP), heavy metals and depressive symptoms in the NHANES National Epidemiological Survey. *BMJ Open* 4 (7), e005142.
- Bramer SL, Kallungal BA, 2003 Clinical considerations in study designs that use cotinine as a biomarker. *Biomarkers* 8 (3–4), 187–203. [PubMed: 12944172]
- Ellingsen DG, Thomassen Y, Aaseth J, Alexander J, 1997 Cadmium and selenium in blood and urine related to smoking habits and previous exposure to mercury vapour. *J. Appl. Toxicol* 17 (5), 337–343. [PubMed: 9339747]
- Han C, Lim YH, Hong YC, 2016 Does cadmium exposure contribute to depressive symptoms in the elderly population? *Occup. Environ. Med* 73 (4), 269–274. [PubMed: 26769897]
- Jarup L, Akesson A, 2009 Current status of cadmium as an environmental health problem. *Toxicol. Appl. Pharmacol* 238 (3), 201–208. [PubMed: 19409405]

- Jarup L, Berglund M, Elinder CG, Nordberg G, Vahter M, 1998 Health effects of cadmium exposure – a review of the literature and a risk estimate. *Scand. J. Work Environ. Health* 24 (Suppl 1), 1–51.
- Kroenke KS, Spitzer Robert L., 2002 The PHQ-9: a new depression diagnostic and severity measure. *Psychiatr. Ann* 32 (9), 1–7.
- Mendez-Armenta M, Rios C, 2007 Cadmium neurotoxicity. *Environ. Toxicol. Pharmacol* 23 (3), 350–358. [PubMed: 21783780]
- Milette K, Hudson M, Baron M, Thombs BD, 2010 Comparison of the PHQ-9 and CES-D depression scales in systemic sclerosis: internal consistency reliability, convergent validity and clinical correlates. *Rheumatology* 49 (4), 789–796. [PubMed: 20100794]
- NCHS, 2016a Continuous National Health and Nutrition Examination Survey (NHANES) Web Tutorial Centers for Disease Control and Prevention, National Center for Health Statistics, Atlanta, GA.
- NCHS, 2016b NHANES- National Health and Nutrition Examination Survey Homepage Centers for Disease Control and Prevention, National Center for Health Statistics, Atlanta, GA.
- Olabanji ON, J. C, Msagati TAM, Oluyemi EA, Fatoye FO, Mamba BB, 2011 Effect of metal poisoning and the implications of gender and age on the elemental composition in patients with mental behavioural disorders. *Afr. J. Biotechnol* 10 (8), 3585–3593.
- Orisakwe OE, 2014 The role of lead and cadmium in psychiatry. *N. Am. J. Med. Sci* 6 (8), 370–376. [PubMed: 25210669]
- Scinicariello F, Buser MC, 2015 Blood cadmium and depressive symptoms in young adults (aged 20–39 years). *Psychol. Med* 45 (4), 807–815. [PubMed: 25115444]
- Shiue I, 2015 Urinary heavy metals, phthalates and polyaromatic hydrocarbons independent of health events are associated with adult depression: USA NHANES, 2011–2012. *Environ. Sci. Pollut. Res. Int* 22 (21), 17095–17103. [PubMed: 26126689]
- Yeager DS, Krosnick JA, 2010 The validity of self-reported nicotine product use in the 2001–2008 National Health and Nutrition Examination Survey. *Med. Care* 48 (12), 1128–1132. [PubMed: 20940652]

Table 1Demographics of cohort ($n = 11,209$).

	Depressive symptoms +	Depressive symptoms –	<i>p</i> -value (Chi-square test or <i>t</i> -test)
Total <i>n</i> (%)	876 (8)	10,333 (92)	
Female <i>n</i> (%)	532 (61)	4621 (45)	<0.01
Male <i>n</i> (%)	344 (39)	5712 (55)	
Race <i>n</i> (%)			<0.01
Caucasian	392 (45)	5307 (51)	
Black	212 (24)	1947 (19)	
Mexican- American	127 (14)	1584 (15)	
Other Hispanic	102 (12)	825 (8)	
Other	43 (5)	670 (7)	
PIR mean $\pm$ SD	1.79 $\pm$ 1.47	2.84 $\pm$ 1.65	<0.01
Age mean $\pm$ SD	44.06 $\pm$ 14.44	46.70 $\pm$ 17.35	<0.01
Blood Cd μ g/L mean $\pm$ SD	0.79 $\pm$ 0.84	0.53 $\pm$ 0.62	<0.01
Pack years mean $\pm$ SD	12.1 $\pm$ 20.3	8.2 $\pm$ 17.4	<0.01
Blood cotinine ng/ml mean $\pm$ SD	110.5 $\pm$ 156.3	60.9 $\pm$ 127.8	<0.01
Current smokers <i>n</i> (%)	406 (46)	2480 (24)	<0.01
Never smokers <i>n</i> (%)	322 (37)	5372 (52)	
Former smokers <i>n</i> (%)	148 (17)	2481 (24)	

Depressive symptoms + was defined by a score greater than or equal to 10 on the PHQ-9.

PIR = Poverty Index Ratio.

Cd = Cadmium.

Table 2

Logistic regression of the association between blood cadmium and depressive symptoms.

	<i>N</i>	<i>N</i>	β	OR (95% CI)
	Depressive symptoms +			
Unadjusted	11,209	876	0.43	1.53 (1.37, 1.71)
Model 1- Adjusted for age, sex, race, PIR, and alcohol use	11,209	876	0.31	1.37 (1.24, 1.51)
Model 2- Also adjusted for blood cotinine and pack-years smoking	11,209	876	0.15	1.16 (1.04, 1.30)
Model 2, Stratified by Smoking Status				
Never Smokers ^a	5694	322	– 0.36	0.70 (0.33, 1.47)
Former Smokers	2629	148	– 0.41	0.67 (0.28, 1.58)
Current Smokers	2886	406	0.07	1.08 (0.95, 1.22)

Depressive symptoms + was defined by a score greater than or equal to 10 on the PHQ-9.

PIR = Poverty Index Ratio.

^aNot also adjusted for pack years since all were equal to zero among never smokers.

Table 3

Blood cadmium levels in $\mu\text{g/L}$ (mean $\pm$ SD) stratified by smoking status and depressive symptoms.

	All	Depressive symptoms +	Depressive symptoms –	<i>p</i> -value (<i>t</i> -test)
All	0.55 $\pm$ 0.64	0.79 $\pm$ 0.84	0.53 $\pm$ 0.62	< 0.01
Current smokers	1.18 $\pm$ 0.95	1.32 $\pm$ 0.96	1.16 $\pm$ 0.90	< 0.01
Never smokers	0.29 $\pm$ 0.21	0.29 $\pm$ 0.16	0.29 $\pm$ 0.21	1.0
Former smokers	0.40 $\pm$ 0.27	0.39 $\pm$ 0.28	0.40 $\pm$ 0.26	0.65

Depressive symptoms + was defined by a score greater than or equal to 10 on the PHQ-9. Raw data. Not adjusted for cotinine or pack years.