Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

ORIGINAL RESEARCH

Is hip strengthening the best treatment option for females with patellofemoral pain? A randomized controlled trial of three different types of exercises

Marcelo Camargo Saad^{a,b,*}, Rodrigo Antunes de Vasconcelos^c,
Letícia Villani de Oliveira Mancinelli^d, Matheus Soares de Barros Munno^d,
Rogério Ferreira Liporaci^b, Débora Bevilaqua Grossi^a

^a Programa de Pós-graduação em Reabilitação e Desempenho Funcional, Faculdade de Medicina de Ribeirão Preto (FMRP), Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

^b Programa de Biomecânica, Medicina e Reabilitação do Aparelho Locomotor, Faculdade de Medicina de Ribeirão Preto (FMRP), Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

^c Núcleo de Estudos Instituto Wilson Mello, Campinas, SP, Brazil

^d Curso de Fisioterapia, Faculdade de Medicina de Ribeirão Preto (FMRP), Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

Received 26 September 2017; received in revised form 14 February 2018; accepted 22 March 2018

Available online 4 April 2018

KEYWORDS

Anterior knee pain;
Rehabilitation;
Step up and down
kinematics;
Quadriceps and hip
muscles.

Abstract

Objective: To evaluate the effect of three types of exercise intervention in patients with patellofemoral pain and to verify the contributions of each intervention to pain control, function, and lower extremity kinematics.

Methods: A randomized controlled, single-blinded trial was conducted. Forty women with patellofemoral pain were randomly allocated into four groups: hip exercises, quadriceps exercises, stretching exercises and a control group (no intervention). Pain (using a visual analog scale), function (using the Anterior Knee Pain Scale), hip and quadriceps strength (using a handheld isometric dynamometer) and measuring lower limb kinematics during step up and down activities were evaluated at baseline and 8 weeks post intervention.

Results: All treatment groups showed significant improvements on pain and Anterior Knee Pain Scale after intervention with no statistically significant differences between groups except when compared to the control group. Only hip and quadriceps groups demonstrated improvements in muscle strength and knee valgus angle during the step activities.

* Corresponding author at: Prédio da Fisioterapia e Terapia Ocupacional da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo, Avenida Bandeirantes, 3900, Bairro Monte Alegre, CEP: 14049-900, Ribeirão Preto, SP, Brazil.

E-mail: marfisiousp@yahoo.com.br (M.C. Saad).

Conclusion: Hip strengthening exercises were not more effective for pain relief and function compared to quadriceps or stretching exercises in females with patellofemoral pain. Only hip and quadriceps groups were able to decrease the incidence of dynamic valgus during step-down activity. This study was approved by Brazilian Clinical Trials Registry registration number: RBR-6tc7mj ([http://www.ensaiosclinicos.gov.br/rg/RBR-6tc7mj/](http://www ensaiosclinicos.gov.br/rg/RBR-6tc7mj/)).

© 2018 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Patellofemoral pain (PFP) is a prevalent clinical condition that is very commonly observed in orthopedic practice.^{1–4} Dysfunctions in the patellofemoral joint are especially common among physically active young adults, with a 2.2 times higher prevalence in women, although it occurs frequently in men as well.^{5,6} Symptoms related to PFP include pain associated with movement, particularly with activities such as kneeling, ascending and descending stairs, running and squatting.⁷

The rehabilitation of these patients is increasingly important, as 45% of those with PFP may suffer from osteoarthritis with advanced age⁸ and 22% of individuals who have undergone unicompartmental arthroplasty of the knee reported a history of PFP at a younger age.⁹

In recent years, the role of hip stabilizers has been the object of several studies and rehabilitation protocols.^{10–13} Patients with PFP often present with pelvic misalignment and increased internal rotation of the femur during the execution of functional activities, such as ascending and descending stairs that are functionally two of the most painful movements of daily living for patients with PFP.¹⁴

The theory about the changed position of patella not being caused solely by quadriceps imbalance, but by changing position of the femur in closed kinetic chain activities due to a hip weakness gained strength and it has been hypothesized that the cause of PFP include exercises to the hip and pelvis which would be the key to satisfactory clinical outcomes for PFP patients.¹⁵ Thus, strengthening of hip stabilizing muscles, especially the abductors and lateral rotators, as indicated by Powers,¹⁵ has changed the opinion and conduct of treatment proposed by clinicians and researchers.

It has not been established which type of conservative treatment is more effective because the majority of studies have used different groups of treatment with a common protocol of exercises to all groups or the association of different types of exercise in the same protocol.^{16–18} Other studies have proposed the strengthening of the same muscle group, distinguishing only the method of performance.^{19–21} Besides, no studies were found that assessed the effect of stretching exercises alone, only those that included stretching muscle groups that were common to all groups of treatment.^{18,21,22} So we do not know what the real effect, in an isolated way, that a treatment based solely on stretching exercises can bring to individuals with PFP. Thus, no study has evaluated

each form of treatment in an isolated way, without any combination of them.

There are currently seven prospective randomized controlled trials in the literature that have investigated the inclusion of hip and pelvic muscle strengthening for PFP^{12,18,20,23–26} patients, but the conclusions of these studies regarding the benefits of proximal muscles strengthening are questionable. Because common exercises for hip and quadriceps, as well as stretching were associated in these protocols, and results were either only compared to no intervention groups²⁶ or included exercises to strengthen adductor muscles, contrary to idea of strengthening abductors and external rotators in order to stabilize the pelvis.^{23,27} However the design of these studies was based on the knowledge available at the time of those studies.^{12,20,25}

Thus, our purpose was to evaluate the contribution of three types of individual interventions, in patients with PFP and to verify the contributions of each one in terms of decreasing pain, and improving function and lower extremity kinematics.

Methods

A 4arm, randomized controlled assessor-blinded trial was performed.

Patients

Forty recreational female athletes between the ages of 18 and 28 years with PFP were selected from the university campus and randomly assigned to groups for treatments. A recreational athlete was defined as anyone participating in aerobic or athletic activity at least 3 times per week for at least 30 min.²³

Subjects were included in the study if they were female and had anterior knee pain with a minimum intensity of 3 or greater on the 10-cm visual analog scale (VAS)^{1,23} for at least three months before the study assessment. Additional inclusion and exclusion criteria were followed according to Nakagawa et al.²⁸ Inclusion criteria: (1) insidious onset of symptoms; (2) retropatellar or peripatellar pain with at least 2 of the following activities (ascending/descending stairs, running, kneeling, squatting, prolonged sitting or jumping). Exclusion criteria: (1) previous history of knee surgery; (2) history of back, hip, or ankle joint injury or pain; (3) patellar instability; (4) lesion or pain during palpation or test of any

structure of knee and (6) any neurological involvement that would affect gait.

Of the 40 volunteer subjects included in the project, all subjects first underwent a physical, kinematic, and strength assessment. This study and the informed consent form related to this same study were approved by the Ethics Committee of the Ribeirão Preto Medical School, Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil (reference number: 10371/2010). All volunteers read and signed the informed consent document.

This study was prospectively registered in the Brazilian Clinical Trials Registry (REBEC), registration number: RBR-6tc7mj.

Randomization

The PFP subjects were randomly allocated to one of four groups: quadriceps strengthening group (QG), hip strengthening group (HG), stretching group (SG) or a control group (CG) (no treatment). The randomization schedule was generated using R 2.7.2 statistical software. The allocation was concealed by the use of consecutively numbered, sealed and opaque envelopes (Fig. 1).

Outcome measures

All kinematic assessments were performed by RFL (author), who had at least 4 years experience doing these procedures. Only LVOM, MSBM and RFL (authors) were blinded to the group allocation, because MCS (author) provided all treatments to the groups. LVOM and MSBM were also responsible for the administration of the questionnaires, intensity of pain assessment and muscle strength assessment tests.

There were two evaluations. The initial and final evaluation post 8 weeks consisted of an assessment of pain,²⁹ a functional questionnaire,² assessment of muscle strength through a handheld isometric dynamometer,^{30,31} and dynamic valgus were assessed during functional activities (ascending and descending stairs).

Pain

Intensity of pain, the primary endpoint of our study, was assessed using a visual analog scale (VAS) on a 10 cm scale, where 0 indicated "no pain" and 10 indicated "worst possible pain". This scale is considered a reliable, valid, and responsive method to examine individuals with PFP.²⁹

Function

Physical function was assessed by an Anterior Knee Pain Scale (AKPS),² which is a specific questionnaire for individuals with PFP. This tool is valid for individuals with PFP and has proved to be highly reliable in test-retest,²⁹ in addition, it was translated, adapted, and validated in Brazil.²

Kinematics

For kinematic data analysis, the *Qualisys® Pro-Reflex* video camera system³² with six infrared video cameras was used,

which allowed for the reconstruction in 3D of passive reflective markers located at specific bony prominences (i.e. hip, knee, and ankle).¹⁶ The frequency of data capture was 120 Hz, and standard deviation errors below 10 mm were allowed during the initial calibration.

Participants were asked to ascend and descend a three-step stair at normal speed, simulating a usual everyday situation. The mean values obtained from three ascending and descending attempts were used for statistical analyses.

Dynamic valgus was measured in the frontal-plane at 45 degrees of tibiofemoral flexion because this was the best moment to assess the valgus, and altered patterns of movement could be exacerbated in this position.^{33,34}

Strength evaluation

Using a load cell (KRATOS®) adapted (handheld dynamometer) to a digital reader, the isometric muscle strength of the hip abductors (gluteus medius), adductors, lateral rotators, medial rotators, and hip extensors (gluteus maximus) and flexors (iliopsoas and rectus femoris) was measured. For knee extensors (quadriceps) and flexors (hamstrings), the load cell was adapted and fixed with chains to a flexor-extensor machine and the MVC isometric muscle strength was measured in a favorable position^{30,35–37,60} and properly stabilized with a belt so as to avoid any possible influence by the examiner. The values were normalized to each individual's body mass.

Interventions

Patients included in the treatment groups participated in two sessions per week for eight weeks with a minimum break of 24 h between sessions. Individuals were asked and requested not to perform exercises outside the program and all treatment sessions were performed in the university's posture and human movement analysis laboratory (LAPOMH). Each treatment session was approximately 50 min in duration, and all sessions were performed individually and supervised by the same physical therapist. For all groups of treatment the weights were increasing as the patient's reported changes in their following the Rating of Perceived Exertion (RPE) scale³⁸ based on Borg's Scale of effort.³⁹ This is common method used to assess the intensity of exercise and this scale has been used and researched for its use in both clinical and exercise settings.^{39,40} The exercises chosen for each group should strengthen the muscles of the hip & knee & lead to increased stabilization of the hip & knee joints:

- *Quadriceps strengthening group (QG)*. The exercises in this group focused specifically on quadriceps strengthening (see Appendix A).
- *HIP strengthening group (HG)*. This group performed exercises to strengthen hip stabilizing muscles (see Appendix B).
- *Stretching group (SG)*. In this group, the physical therapist monitored and stabilized the patients during the stretching exercises for all muscles involved in knee and hip stabilization (see Appendix C).

Figure 1 CONSORT flowchart for subject selection for hip & knee muscle strength in PFP subjects. ITT, intention to treat.

- **Control group (CG).** Patients included in this group did not have any kind of intervention for eight weeks, but they were tested at the start of the program & at the end like the other 3 groups. At the end of the experiment, a rehabilitation program was made available to all patients.

Statistical analysis

Sample size calculation: forty PFP patients (10 in each group) were needed to provide 80% power, based on a 1-sided test, type I error of 0.05, standard deviation of 1.8 (from prior data), and the ability to detect a 3-cm between-group difference in pain intensity based on a 10-cm VAS.²³

A linear regression model with mixed effects (i.e. random and fixed) was used. Mixed-effect linear models are used in the analysis of data in which the answers are grouped (i.e. repeated values for the same individual) and the assumption of independence of observations within a group is not adequate.⁴¹ A significance level of 5% was defined.

For analysis of dynamic valgus, a qualitative analysis of data was chosen, which was performed using the McNemar's test,⁴² and which aimed to assess the real effect of a therapeutic intervention within treatment groups; results are expressed in absolute values and a percentage.

All participants attended all treatment sessions except the quadriceps group that computed a loss. For the quadriceps group, the statistical analysis took into account the loss of one individual from the group. An intention-to-treat (ITT) analysis approach was carried out. The missing data was dealt with by using a multiple imputation.⁴³ In addition, a "per protocol" analysis was also performed using patients with complete data only. For statistical analysis, only the symptomatic leg was considered.

Clinical relevance

In order to assess clinical relevance, the MCID (minimal clinically important difference) was performed and an average improvement exceeding the minimum of 2 cm on the VAS is considered clinically relevant.²⁹

Results

Table 1 summarizes the demographic characteristics of the groups at the baseline.

Pain

VAS revealed that, except for the CG, all groups had improvements in post-treatment period. Between-group comparison showed no differences between treatment groups after the 8 week exercise period and all groups showed better performance than the CG (**Table 2**).

All treatments groups had a significant MCID for pain outcomes at the end of 8 week exercise program, except the CG. Between-group comparisons were performed in order to compare which treatment protocol was more effective and none of the three groups had a greater treatment effect than the other two except when compared to the CG (**Table 2**).

Function

For between-group comparison questionnaire scores pre and post-treatment, all groups except the CG improved their scores ($p < 0.01$). No difference was found between the 3 treatment groups. All 3 groups showed better performance than the CG (**Table 2**).

Table 1 Female patient demographics for muscle strengthening & stretching in PFP patients at baseline.

Demographic characteristics	Quadriceps (n = 10)	HIP (n = 10)	Stretching (n = 10)	Control (n = 10)
Age (years)	23.2 ± 2.53	22.5 ± 1.08	21.3 ± 1.16	23.2 ± 1.03
Height (m)	1.61 ± 0.07	1.59 ± 0.03	1.58 ± 0.02	1.61 ± 0.06
Mass (kg)	56.25 ± 5.89	55.32 ± 3.99	54.72 ± 2.24	55.37 ± 2.03
BMI (kg/m ²)	21.80 ± 1.72	22.0 ± 2.0	21.92 ± 1.32	21.34 ± 1.33

Table 2 Pain (VAS) pre and post treatment. Anterior Knee Pain Score (AKPS) pre and post treatment for female PFP patients. Final criteria for clinical relevance on pain results based on MCID to the treatment groups and final results comparison between treatment group.

Group	VAS (cm)		AKPS		
	Pre	Post	Pre	Post	
Quadriceps group (n = 10)	6.34 ± 1.85	0.56 [*] ± 0.89	76.9 ± 8.72	90.11 [*] ± 6.11	
Hip group (n = 10)	5.05 ± 1.27	0.55 [*] ± 0.8	77.4 ± 5.46	91.8 [*] ± 5.67	
Stretching group (n = 10)	4.08 ± 0.82	0.14 [*] ± 0.25	79.8 ± 9.21	91.0 [*] ± 6.62	
Control group (n = 10)	4.03 ± 1.03	3.69 ± 0.6	80.8 ± 10.67	81.9 ± 8.41	
VAS	Between-group postMean difference		95% confidence interval	p	Final criteria
HG × QG	−0.12	−1.03	0.79	0.79	NCR
HG × SG	−0.41	−1.30	0.48	0.36	NCR
HG × CG	3.14	2.25	4.03	<0.01	CR
QG × SG	−0.53	−1.44	0.38	0.24	NCR
QG × CG	3.02	2.11	3.93	<0.01	CR
SG × CG	−3.55	−4.44	−2.66	<0.01	CR

HG, hip group; QG, quadriceps group; SG, stretching group; CG, control group; CR, clinically relevant; NCR, not clinically relevant.

^{*} p < 0.01.

Strength

The post-treatment period revealed strength improvement for HG (i.e. hip abductors, adductors, internal/external rotators and extensors); QG (hip and knee flexors); CG (decrease hip extensors); SG (no significant changes) (Table 3).

Kinematics

Step-down: The QG and HG had improvement in movement patterns in the post-treatment testing. The HG had 80% of individuals with a valgus knee alignment and 20% varus pattern pre-treatment. Post-treatment, 30% of individuals still had a valgus alignment whereas 70% presented with a varus pattern. For the QG, 88.89% of individuals had a knee valgus alignment and 11.11% had a varus pattern pre-treatment. Post-treatment: only 11.11% of subjects still had a valgus alignment whereas 88.89% showed a varus pattern. For the step-up, only the HG had improvement. Pre-treatment, 100% had a knee valgus pattern. Post-treatment, 40% still had this pattern and 60% showed a knee varus pattern (Table 4).

Discussion

Regarding pain and function, our results showed that the three most common forms of treatment for PFP patients,

with focus on quadriceps strengthening, strengthening and stabilization of the hip, and stretching were equally effective and no significant differences were observed among them. These results confirmed previous clinical trials that showed that treatment, regardless of the physical intervention applied, could be beneficial to PFP individuals.^{18,19,44–49}

Our results confirmed that prescription of stretching exercises relieved pain and increased function similarly to effects of strengthening programs and stabilizing hips and knees for PFP patients. Reduced hamstring length can result in increased stress on the patellofemoral joint,⁵⁰ thus, increased flexibility might prevent the development of such pathologies as PFP.^{16,51} Our study is the first to analyze, in an isolated way, the role of muscle stretching in PFP individuals, and the results suggested that stretching exercises were equally effective for pain relief and functional improvements in PFP patients.

Recently, Khayambashi et al.,¹² in a comparative study, concluded that hip strengthening was better than quadriceps strengthening but the quadriceps group were exercised using different isometric angles that were not protective and increased the stress to patellofemoral joint⁵² and used strengthening exercises combined with the strengthening hip adductors that is completely contrary to idea of strengthening hip abductors and external rotators to help stabilize the pelvis.¹⁵ The study design led to a bias toward positive results for the group undergoing hip strengthening. However, Rathleff et al.,⁵³ in a systematic review,

Table 3 Pre and post treatment effects on muscle strength (Kgf) for female PFP patients.

	Quadriceps (N = 10)		Hip (n = 10)	
	Pre	Post	Pre	Post
ABD HIP	10.96 ± 2.99 (8.82–13.1)	11.34 ± 3.74 (8.47–14.22)	11.87 ± 2.68 (9.95–13.78)	15.14 ± 2.48 (13.37–16.92)*
ADD HIP	8.22 ± 3.55 (5.69–10.76)	8.09 ± 2.61 (6.08–10.1)	9.99 ± 2.3 (8.34–11.63)	11.13 ± 1.69 (9.93–12.34)*
EXT HIP	16.82 ± 5.99 (12.54–21.11)	17.74 ± 6.99 (12.37–23.11)	20.25 ± 5.3 (16.47–24.04)	23.67 ± 2.93 (21.58–25.77)*
FLEX HIP	11.82 ± 2.82 (9.8–13.83)	13.97 ± 3.55 (11.25–16.7)*	14.94 ± 2.99 (12.8–17.08)	16.55 ± 3.53 (14.02–19.07)
IR HIP	8.48 ± 2.69 (6.56–10.41)	8.59 ± 2.75 (6.48–10.71)	8.88 ± 3.36 (6.48–11.29)	10.99 ± 2.95 (8.88–13.09)*
ER HIP	7.44 ± 1.52 (6.35–8.52)	7.76 ± 1.87 (6.32–9.19)	7.15 ± 1.21 (6.28–8.01)	8.9 ± 1.07 (8.13–9.67)*
KNEE EXT	20.86 ± 9.17 (14.3–27.42)	25.26 ± 11.16 (16.68–33.85)	30.38 ± 10.69 (22.73–38.03)	35.23 ± 7.28 (30.02–40.44)
KNEE FLEX	8.95 ± 4.02 (6.08–11.83)	13.46 ± 11.02 (4.99–21.93)*	12.76 ± 4.09 (9.83–15.69)	14.07 ± 3.02 (11.91–16.23)
	Stretching (N = 10)		Control	
	Pre	Post	Pre	Post
ABD HIP	13.02 ± 6.02 (8.71–17.33)	13.29 ± 5.45 (9.4–17.19)	12.06 ± 3.58 (9.5–14.62)	11.4 ± 3.22 (9.1–13.7)
ADD HIP	9.61 ± 4.43 (6.44–12.78)	9.62 ± 4.57 (6.36–12.89)	9.32 ± 1.43 (8.3–10.34)	8.77 ± 1.68 (7.57–9.97)
EXT HIP	19.3 ± 7.57 (13.88–24.71)	21.25 ± 8.33 (15.29–27.21)	22.74 ± 6.03 (18.43–27.05)	19.76 ± 6.18 (15.34–24.19)*
FLEX HIP	14.6 ± 4.95 (11.06–18.14)	15.02 ± 4.49 (11.81–18.23)	13.42 ± 3.29 (11.07–15.77)	13.42 ± 4.03 (10.54–16.3)
IR HIP	9.61 ± 4.18 (6.63–12.6)	10.4 ± 4.18 (7.41–13.39)	10.0 ± 2.14 (8.47–11.53)	9.14 ± 1.86 (7.81–10.48)
ER HIP	8.03 ± 3.49 (5.53–10.52)	8.15 ± 2.86 (6.1–10.2)	7.72 ± 1.27 (6.81–8.62)	7.09 ± 1.61 (5.94–8.25)
KNEE EXT	32.07 ± 13.56 (22.37–41.77)	31.85 ± 15.26 (20.93–42.77)	39.71 ± 9.54 (32.89–46.53)	36.76 ± 10.44 (29.29–44.22)
KNEE FLEX	14.15 ± 7.08 (9.08–19.21)	13.71 ± 5.07 (10.09–17.34)	12.53 ± 3.46 (10.06–15.01)	11.39 ± 2.98 (9.26–13.53)

ABD HIP, abductors of hip; ADD HIP, adductors of hip; EXT HIP, extensors of hip; FLEX HIP, flexors of hip; IR HIP, internal rotators of hip; ER HIP, external rotators of hip; KNEE EXT, knee extensors; KNEE FLEX, knee flexors.

Values are mean ± SD; Values in parentheses are 95% confidence interval.

* $p < 0.01$.

Table 4 Pre and post treatment effects on lower limb kinematics for female PFP patients.

Group	HIP kinematics (n(%)) [†]							
	Step up				Step down			
	Valgus		Varus		Valgus		Varus	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post
PFP (n = 39)	34 (87.18)	26 (66.67)*	5 (12.82)	13 (33.33)*	32 (82.05)	19 (48.72)*	7 (17.95)	20 (51.28)*
Quadriceps	8 (88.89)	5 (55.56)	1 (11.11)	4 (44.44)	8 (88.89)	1 (11.11)*	1 (11.11)	8 (88.89)*
Hip	10 (100)	4 (40)*	0 (0)	6 (60)*	8 (80)	3 (30)*	2 (20)	7 (70)*
Stretching	7 (70)	8 (80)	3 (30)	2 (20)	7 (70)	5 (50)	3 (30)	5 (50)
Control	9 (90)	9 (90)	1 (10)	1 (10)	9 (90)	10 (100)	1 (10)	0 (0)

* Statistically different from pre-treatment ($p < 0.01$).

[†] Valgus or Varus presence, values expressed in number and percentage of individuals.

found no association between weakness of the hip muscles and the development of PFP, therefore reduced hip strength could be a result and not the cause of PFP. Rabelo and Lucareli⁵⁴ agreed with this hypothesis. In two other systematic reviews performed by Lack et al.⁵⁵ and Santos et al.,⁵⁶ it was observed that the strengthening of proximal structures (hip) combined with quadriceps strengthening, showed strong evidence of improvement in pain(decreased) and function(improved) in the short term, but with no evidence of improvement or superiority in the long term when compared to isolated quadriceps strengthening. These reviews showed that hip muscle strengthening was effective in reducing the intensity of pain and improving functional

capabilities in patients with PFP, despite the lack of evidence for its ability to increase muscle strength.

Our study had the objective of completely differentiating the treatment groups, so none of the protocols had any exercises in common, to assess the real effect of specific exercises. In previous studies, some protocols often used exercises that were common to different protocols.^{16–18,23} Others focused on one muscle group and distinguished only the way the strengthening was carried out,^{19–21} or included stretching exercises for muscle groups that were common to all treatment groups.^{18,21,22} Therefore, the question is whether the improvement observed could have occurred as a result of a specific exercise for strengthened muscle,

regardless of the way it was strengthened, or whether this improvement occurred due to the common stretching exercises proposed. As a consequence, this possibly resulted in no significant differences being observed between treatments.

The qualitative analysis of movement showed at baseline that PFP individuals had a valgus movement pattern of around 87.18% and 82.05% during step-up and step-down, respectively. At the end of the intervention period, these values decreased to 66.67% and 48.72% respectively. Stratifying into subgroups, only the HG and QG had significant improvements at the end of the 8 weeks intervention period. Step-up had a greater incidence of dynamic valgus, agreeing with a previous study⁵⁷ that showed step-up was more unstable for PFP individuals and could intensify altered movement patterns.

Recent studies reported that strengthening proximal structures could alter incorrect patterns of movements that resulted in excessive dynamic valgus and consequently increased pain in PFP patients. However, these were conceptual statements and kinematics were not assessed.^{11,15,24,26,33} Herman et al.⁵⁸ and Earl and Hoch²⁷ also reported that individual hip strengthening exercises did not alter movement patterns. Recently, an exercise program proposed by Baldon et al.²³ was able to change movement patterns in PFP individuals, agreeing with our results. Our study also found that only the strengthening groups (i.e. HG and QG) had significant improvement in their movement patterns. Nevertheless, despite this improvement in the HG and QG, results in pain and function did not show greater improvement than those in the SG.

In the assessment of muscle strength, only the HG had significant improvement in almost all muscle groups of the hip (i.e. abductors, adductors, extensors, and internal and external rotators). This difference was probably due to the joint position at the moment of evaluation because muscles such as the gluteus maximus and medius which are lateral rotators in the orthostatic position, and anterior fibers of these muscles become medial rotators in the sitting position (i.e. the position chosen during the strength test).⁵⁹ The QG showed strength improvement only for hip flexors and knee flexors. This can be explained by the strengthening of the quadriceps themselves, which act as a biarticular muscle that also facilitates hip flexion.⁵⁹

All groups that received any kind of treatment showed statistical and clinically relevant results in pain relief at the end of the 8 week intervention period, with average improvement greater than the minimum clinically important difference (MCID) for pain of 2 cm VAS.²⁹ At the conclusion of the 8 week intervention period, in order to compare which protocol of treatment was more effective, final results for pain were compared and no group was superior to the others except when compared to the control group that received no treatment.

Above all, the present study highlighted the benefits of physiotherapy treatment that, regardless of the form of intervention, contribute to relieve pain, either by stabilizing and strengthening the hip and quadriceps muscles, or by improving the flexibility of the structures involved. No studies were found in the literature that compared individual treatment protocols to completely different ones, as was done in this study. Some more recent studies have

stated that an association of programs would be even more beneficial to patients with PFP than an isolated program that focuses on only one type of treatment. Thus, the most indicated and effective rehabilitation for each individual should be based on the deficits observed in the initial evaluation of each individual.

Limitations

Our study had some limitations. First, despite the large number of total patients with PFP, the stratified treatment groups were eventually small; nevertheless, sample calculation allowed us to use this number of individuals. Second, participants' baseline characteristics seemed to be different, especially for pain, probably as a consequence of low number of participants. Third, the current findings should not be extrapolated to males with PFP, as only women participated in this study. And third, 8 weeks was a short time for a follow-up but we know that this is a period that we can already observe pain relief in these patients.^{12,23}

Conclusion

Hip strengthening exercises was no more effective on pain and function compared to quadriceps exercises or stretching exercises in females with PFP. Both hip and quadriceps strengthening exercises improved lower extremity kinematics after 8 weeks of intervention.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgement

This study received financial support of The State of São Paulo Research Foundation – FAPESP (process number: 2010-/12561-9).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.bjpt.2018.03.009](https://doi.org/10.1016/j.bjpt.2018.03.009).

References

1. Cowan SM, Crossley KM, Bennell KL. Altered hip and trunk muscle function in individuals with patellofemoral pain. *Br J Sports Med*. 2009;43(8):584–588.
2. Da Cunha RA, Costa LO, Hespanhol Junior LC, Pires RS, Kujala UM, Lopes AD. Translation, cross-cultural adaptation, and clinical testing of instruments used to assess patients with patellofemoral pain syndrome in the Brazilian population. *J Orthop Sports Phys Ther*. 2013;43(5):332–339.
3. Davis IS, Powers CM. Patellofemoral pain syndrome: proximal, distal, and local factors, an international retreat, April 30–May 2, 2009, Fells Point, Baltimore, MD. *J Orthop Sports Phys Ther*. 2010;40(3):A1–A16.

4. Wood L, Muller S, Peat G. The epidemiology of patellofemoral disorders in adulthood: a review of routine general practice morbidity recording. *Prim Health Care Res Dev*. 2011;12:157–164.
5. Boling M, Padua D, Marshall S, Guskiewicz K, Pyne S, Beutler A. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. *Scand J Med Sci Sports*. 2010;20:725–730.
6. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther*. 2010;40:42–51.
7. Cook C, Hegedus E, Hawkins R, Scovell F, Wyland D. Diagnostic accuracy and association to disability of clinical test findings associated with patellofemoral pain syndrome. *Physiother Can*. 2010;62:17–24.
8. Stathopulu E, Baildam E. Anterior knee pain: a long-term follow-up. *Rheumatology (Oxford)*. 2003;42(2):380–382.
9. Utting MR, Davies GJ, Newman JH. Is anterior knee pain a predisposing factor to patellofemoral osteoarthritis? *Knee*. 2005;12:362–365.
10. Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Lattermann C, Uhl TL. Hip strengthening prior to functional exercises reduces pain sooner than quadriceps strengthening in females with patellofemoral pain syndrome: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2011;41:560–570.
11. Fukuda TY, Melo WP, Zaffalon BM, et al. Hip posterolateral musculature strengthening in sedentary women with patellofemoral pain syndrome: a randomized controlled clinical trial with 1-year follow-up. *J Orthop Sports Phys Ther*. 2012;42(Oct (10)):823–830.
12. Khayambashi K, Fallah A, Movahedi A, Bagwell J, Powers C. Posterolateral hip muscle strengthening versus quadriceps strengthening for patellofemoral pain: a comparative control trial. *Arch Phys Med Rehabil*. 2014;95(May (5)):900–907.
13. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther*. 2009;39(1):12–19.
14. Brechter JH, Powers CM. Patellofemoral joint stress during stair ascent and descent in persons with and without patellofemoral pain. *Gait Posture*. 2002;16:115–123.
15. Powers CM. The influence of altered lower-extremity kinematics on patellofemoral joint dysfunction: a theoretical perspective. *J Orthop Sports Phys Ther*. 2003;33(11):639–646.
16. Collins N, Crossley K, Beller E, Darnell R, McPoil T, Vicenzino B. Foot orthoses and physiotherapy in the treatment of patellofemoral pain syndrome randomised clinical trial. *Br J Sports Med*. 2009;43:163–168.
17. Collins NJ, Crossley KM, Darnell R, Vicenzino B. Predictors of short and long term outcome in patellofemoral pain syndrome: a prospective longitudinal study. *BMC Musculoskelet Disord*. 2010;19(11):11.
18. Nakagawa TH, Muniz TB, Baldon Rde M, Dias Maciel C, de Menezes Reiff RB, Serrão FV. The effect of additional strengthening of hip abductor and lateral rotator muscles in patellofemoral pain syndrome: a randomized controlled pilot study. *Clin Rehabil*. 2008;22(Dec (12)):1051–1060.
19. Herrington L, Al-Sherhi A. A controlled trial of weight-bearing versus non-weight-bearing exercises for patellofemoral pain. *J Orthop Sports Phys Ther*. 2007;37(Apr (4)):155–160.
20. Song CY, Lin YF, Wei TC, Lin DH, Yen TY, Jan MH. Surplus value of hip adduction in leg-press exercise in patients with patellofemoral pain syndrome: a randomized controlled trial. *Phys Ther*. 2009;89(5):409–418.
21. Syme G, Rowe F, Martin D, Daly G. Disability in patients with chronic patellofemoral pain syndrome: a randomised controlled trial of VMO selective training versus general quadriceps strengthening. *Manual Ther*. 2009;14(3):252–263.
22. Brantingham JW, Globe GA, Jensen ML, et al. A feasibility study comparing two chiropractic protocols in the treatment of patellofemoral pain syndrome. *J Manipulative Physiol Ther*. 2009;32(7):536–548.
23. Baldon RM, Serrão FV, Scattone Silva R, Piva SR. Effects of functional stabilization training on pain, function, and lower extremity biomechanics in women with patellofemoral pain: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2014;44(4), 240–A8.
24. Fukuda TY, Rossetto FM, Magalhães E, Bryk FF, Lucareli PR, de Almeida Aparecida Carvalho N. Short-term effects of hip abductors and lateral rotators strengthening in females with patellofemoral pain syndrome: a randomized controlled clinical trial. *J Orthop Sports Phys Ther*. 2010;40(11):736–742.
25. Ismail MM, Gamaleldein MH, Hassa KA. Closed kinetic chain exercises with or without additional hip strengthening exercises in management of patellofemoral pain syndrome: a randomized controlled trial. *Eur J Phys Rehabil Med*. 2013;49(5):687–698.
26. Khayambashi K, Mohammadkhani Z, Ghaznavi K, Lyle MA, Powers CM. The effects of isolated hip abductor and external rotator muscle strengthening on pain, health status, and hip strength in females with patellofemoral pain: a randomized controlled trial. *J Orthop Sports Phys Ther*. 2012;42(1):22–29.
27. Earl JE, Hoch AZ. A proximal strengthening program improves pain, function, and biomechanics in women with patellofemoral pain syndrome. *Am J Sports Med*. 2011;39(1):154–163.
28. Nakagawa TH, Moriya ET, Maciel CD, Serrão FV. Trunk, pelvis, hip, and knee kinematics, hip strength, and gluteal muscle activation during a single-leg squat in males and females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther*. 2012;42(Jun (6)):491–501.
29. Crossley KM, Bennell KL, Cowan SM, Green S. Analysis of outcome measures for persons with patellofemoral pain: which are reliable and valid? *Arch Phys Med Rehabil*. 2004;85:815–822.
30. Magalhães E, Fukuda TY, Sacramento SN, Forgas A, Cohen M, Abdalla RJ. A comparison of hip strength between sedentary females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther*. 2010;40(10):641–647.
31. Martin HJ, Yule V, Syddall HE, Dennison EM, Cooper C, Aihie Sayer A. Is handheld dynamometry useful for the measurement of quadriceps strength in older people? A comparison with the gold standard Biodex dynamometry. *Gerontology*. 2006;52(3):154–159.
32. QUALISYS, AB Qualisys. Track Manger Use Manual. Gothenburg, Suécia, v.1.9.2; 2006.
33. Munro A, Herrington L, Carolan M. Reliability of 2-dimensional video assessment of frontal-plane dynamic knee valgus during common athletic screening tasks. *J Sport Rehabil*. 2012;21(Feb (1)):7–11.
34. Nakagawa TH, Moriya ET, Carlos Dias CD, Serrao FV. Frontal plane biomechanics in males and females with and without patellofemoral pain. *Med Sci Sports Exerc*. 2012;44(Sep (9)):1747–1755.
35. Roy MA, Doherty TJ. Reliability of hand-held dynamometry in assessment of knee extensor strength after hip fracture. *Am J Phys Med Rehabil*. 2004;83(11):813–818.
36. Vasconcelos RA, Grossi DB, Shimano AC, et al. Reliability and validity of a modified isometric dynamometer in the assessment of muscular performance in individuals with anterior cruciate ligament reconstruction. *Rev Bras Ortop*. 2009;44(3).
37. McNemar Q. Note on the sampling error of the difference between correlated proportions or percentages. *Psychometrika*. 1947;12(Jun (2)):153–157.

38. Egan DA, Winchester BJ, Foster C, McGuigan RM. Using session RPE to monitor different methods of resistance exercise. *J Sports Sci Med*. 2006;5(Jun (2)):289–295.
39. Borg G. *Borg's Perceived Exertion and Pain Scales*. Champaign, IL: Human Kinetics; 1998.
40. Noble BJ, Robertson RJ. *Perceived Exertion*. Champaign, IL: Human Kinetics; 1996.
41. Schall R. Estimation in generalized linear models with random effects. *Biometrika*. 1991;78(4):719–727.
42. Mascal CL, Landel R, Powers C. Management of patellofemoral pain targeting hip, pelvis, and trunk muscle function: 2 case reports. *J Orthop Sports Phys Ther*. 2003;33(11):647–660.
43. Schafer JL, Graham JW. Missing data: our view of the state of the art. *Psychol Methods*. 2002;7(Jun (2)):147–177.
44. Boling MC, Bolgla LA, Mattacola CG, Uhl TL, Hosey RG. Outcomes of a weight-bearing rehabilitation program for patients diagnosed with patellofemoral pain syndrome. *Arch Phys Med Rehabil*. 2006;87:1428–1435.
45. Chiu JK, Wong YM, Yung PS, Ng GY. The effects of quadriceps strengthening on pain, function, and patellofemoral joint contact area in persons with patellofemoral pain. *Am J Phys Med Rehabil*. 2012;91(2):98–106.
46. Van der Heijden RA, Lankhorst NE, van Linschoten R, Bierma-Zeinstra SM, van Middelkoop M. Exercise for treating patellofemoral pain syndrome. *Cochrane Database Syst Rev*. 2015;1:CD010387, <http://dx.doi.org/10.1002/14651858.CD010387.pub2>. Review.
47. Van Linschoten R, van Middelkoop M, Berger MY, et al. Supervised exercise therapy versus usual care for patellofemoral pain syndrome: an open label randomised controlled trial. *BMJ*. 2009;339, b4074.
48. Witvrouw E, Lysens R, Bellemans J, Peers K, Vanderstraeten G. Open versus closed kinetic chain exercises for patellofemoral pain. A prospective, randomized study. *Am J Sports Med*. 2000;28(Sep–Oct (5)):687–694.
49. Witvrouw E, Danneels L, Van Tiggelen D, Willems TM, Cambier D. Open versus closed kinetic chain exercises in patellofemoral pain: a 5-year prospective randomized study. *Am J Sports Med*. 2004;32(Jul–Aug (5)):1122–1130.
50. Whyte EF, Moran K, Shortt CP, Marshall B. The influence of reduced hamstring length on patellofemoral joint stress during squatting in healthy male adults. *Gait Posture*. 2010;31(1):47–51.
51. Witvrouw E, Werner S, Mikkelsen C, Van Tiggelen D, Vanden Bergh L, Cerulli G. Clinical classification of patellofemoral pain syndrome: guidelines for nonoperative treatment. *Knee Surg Sports Traumatol Arthrosc*. 2005;13:122–130.
52. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sports Med*. 1993;21(May–Jun (3)):438–444.
53. Rathleff MS, Rathleff CR, Crossley KM, Barton CJ. Is hip strength a risk factor for patellofemoral pain? A systematic review and meta-analysis. *Br J Sports Med*. 2014;48(Jul (14)):1088.
54. Rabelo ND, Lucareli PR. Do hip muscle weakness and dynamic knee valgus matter for the clinical evaluation and decision-making process in patients with patellofemoral pain? *Braz J Phys Ther*. 2017, <http://dx.doi.org/10.1016/j.bjpt.2017.10.002>. pii:S1413-3555(17)30522-1 [Epub ahead of print].
55. Lack S, Barton C, Crossley K, Morrissey D. Proximal muscle rehabilitation is effective for patellofemoral pain: a systematic review with meta-analysis. *Br J Sports Med*. 2015;49(Nov (21)):1365–1376.
56. Santos TRT, Oliveira BA, Ocarino JM, Holt KG, Fonseca ST. Effectiveness of hip muscle strengthening in patellofemoral pain syndrome patients: a systematic review. *Braz J Phys Ther*. 2015;19(May–Jun (3)):167–176.
57. Saad MC, Felício LR, Masullo Cde L, Liporaci RF, Bevilacqua Grossi D. Analysis of the center of pressure displacement, ground reaction force and muscular activity during step exercises. *J Electromyogr Kinesiol*. 2011;21(Oct (5)):712–718.
58. Herman DC, Weinhold PS, Guskiewicz KM, Garrett WE, Yu B, Padua DA. The effects of strength training on the lower extremity biomechanics of female recreational athletes during a stop-jump task. *Am J Sports Med*. 2008;36:733–740.
59. Neumann DA. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther*. 2010;40(Feb (2)):82–94.
60. Oliveira LV, Saad MC, Felício LR, Grossi DB. Muscle strength analysis of hip and knee stabilizers in individuals with Patellofemoral Pain Syndrome. *Fisioter Pesqui*. 2014;21(4), <http://dx.doi.org/10.590/1809-2950/11903021042014>.