Clinician's Commentary on Richards et al.¹

Over the next two decades, the number of people living with stroke in Canada is expected to rise significantly—from 405,000 in 2013 to an estimated 726,000 in 2038—as a result of aging, reduced stroke mortality, and population growth.² Although stroke mortality is decreasing, stroke survivors are living with disabilities;^{2, 3} 40% of them will have moderate to severe impairments,⁴ and as much as 40% of them will have limited to no walking ability.⁵ Thus, the recovery of functional walking capacity is a primary goal of post-stroke survivors as well as neuro-rehabilitation therapists.⁶

The past 3 decades have seen several technological advancements in the rehabilitation of locomotion with the introduction of body weight-supported treadmill training,⁷ robotic devices such as the Lokomat,⁶ and human augmentation through exoskeleton devices.⁸ Simultaneously, virtual reality (VR)—the ability to simulate real-world objects and events—and virtual environments—providing interactive, context-specific environments simulating everyday events—have been developed as rehabilitation tools,⁹ enabling the manipulation of sensory-motor experiences in a safe and progressive environment.

Richards and colleagues¹ have developed a novel rehabilitation tool consisting of a VR-coupled treadmill system that provides the person post-stroke with a training environment that simulates both the visual and the physical demands of a real-life complex environment, simultaneously challenging the sensory-motor and cognitive components of locomotion. The decision to investigate this novel intervention with a person 32 months post-stroke is a welcome addition to the evidence base, which is limited for the population with chronic stroke,¹⁰ and it demonstrates clinical improvements that identify that ongoing recovery is possible in this population. Unfortunately, access to neuro-rehabilitation is scarce for community-dwelling chronic stroke survivors.

This proof-of-principle study highlights the many challenges inherent in neuro-rehabilitation research—in particular, the interface with clinical practice. The research design was creative and thorough, using a factorial case study design. The participant was a community-dwelling independent ambulator with no significant cognitive, perceptual, or communication difficulties or other comorbidities that affected his mobility. This clinical presentation will likely be difficult to replicate with larger numbers of post-stroke participants because approximately 50%–70% of stroke survivors present with cognitive deficits.¹¹ In addition, the expanding aging population in Canada brings clinical complexity because of the increased likelihood of the co-existence of two or more chronic conditions.¹²

A major challenge for all health care providers is managing multi-morbidity in their scope of practice.¹² Likewise, access to and use of expensive, complex technologies requires careful consideration of access to capital and training investments. Understanding the facilitators of, and barriers to, the use of complex technology in clinical practice is essential.¹³ Time is critical in clinical practice, and, therefore, the challenge for the clinician is weighing the training time required to use the technology effectively and safely, and the time required for appropriate set-up, along with the potential benefits and risks of using

the technology.¹⁴ Thus, consideration needs to be given to intervention eligibility (for whom will this intervention be relevant, and in what clinical setting might it be available?) to ensure that newly developed interventions are relevant and accessible to a large proportion of the stroke population during neuro-rehabilitation.

Although Richards and colleagues¹ used a follow-up and washout period, the study participant did not return to baseline clinical measures (except for the Activities-Specific Balance Confidence scale, Canadian French version) after the control intervention. Therefore, the improvements that occurred after the experimental VR intervention may have been, in part, because of the combined intervention effects. This is not unusual in clinical practice and is, in fact, the essence of clinical practice, where interventions are not provided in isolation but rather as a coordinated, progressive combination of interventions that simultaneously address the multiple aspects of the clinical presentation.¹⁵ Herein lies the dichotomy between neuro-rehabilitation from a clinical practice perspective and neuro-rehabilitation research: No one intervention will solve the complex movement, cognitive, and perceptual problems sustained as the result of a stroke, but to date, interventions are generally researched in isolation, and, as a result, the evidence base does not reflect the persons or the complex movement problems encountered every day by neurorehabilitation therapists.

The tools developed over the past three decades, including the one by Richards and colleagues,¹ primarily address one aspect of locomotion rehabilitation—that is, determining the most effective way to improve walking for stroke survivors who are already walking. However, determining the most effective way to potentiate walking for stroke survivors who are not able to walk and who may have a combination of physical, cognitive, and perceptual deficits, remains a fundamental and pressing question for clinicians and stroke survivors alike.

*Julie Vaughan-Graham, PhD, PT
Physiotherapist, Physio-Logic Rehabilitation Services;
Scientific Associate, Toronto Rehab Research Institute;
and Adjunct Lecturer, Department of Physical Therapy,
University of Toronto, Toronto;
julie.vaughan.graham@utoronto.ca.*

REFERENCES

1. Richards CL, Malouin F, Lamontagne AJ, et al. Gait training after stroke on a self-paced treadmill with and without virtual environment scenarios: A proof-of-principle study. *Physiother Can*. 2018;70(3):221–30. <https://doi.org/10.3138/ptc.2016-97>.
2. Krueger H, Koot J, Hall RE, et al. Prevalence of individuals experiencing the effects of stroke in Canada: trends and projections. *Stroke*. 2015;46(8):2226–31. <https://doi.org/10.1161/STROKEAHA.115.009616>. Medline:26205371
3. Edwards JD, Koehoorn M, Boyd LA, et al. Is health-related quality of life improving after stroke? A comparison of health utilities indices

- among Canadians with stroke between 1996 and 2005. *Stroke*. 2010;41(5):996–1000. <https://doi.org/10.1161/STROKEAHA.109.576678>. Medline:20360545
4. Teasell R, Meyer MJ, Foley N, et al. Stroke rehabilitation in Canada: a work in progress. *Top Stroke Rehabil*. 2009;16(1):11–9. <https://doi.org/10.1310/tsr1601-11>. Medline:19443343
 5. Kollen B, Kwakkel G, Lindeman E. Longitudinal robustness of variables predicting independent gait following severe middle cerebral artery stroke: a prospective cohort study. *Clin Rehabil*. 2006;20(3):262–8. <https://doi.org/10.1191/0269215506cr910oa>. Medline:16634346
 6. Calabrò RS, Cacciola A, Bertè F, et al. Robotic gait rehabilitation and substitution devices in neurological disorders: where are we now? *Neurol Sci*. 2016;37(4):503–14. <https://doi.org/10.1007/s10072-016-2474-4>. Medline:26781943
 7. Mehrholz J, Pohl M, Elsner B. Treadmill training and body weight support for walking after stroke. *Cochrane Database Syst Rev*. 2014;(1):CD002840. <https://doi.org/10.1002/14651858.CD002840.pub3>. Medline:24458944
 8. Louie DR, Eng JJ. Powered robotic exoskeletons in post-stroke rehabilitation of gait: a scoping review. *J Neuroeng Rehabil*. 2016;13(1):53. <https://doi.org/10.1186/s12984-016-0162-5>. Medline:27278136
 9. Sangani S, Lamontagne A, Fung J. Cortical mechanisms underlying sensorimotor enhancement promoted by walking with haptic inputs in a virtual environment. In: Dancause N, Nadeau S, Rossignol S, editors. *Progress in brain research*. Vol. 218. New York: Elsevier; 2015. p. 313–30. <https://doi.org/10.1016/bs.pbr.2014.12.003>.
 10. Vaughan-Graham J, Cott C, Wright FV. The Bobath (NDT) concept in adult neurological rehabilitation: what is the state of the knowledge? A scoping review. Part II: intervention studies perspectives. *Disabil Rehabil*. 2015b;37(21):1909–28. <https://doi.org/10.3109/09638288.2014.987880>. Medline:25427891
 11. Mahon S, Parmar P, Barker-Collo S, et al. Determinants, prevalence, and trajectory of long-term post-stroke cognitive impairment: results from a 4-year follow-up of the ARCOS-IV study. *Neuroepidemiology*. 2017;49(3-4):129–34. <https://doi.org/10.1159/000484606>. Medline:29145207
 12. Upshur RE, Tracy S. Chronicity and complexity: is what's good for the diseases always good for the patients? *Can Fam Physician*. 2008;54(12):1655–8. Medline:19074692
 13. Schmid L, Glässel A, Schuster-Amft C. Therapists' perspective on virtual reality training in patients after stroke: a qualitative study reporting focus group results from three hospitals. *Stroke Res Treat*. 2016;2016:6210508. <https://doi.org/10.1155/2016/6210508>. Medline:28058130
 14. Glegg SM, Holsti L, Velikonja D, et al. Factors influencing therapists' adoption of virtual reality for brain injury rehabilitation. *Cyberpsychol Behav Soc Netw*. 2013;16(5):385–401. <https://doi.org/10.1089/cyber.2013.1506>. Medline:23713844
 15. Cott CA, Graham JV, Brunton K. When will the evidence catch up with clinical practice? *Physiother Can*. 2011;63(3):387–90. <https://doi.org/10.3138/physio.63.3.387>. Medline:22654245

DOI:10.3138/ptc.2016-97-cc