Echocardiographic Evaluation of Aortic Stenosis – Normal Flow and Low Flow Scenarios

Ian G Burwash

University of Ottawa Heart Institute, University of Ottawa, Ottawa, Ontario, Canada

Abstract

The echocardiographic evaluation of the patient with aortic stenosis (AS) has evolved in recent years, beyond confirming the diagnosis and measuring the resting mean pressure gradient or valve area. New echocardiographic approaches have developed to address the clinical dilemmas related to discordant haemodynamic data, asymptomatic haemodynamically severe AS and low-flow, low-gradient AS in order to better evaluate the disease severity, enhance the risk stratification of patients and provide important prognostic information. This article reviews the echocardiographic evaluation of the AS patient and focuses on the echocardiographic assessment of the haemodynamic severity, the prediction of clinical outcome and the use of echocardiography to guide patient management in the presence of normal flow and low flow scenarios.

Keywords

Aortic stenosis, echocardiography, stress echocardiography, exercise echocardiography, dobutamine echocardiography, low-flow aortic stenosis, low-gradient aortic stenosis

Disclosure: The author has no conflicts of interest to declare.

Received: 10 October 2014 **Accepted:** 11 November 2014 **Citation:** *European Cardiology Review*, 2014;9(2):92–9

Correspondence: Ian G Burwash, Department of Medicine, University of Ottawa Heart Institute, 40 Ruskin Street, Ottawa, Ontario, K1Y 4W7, Canada
E: iburwash@ottawaheart.ca

Aortic stenosis (AS) is a progressive disease of worsening left ventricular outflow obstruction. In the early stage of the disease, the prognosis is excellent with a very small risk of sudden death (<1 % per year).^{1–3} However, the development of a haemodynamically severe stenosis and the onset of symptoms herald a dramatic deterioration in prognosis. At this point, aortic valve replacement (AVR) provides symptomatic and survival benefits that generally outweigh the associated surgical risks.^{4–10}

Echocardiography plays a critical role in the management of patients with suspected and documented AS by confirming the diagnosis, quantifying the haemodynamic severity and providing prognostic information important for predicting the clinical course and guiding management.^{9–11} Resting transthoracic echocardiography provides the mainstay technique for these purposes; however, transoesophageal and stress echocardiography can provide additional important information when encountering patients with discordant haemodynamic data, apparently asymptomatic haemodynamically severe AS and severe low-flow, low-gradient AS with or without a reduced left ventricular ejection fraction (LVEF). The purpose of this article is to review the echocardiographic assessment of the AS patient, highlighting important issues related to the assessment of haemodynamic severity, the prediction of clinical outcome and the use of echocardiography to guide patient management in both normal flow and low flow scenarios.

Criterion for Haemodynamically Severe Aortic Stenosis

Management of the AS patient is critically dependent on an accurate evaluation of the patient's symptomatic status and the haemodynamic

severity of the valve disease.^{9,10} Haemodynamic severity is used to decide whether symptoms are likely attributable to the valve and dictate the frequency of follow-up evaluations. Numerous haemodynamic indices have been proposed for this purpose; however, transvalvular velocity (V_{AS}), transvalvular mean gradient (MG) and aortic valve area (AVA) derived by the continuity equation are the parameters recommended for routine clinical use due to their ability to predict clinical outcome.^{9–11} Haemodynamically severe AS has been defined by $V_{AS} \geq 4$ m/s, $MG \geq 40$ mmHg and $AVA \leq 1.0$ cm², because these findings are associated with a poor prognosis.^{9–11} The fulfilment of all three criteria essentially confirms the presence of severe AS; however, discordant findings occur in at least 30 % of AS patients with a normal LVEF, casting doubt about the disease severity.^{12–14} In patients with a reduced LVEF, discordant findings are even more common.^{15–19} In both groups, the vast majority of the discordance relates to an AVA in the severe range, but MG or V_{AS} in the non-severe range. In only a small minority of cases (<10 %), AVA is in the non-severe range, but MG or V_{AS} is severe.^{12,14}

Sorting Out Discrepant Haemodynamic Data – Aortic Valve Area is Severe, but the Mean Gradient is Not!

The Normal Left Ventricular Ejection Fraction Patient Technical Reasons (Measurement Error)

Accurate measurement of the left ventricular outflow tract (LVOT) velocity, LVOT diameter and V_{AS} are critical to the calculation of AVA using the continuity equation. Underestimation of the LVOT time velocity integral or diameter can lead to an erroneously small

stroke volume (SV) and underestimation of AVA. Optimal LVOT velocity signals require the ultrasound beam to be aligned parallel to left ventricular (LV) outflow, the sample volume to be positioned immediately inferior to the aortic valve (5–10 mm) and the signal to show laminar flow with minimal spectral dispersion.¹¹ The LVOT diameter measurement should be acquired on a zoomed parasternal long-axis image in mid-systole and measured 5 mm below cusp insertion.¹¹ While measurements obtained at the annulus improve reproducibility, they tend to result in a larger AVA. The circular assumption used to derive the LVOT area from the diameter measurement is an important cause for underestimation of AVA as many patients have an elliptically shaped LVOT.²⁰ The LVOT long-axis diameter often approximates the minor axis diameter and can lead to a 26 % underestimation of SV and AVA.^{21,22} Identification of the highest V_{AS} is required to accurately measure MG and AVA, and Doppler interrogation of the aortic valve should be performed from the apical, right parasternal and suprasternal windows.²³ Failure to identify the highest V_{AS} will lead to an underestimation of MG, an overestimation of AVA and an underestimation of AS severity that tends to have a concordant rather than discordant MG and AVA.

When discrepant data are encountered, the initial approach should be to rule out measurement error. Echocardiographic images should be carefully reviewed for proper data acquisition and repeated if improper technique was used. The SV measurement should be confirmed by using another technique such as three-dimensional (3D)-echocardiography, or corroborating the continuity equation AVA by planimetry of the anatomic orifice using two-dimensional (2D) or preferably 3D-transoesophageal echocardiography.^{22,24} An $AVA \leq 1.0 \text{ cm}^2$ on planimetry confirms the presence of severe AS. However, an AVA slightly greater than 1.0 cm^2 does not definitively exclude severe AS since the effective orifice area derived by continuity equation is usually slightly smaller than the anatomic AVA.

Patient Body Size

Cardiac output and AVA requirements are dependent on body size. In a small individual, an $AVA \leq 1.0 \text{ cm}^2$ may be relatively adequate for the cardiac output and generate a $V_{AS} < 4 \text{ m/s}$ or $MG < 40 \text{ mmHg}$. Conversely, an $AVA > 1.0 \text{ cm}^2$ may be inadequate for a large individual, resulting in severe obstruction and a $V_{AS} \geq 4 \text{ m/s}$ or $MG \geq 40 \text{ mmHg}$. Current guidelines suggest that indexing AVA for body surface area (aortic valve area index [AVAi]) to account for different body sizes may be helpful and propose an $AVAi \leq 0.6 \text{ cm}^2/\text{m}^2$ to identify severe AS.^{9–11} However, prognostic studies supporting this approach are limited.²⁵ Indexing AVA to body surface area increases the prevalence of severe AS, but also increases the prevalence of discrepant haemodynamic data.^{12,14,26,27} In the Simvastatin and Ezetimibe in Aortic Stenosis (SEAS) trial, AVAi did not improve the prediction of aortic valve events or cardiovascular death compared to AVA.^{26,27} A linear relationship between AVA and body surface area may not be the optimal correction factor and likely oversimplifies the different implications of body size, overweight status and obesity that might require different correction factors.^{14,28} Nevertheless, body size should be considered when confronted with discordant haemodynamic data.

Inconsistencies Between Different Criteria for Severe Aortic Stenosis in the Valve Guidelines

Inherent in the definition of severe AS is that a MG of 40 mmHg equates to an AVA of 1.0 cm^2 . However, the relationship between

MG and AVA is a function of cardiac output, heart rate and systolic ejection period. A patient with a cardiac output of 6 l/min, a heart rate of 70 bpm and a systolic ejection period of 330 msec would have a MG of 34 mmHg if AVA was 1.0 cm^2 , not 40 mmHg as suggested by the guidelines. Potentially, MG could be as low as 26 mmHg with an AVA of 1.0 cm^2 despite a normal cardiac output.²⁹

Low Flow State

MG is directly proportional to the squared function of transvalvular flow and can be small ($< 40 \text{ mmHg}$) despite an $AVA \leq 1.0 \text{ cm}^2$ if transvalvular flow is significantly reduced.^{16,18,19} Low flow is usually considered when the LVEF is reduced, but should not be excluded when the LVEF is normal. Low flow has been defined as an indexed stroke volume (SV_i) $< 35 \text{ ml/m}^2$;^{19,30} however, measures that account for the systolic ejection period (i.e. mean transvalvular flow rate $< 200\text{--}250 \text{ ml/s}$) are theoretically likely more appropriate. Hypertension, significant mitral or tricuspid valve disease, pulmonary artery hypertension and atrial fibrillation are clinical entities often co-existing with AS and associated with low flow despite a normal LVEF. Elevated blood pressure can reduce transvalvular flow and result in a smaller MG, smaller AVA and discordant measurements.³¹ In patients with uncontrolled hypertension, low flow and discordant data, blood pressure should be corrected and the echocardiogram repeated as normalisation of flow may resolve the discrepancy.

Low transvalvular flow can also be present in AS patients without concomitant valve disease or pulmonary artery hypertension, a clinical entity often referred to as paradoxical low flow AS, because of the paradoxical association of a low flow state despite a preserved LVEF.³⁰ These patients tend to be older, female and have associated hypertension, metabolic syndrome, diabetes or atrial fibrillation.^{19,30,32–34} They also tend to have smaller left ventricular cavities with concentric remodeling, myocardial fibrosis, reduced compliance and restrictive physiology.^{19,34,35} Despite the preserved LVEF, they have evidence of intrinsic systolic dysfunction with impaired longitudinal shortening.^{34–37}

The Reduced Left Ventricular Ejection Fraction Patient

The presence of a low flow state is often the first assumption when encountering discrepant haemodynamic data in the AS patient with a reduced LVEF. However, not all patients with a reduced LVEF have a low flow state. SV_i or mean transvalvular flow rate ($< 35 \text{ ml/m}^2$ and $< 200\text{--}250 \text{ ml/s}$, respectively) should be calculated to confirm the presence of low flow and measurement error excluded.

Additional Complementary Investigations

Discordant measurements can usually be resolved by systematically considering the above issues. However, multi-detector computed tomography (MDCT), magnetic resonance imaging (MRI) and cardiac catheterisation can be helpful when the cause for the discrepancy is not apparent and the patient has symptoms or unexplained left ventricular dysfunction. AVA, LVOT area and SV can be obtained non-invasively using MDCT or MRI and compared with the echocardiography data.^{22,38–40} Aortic valve calcification (AVC) can be quantified using MDCT and may help clarify discordant data. $AVC > \sim 1,250$ Agatston units (AU) in women and $> \sim 2,050$ AU in men suggests the presence of severe AS.⁴¹ Cardiac catheterisation provides a direct measurement of MG or AVA by Gorlin equation, and can evaluate for coronary artery disease as an alternate explanation for symptoms or left ventricular dysfunction.³⁹

Figure 1: Survival of Medically Managed (Censored at Aortic Valve Replacement) Severe Aortic Stenosis Patients with Left Ventricular Ejection Fraction ≥ 50 % (n=1,704) Based on the Flow Pressure Gradient Classification

The NF/LG group had the best survival and the LF/LG group had the worst ($p < 0.0001$).
 AVR = aortic valve replacement; HG = high gradient; LF = low flow; LG = low gradient;
 NF = normal flow. Source: Eleid et al., 2013,³² reproduced with permission from the publisher (Wolters Kluwer Health).

Risk Stratification and the Prediction of Outcome Using Echocardiography

Asymptomatic Aortic Stenosis with Normal Left Ventricular Ejection Fraction

Peak Aortic Stenosis Velocity and the Rate of Haemodynamic Progression

V_{AS} is the strongest predictor of clinical outcome, even in the subgroup with severe AS.^{1,2,42-45} For asymptomatic AS patients with $V_{AS} < 3$ m/s, 3–4 m/s and > 4 m/s, approximately 15 %, 35 % and 75 % of patients, respectively, will develop symptoms requiring AVR within three years.^{1-3,42,45} $V_{AS} > 5.5$ m/s is associated with a poor three year event-free survival of only 11 %, and the European Society of Cardiology (ESC) and American Heart Association (AHA)/American College of Cardiology (ACC) valve guidelines consider AVR reasonable in these asymptomatic patients if the surgical risk is low (IIa recommendation).^{9,10} AVA also predicts event-free survival, although the relationship is not as robust as V_{AS} .^{1,42,43,46} This may relate in part to an inherent bias by which clinicians may be more likely to refer patients with a higher V_{AS} or MG for AVR. Rapid haemodynamic progression, defined as an increase in $V_{AS} \geq 0.3$ m/s/yr, identifies patients at increased risk for clinical events.^{1,3,45} Severe AS patients with rapid haemodynamic progression have an event rate as high as 80 % in two years,³ and according to the ESC and AHA/ACC valve guidelines, AVR may be considered in an asymptomatic patient at low surgical risk (IIa and IIb recommendation, respectively).^{9,10}

Valve Calcification

Valve calcification is best evaluated clinically using MDCT. However, valve calcification assessed by echocardiography is a strong predictor of haemodynamic and clinical progression across the spectrum of disease severity, and independent of the haemodynamic severity.^{3,45} Calcification can be graded as none, mild (isolated small spots), moderate (multiple bigger spots) and severe (extensive thickening/calcification of all cusps). Severe AS patients with moderate or severe valve calcification have an event rate of 80 % at four years, but only 25 % with no or mild calcification.³

Flow Pressure Gradient Classification

Patients with $AVA \leq 1.0$ cm² and a normal LVEF can be classified into four subtypes based on the presence of:

- normal flow (NF) or low flow (LF) ($SVi \geq$ or < 35 ml/m²); and
- high MG (HG) or low MG (LG) ($MG \geq$ or < 40 mmHg).^{36,37}

Most patients (52–73 %) have the NF/HG pattern, in which MG and AVA are concordant for severe AS and SVi is normal.^{30,32,36,37} The NF/LG pattern is present in 15–31 %, and encompasses patients with inherent inconsistencies in the cut-points for severe AS, small body size and prolonged ejection times in which SVi is normal but transvalvular flow rate may be reduced. The LF/HG pattern is observed in 3–13 %, and includes patients with concordant indices but intrinsic left ventricular dysfunction despite a normal LVEF. The LF/LG pattern is present in 3–19 % of patients, a group characterised by discordant haemodynamic indices and a low flow state.

The flow pressure gradient classification provides important prognostic information for severe AS patients with a normal LVEF. The NF/LG pattern has the highest event-free survival, while the LF/LG pattern has the worst outcome.^{32,34,37,47} The NF/HG and LF/HG patterns have intermediate survival. In Eleid et al.,³² survival with medical management (censored at AVR) at two years was 82 % in NF/LG patients, 81 % in LF/HG patients, 67 % in NF/HG patients and 44 % in LF/LG patients ($p < 0.0001$) (see Figure 1). AVR conferred a strong survival benefit in LF/LG and NF/HG patients, but no benefit in NF/LG patients. However, other investigators have observed a survival benefit of AVR in both symptomatic LF/LG and NF/LG patients.⁴⁸⁻⁵¹ SVi appears to be an important marker of prognosis and has a graded association with mortality in severe low gradient AS patients with preserved LVEF.^{52,53}

Left Ventricular Function

Only a minority of severe AS patients present with a reduced LVEF and the prognosis is poor without AVR.^{9,54} However, a large proportion of severe AS patients with normal LVEF have intrinsic systolic dysfunction with abnormal global longitudinal strain on speckle tracking imaging.⁵⁵⁻⁵⁷ Patients with moderate or severe AS and global longitudinal strain ≤ 15 % have a poor two-year event-free survival of ≈ 30 %, as compared with 65 % if > 15 %.^{56,57} There are no data to justify AVR in asymptomatic patients with impaired longitudinal strain, although close follow-up is likely warranted.

Valvular Arterial Impedance

Valvular arterial impedance (Zva) provides a measure of global LV load due to the valve stenosis and vascular afterload. Zva is calculated as $(SBP + MG) / SVi$, where SBP is the systolic blood pressure.⁵⁸ In asymptomatic patients with moderate or severe AS, $Zva > 4.5$ –5.0 mmHg/ml/m² predicts reduced event-free survival and a 2.8-fold increased risk of death;^{56,58} however, Zva does not identify the relative contributions of the valve and vasculature that is necessary to guide management (i.e. AVR versus blood pressure [BP] control).

Exercise Echocardiography

More than 50 % of patients with asymptomatic severe AS develop symptoms within three years, and the subsequent prognosis is poor without timely AVR.^{1-3,8} Exercise testing can unmask symptoms in as many as one third of patients and a positive exercise test predicts the need for AVR or death.⁵⁹⁻⁶¹ The ESC and AHA/ACC valve guidelines

consider AVR indicated in asymptomatic severe AS patients with symptoms on exercise testing (Class I recommendation) or a drop in BP (IIa recommendation).^{9,10}

Exercise echocardiography may provide additional prognostic information beyond the exercise test. An exercise increase in $MG > 18$ – 20 mmHg potentially identifies severe AS patients at increased risk for symptoms, AVR or death, even with a normal exercise test (see Figure 2).^{62,63} The adverse prognosis of patients with greater increases in MG likely reflects a more severely stenotic or non-compliant valve with minimal opening reserve. In the ESC guidelines, AVR may be considered in these patients if the surgical risk is low (IIb recommendation).¹⁰ Failure of LVEF to increase with exercise is also associated with an abnormal exercise test and reduced event-free survival (event-free survival $< 40\%$ at two years).⁶⁴ Although not specifically addressed in either valve guidelines, these patients should at a minimum be observed closely for surgical indications. Resting and exercise changes in global longitudinal strain are associated with an abnormal stress test and reduced event-free survival; however, their incremental prognostic value requires further study.^{55,65}

Pulmonary hypertension is associated with poor outcomes and higher complication rates with valve intervention.^{66,67} The prevalence of pulmonary hypertension at rest in asymptomatic AS patients is low; however, exercise-induced pulmonary hypertension may develop in $> 50\%$ of severe AS patients.⁶⁸ Pulmonary systolic pressure > 60 mmHg with exercise is associated with a twofold increased risk of cardiac events and a reduced cardiac event-free survival at three years (22 ± 7 versus $55 \pm 9\%$; $p = 0.014$), independent of the change in MG with exercise.⁶⁸ Exercise pulmonary hypertension appears to unmask a severe impairment of diastolic function. Close follow-up of these patients is likely warranted.

Low-flow, Low-gradient Aortic Stenosis

Low-flow, Low-gradient Severe Aortic Stenosis with Reduced Left Ventricular Ejection Fraction

Definition

Low-flow, low-gradient severe AS with reduced LVEF, or classical low-flow, low-gradient AS,¹⁹ is defined by a $MG < 40$ mmHg or $V_{AS} < 4$ m/s, $AVA \leq 1.0$ cm², and a $LVEF < 40$ – 50% .^{9–11} Inherent in the definition is that flow is reduced, although no specific criteria are included in the guidelines. Individual studies have used a cardiac index < 3.0 l/min/m² or $SVi < 35$ ml/m²,^{18,19} although measures such as transvalvular flow rate (< 200 – 250 ml/s) may be more appropriate. These patients represent a diagnostic and management challenge since the severity of the underlying valve stenosis is often unclear, AVR is associated with a high operative risk and the benefits of AVR to alleviate symptoms or significantly improve prognosis may be limited in some patients.^{16–19,54,69–71}

True Versus Pseudo-severe Aortic Stenosis

AVA, whether measured by continuity equation or orifice planimetry, is affected by flow and becomes smaller when transvalvular flow is reduced.^{72–74} This phenomenon has been attributed to flow dependence of the flow convergence,^{75–77} flow-mediated effects on the vena contracta spatial velocity profile,⁷⁶ reductions in anatomic orifice area due to smaller flow-mediated forces,^{78,79} or a combination of these factors. The potential effect is that AVA may appear only mild or moderately stenotic when assessed under normal flow conditions, but potentially severe when assessed under low flow conditions. The challenge when

Figure 2: Event-free Survival (Cardiovascular Death or Aortic Valve Replacement) of Aortic Stenosis Patients with a Normal Exercise Test (n=135) Based on the Rest Mean Gradient and Exercise Increase in Mean Gradient

AVR = aortic valve replacement; Exer. MG = exercise-induced increase in mean pressure gradient; MG = mean pressure gradient. Source: Marechaux et al., 2010,⁶³ reproduced with permission from Oxford University Press.

encountering the patient with low-flow, low-gradient severe AS with reduced LVEF is to distinguish the patient with 'true' severe AS (TSAS), where the valve is severely stenotic and afterload mismatch exists, from the patient with 'pseudo-severe' AS (PSAS), where the underlying valve is only mild or moderately stenotic, but appears severe due to low flow and the inability of a poorly functioning ventricle to sufficiently open the valve. AVR is likely beneficial in the TSAS patient, but may not be beneficial in the PSAS patient since the valve is not the primary problem. PSAS is reported to be present in 5–35% of these patients depending on the specific diagnostic criteria used.^{69,70,80–82}

TSAS and PSAS can be distinguished by augmenting transvalvular flow using dobutamine stress echocardiography (DSE) and re-evaluating the valve haemodynamics at a normal flow state. A commonly used DSE protocol employs an initial infusion of 2.5–5.0 µg/kg/min, followed by incremental infusions of 2.5–5.0 µg/kg/min at 5–8 minute stages to a maximum dose of 20 µg/kg/min.^{71,82} This low dose, longer stage protocol minimises the potential for myocardial ischaemia and allows for steady state acquisition of LVOT and transvalvular velocities. Patients with TSAS and an underlying 'fixed' valve stenosis would be expected to have little change in AVA and a large increase in MG during DSE. In contrast, patients with PSAS would be expected to have an increase in AVA outside of the severe range and only a small increase in MG due to the larger AVA.

Criteria used to diagnose TSAS have included a DSE peak $MG > 30$ – 40 mmHg, an AVA increase < 0.3 cm², and/or a DSE peak $AVA \leq 1.0$ – 1.2 cm².^{18,19,69,70,80–82} PSAS is suggested if the DSE peak AVA is > 1.0 – 1.2 cm² and the peak MG is ≤ 30 mmHg or the AVA increase is ≥ 0.3 cm². In the AHA/ACC valve guidelines, AVR is considered reasonable in symptomatic patients with low-flow, low-gradient severe AS with reduced LVEF if V_{AS} is ≥ 4.0 m/s or MG is ≥ 40 mmHg with an $AVA \leq 1.0$ cm² at any time during DSE (IIa recommendation).⁹ While this criteria is specific for TSAS, many

Figure 3: Survival Curves for Low-flow, Low-gradient Aortic Stenosis Patients with Reduced Left Ventricular Ejection Fraction (n=136) and Contractile Reserve (Group I) or No Contractile Reserve (Group II) Based on Whether They Underwent Valve Replacement or Medical Treatment

Source: Monin et al., 2003,⁷⁰ reproduced with permission from Wolters Kluwer Health.

patients with TSAS have a $MG < 40$ mmHg during DSE because of poor contractile reserve or inherent inconsistencies in the criteria for severe AS.^{69,82,83} A lower cut-point for MG (i.e. >30 mmHg) may be justified to avoid underdiagnosing TSAS patients in whom the prognosis without AVR is extremely poor.^{69-71,82,83}

A significant limitation to distinguishing TSAS and PSAS patients relates to the marked inter-individual variability to which flow augmentation occurs with DSE.^{74,82} Some patients have very little increase in transvalvular flow during DSE and a low flow state persists. Other patients have a large increase in transvalvular flow with supranormal flow rates. Since the increase in MG and AVA are largely dependent on the magnitude of flow augmentation, interpretation of the underlying AS severity without consideration of the transvalvular flow is problematic. To address this issue, the projected AVA (AVAproj) at 250 ml/s has been proposed to standardise the calculation of AVA at a normal flow rate and better distinguish TSAS and PSAS.^{82,83} Transvalvular flow rate and AVA generally display a linear relationship in AS patients with the slope varying between patients.^{72,74} AVAproj can be calculated by measuring AVA and transvalvular flow rate at a minimum of two different flow rates during DSE using the equation:

$$AVA_{proj} = AVA_{rest} + VC \times (250 - Q_{rest})$$

where AVA_{rest} is the resting AVA, Q_{rest} is the resting mean transvalvular flow rate and VC is the 'valve compliance' or slope of the valve area – transvalvular flow relationship.⁸²

In an initial 23 patients undergoing surgical inspection, $AVA_{proj} \leq 1.0$ cm² and an $AVA_{proj} \leq 0.55$ cm²/m² correctly classified 83 % and 91 % of patients, respectively.⁸² In a larger study of 52 patients, a simplified approach was used to calculate VC that only required the measurement of resting and peak DSE transvalvular flow rate and AVA:

$$VC = (AVA_{peak} - AVA_{rest}) / (Q_{peak} - Q_{rest})$$

$AVA_{proj} \leq 1.0$ cm² correctly classified 92 % of patients, and was superior to DSE peak AVA < 1.2 cm² (77 %), peak MG > 30 mmHg (73 %) and AVA increase < 0.3 cm² (60 %).⁸³ However, AVAproj needs to be interpreted cautiously if the increase in transvalvular flow rate is < 15 % due to inaccuracies in calculating VC.⁸²

Figure 4: Survival Curves for Low-flow, Low-gradient Aortic Stenosis Patients with Reduced Left Ventricular Ejection Fraction Based on (A) Dobutamine Stress Echocardiography Peak Left Ventricular Ejection Fraction ≤ 35 or > 35 % (n=101) and (B) Dobutamine Stress Echocardiography Peak Global Longitudinal Strain Rate $\leq -0.5s^{-1}$ or $> -0.5s^{-1}$ (n=47)

DSE = dobutamine stress echocardiography; LVEF = left ventricular ejection fraction; PLSR = peak global longitudinal strain rate. Source for Figure 4A: Clavel et al., 2008,⁷¹ reproduced with permission from Wolters Kluwer Health. Source for Figure 4B: Bartko et al. 2013,⁸⁶ reproduced with permission from Wolters Kluwer Health.*

Quantitation of AVC by MDCT may provide an additional technique to potentially distinguish TSAS from PSAS, and be especially helpful in those patients in whom the DSE is non-diagnostic or the augmentation of transvalvular flow is minimal. In 18 patients with classical low-flow, low-gradient AS, $AVC \geq 1,651$ AU correctly classified 89 % of patients as having either TSAS or PSAS.⁸⁴ Gender specific thresholds for identifying TSAS are likely warranted based on studies in AS patients with preserved LVEF ($\sim 1,250$ AU in woman and $\sim 2,050$ AU in men), although this has not been studied in classical low-flow, low-gradient AS.⁴¹

Prediction of Clinical Outcome

Contractile or flow reserve, defined as a 20 % increase in SV with DSE, is a strong predictor of peri-operative mortality rate and long-term survival.^{69,70} AVR is associated with a peri-operative mortality rate as high as 33 % in patients with no contractile reserve, but only 5–8 % in patients with contractile reserve.^{54,69,70} In the European Multicentre study, three-year survival of patients with contractile

reserve undergoing AVR was 79 %, significantly better than the 35 % survival observed in patients with no contractile reserve (see Figure 3).⁷⁰ Patients with contractile reserve did significantly better with AVR than medical management (three-year survival <25 %).

Patients with no contractile reserve have a high operative mortality rate and poor long-term survival; however, their survival is better with AVR than medical therapy when tested using propensity matched analysis.⁵⁴ Furthermore, patients with no contractile reserve have similar improvements in LVEF and New York Heart Association (NYHA) functional class following AVR as patients with contractile reserve.⁸⁵ In essence, absence of contractile reserve identifies patients with a high operative mortality rate and reduced long-term survival who may still benefit from AVR. In the ESC guidelines, AVR should be considered in symptomatic patients with evidence of flow reserve (IIa recommendation) and may be considered in symptomatic patients without flow reserve (IIb recommendation).¹⁰

In the True or Pseudo Severe Aortic Stenosis (TOPAS) study of 101 patients with low-flow, low-gradient severe AS with reduced LVEF, independent echocardiographic predictors of reduced survival were $AVA_{proj} \leq 1.2 \text{ cm}^2$ and DSE peak LVEF $\leq 35 \%$ (see Figure 4A).^{71,83} Contractile reserve was not an independent predictor of survival.⁷¹ The conflicting findings regarding the predictive value of contractile reserve in TOPAS and the European Multicentre Study may have related to different DSE protocols or differences in the study population (TOPAS had a greater prevalence of concomitant diseases such as coronary artery disease).^{70,71} Additionally, DSE peak LVEF is essentially a composite measure of both resting LVEF and 'contractile reserve', and may explain the better predictive value of this parameter.

Longitudinal strain parameters are severely impaired in low-flow, low-gradient severe AS with reduced LVEF, but may improve with DSE in a substantial proportion of patients. In a substudy of TOPAS, patients with a DSE peak global longitudinal strain rate (PLSR) $< -0.5 \text{ s}^{-1}$ had a two-year survival of 91 %, but only 40 % if DSE PLSR was $\geq -0.5 \text{ s}^{-1}$ (see Figure 4B).⁸⁶ DSE PLSR was the strongest predictor of overall survival and added incremental prognostic information to DSE peak LVEF.

Low-flow, Low-gradient Severe Aortic Stenosis with Preserved Left Ventricular Ejection Fraction

Definition

The criteria for low-flow, low-gradient severe AS with preserved LVEF include a $V_{AS} < 4 \text{ m/s}$ or $MG < 40 \text{ mmHg}$, $AVA \leq 1.0 \text{ cm}^2$, $LVEF > 50 \%$ and $SVi < 35 \text{ ml/m}^2$.^{9-11,19}

True Versus Pseudo-severe Aortic Stenosis

Patients with low-flow, low-gradient severe AS with preserved LVEF have similar transvalvular flow rates as classical low-flow, low-gradient severe AS.^{83,87} Since transvalvular flow is reduced, the small AVA may reflect underlying TSAS or PSAS. The prevalence of PSAS is reported to be $\approx 30 \%$ in these patients, similar to what is observed in classical low-flow, low-gradient severe AS.⁸⁷ DSE and exercise echocardiography have been used to distinguish TSAS and PSAS in one small study.⁸⁷ $AVA_{proj} \leq 1.0 \text{ cm}^2$ was 94 % accurate in distinguishing the two conditions when compared to surgical inspection, and performed better than DSE peak $MG \geq 30 \text{ mmHg}$ (75 %) or $\geq 40 \text{ mmHg}$ (81 %) and DSE peak $AVA \leq 1.0 \text{ cm}^2$ (69 %) or $\leq 1.2 \text{ cm}^2$ (63 %). In the AHA/ACC and ESC valve guidelines, AVR is reasonable in symptomatic patients if the clinical, haemodynamic and

Figure 5: Event-free Survival of Low-flow, Low-gradient Severe Aortic Stenosis Patients with Left Ventricular Ejection Fraction >50 % (n=55) Based on (A) AVA_{proj} and (B) Indexed AVA_{proj}

AVA_{proj} = projected aortic valve area; HR = hazard ratio. * on Figure 5B is hazard ratio (HR) adjusted for age and sex. Source: Clavel et al., 2013,⁸⁷ reproduced with permission from Elsevier.

anatomic data support valve obstruction as the most likely cause of the symptoms (IIa recommendation).^{9,10} Reserving surgical intervention only for symptomatic TSAS patients appears theoretically sound, although this management strategy has not been tested.

Measurements of AVC using MDCT may have a role in identifying TSAS and PSAS in patients with low-flow, low-gradient severe AS with preserved LVEF. In 451 patients with at least moderate AS, preserved LVEF, normal SVi and concordant MG and AVA measurements, an AVC threshold of 1,274 AU in women and 2,065 AU in men correctly classified 87 % of patients as having moderate AS or severe AS.⁴¹ A similar 87 % accuracy was obtained using an AVC indexed to aortic annulus area threshold of 292 AU/cm² for women and 476 AU/cm² for men. In a further 172 patients with preserved LVEF and discordant AS measurements ($MG < 40 \text{ mmHg}$ and $AVA \leq 0.6 \text{ cm}^2/\text{m}^2$), $\approx 50 \%$ of patients had an AVC score consistent with moderate rather than severe AS.⁴¹ Low flow ($SVi < 35 \text{ ml/m}^2$) was present in $\approx 16 \%$ of these patients; however, results for this subgroup were not provided and DSE results were not reported. Further studies are warranted to determine whether AVC thresholds can be used to distinguish TSAS and PSAS in low-flow, low-gradient severe AS with preserved LVEF.

Prediction of Clinical Outcome

AVAproj and indexed AVAproj are strong predictors of adverse outcome in patients with low-flow, low-gradient severe AS with preserved LVEF (hazard ratio: 1.32 and 1.59, respectively, for each 0.1 cm² decrease in AVAproj and indexed AVAproj) and is superior to traditional parameters of stenosis severity for predicting outcome.⁸⁷ More than 90 % of patients with AVAproj>1.0 cm² or indexed AVAproj>0.55 cm²/m² remain free of death or AVR at 12 months, whereas only 50 % of patients with AVAproj≤1.0 cm² or AVAproj≤0.55 cm²/m² will be free of death or AVR (see *Figure 5*).⁸⁷

Conclusion

The role of echocardiography in the evaluation of AS patients has progressed beyond the measurement of resting MG and AVA. Challenging situations in which patients have discordant haemodynamic data, asymptomatic haemodynamically severe AS or severe low-flow, low-gradient AS have resulted in new approaches to improve the haemodynamic evaluation, risk stratification and prediction of clinical outcome for patients with both normal-flow and low-flow AS. Integration of these advances will help ensure that AS patients receive the most appropriate management. ■

- Otto CM, Burwash IG, Leggett ME, et al., Prospective study of asymptomatic valvular aortic stenosis. Clinical, echocardiographic, and exercise predictors of outcome, *Circulation*, 1997;95:2262–70.
- Pellikka PA, Sarano ME, Nishimura RA, et al., Outcome of 622 adults with asymptomatic, hemodynamically significant aortic stenosis during prolonged follow-up, *Circulation*, 2005;111:3290–5.
- Rosenhek R, Binder T, Porenta G, et al., Predictors of outcome in severe, asymptomatic aortic stenosis, *N Engl J Med*, 2000;343:611–7.
- Ross J Jr, Braunwald E, Aortic stenosis, *Circulation*, 1968;38:61–7.
- Lund O, Preoperative risk evaluation and stratification of long-term survival after valve replacement for aortic stenosis: reasons for earlier operative intervention, *Circulation*, 1990;82:124–39.
- Kvidal P, Bergström R, Hörte LG, Ståhle E, Observed and relative survival after aortic valve replacement, *J Am Coll Cardiol*, 2000;35:747–56.
- Schwarz F, Baumann P, Manthey J, et al., The effect of aortic valve replacement on survival, *Circulation*, 1982;66:1105–10.
- Leon MB, Smith CR, Mack M, et al., Transcatheter aortic-valve implantation for aortic stenosis in patients who cannot undergo surgery, *N Engl J Med*, 2010;363:1597–607.
- Nishimura RA, Otto CM, Bonow RO, et al., 2014 AHA/ACC Guideline for the Management of Patients With Valvular Heart Disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. ACC/AHA Task Force Members, *Circulation*, 2014;129:e521–643.
- Vahanian A, Alfieri O, Andreotti F, et al., Guidelines on the management of valvular heart disease (version 2012), *Eur Heart J*, 2012;33:2451–96.
- Baumgartner H, Hung J, Bermejo J, et al., Echocardiographic assessment of valve stenosis: EAE/ASE recommendations for clinical practice, *J Am Soc Echocardiogr*, 2009;22:1–23.
- Minners J, Allgeier M, Gohlke-Baerwolf C, et al., Inconsistencies of echocardiographic criteria for the grading of aortic valve stenosis, *Eur Heart J*, 2008;29:1043–8.
- Jander N, Minners J, Holme I, et al., Outcome of patients with low-gradient “severe” aortic stenosis and preserved ejection fraction, *Circulation*, 2011;123:887–95.
- Michelena HI, Margaryan E, Miller FA, et al., Inconsistent echocardiographic grading of aortic stenosis: is the left ventricular outflow tract important?, *Heart*, 2013;99:921–31.
- Connolly HM, Oh JK, Orszulik TA, et al., Aortic valve replacement for aortic stenosis with severe left ventricular dysfunction. Prognostic indicators, *Circulation*, 1997;95:2395–400.
- Connolly HM, Oh JK, Schaff HV, et al., Severe aortic stenosis with low transvalvular gradient and severe left ventricular dysfunction: result of aortic valve replacement in 52 patients, *Circulation*, 2000;101:1940–6.
- Kulik A, Burwash IG, Kapila V, et al., Long-term outcomes after valve replacement for low-gradient aortic stenosis: Impact of prosthesis-patient mismatch, *Circulation*, 2006;114:553–8.
- Burwash IG, Low flow, low gradient aortic stenosis: From evaluation to treatment, *Curr Opin Cardiol*, 2007;22:84–91.
- Pibarot P, Dumesnil JG, Low-flow, low-gradient aortic stenosis with normal and depressed left ventricular ejection fraction, *J Am Coll Cardiol*, 2012;60:1845–53.
- Doddamani S1, Grushko MJ, Makaryus AN, et al., Demonstration of left ventricular outflow tract eccentricity by 64-slice multi-detector CT, *Int J Cardiovasc Imaging*, 2009;25:175–81.
- Piazza N, de Jaegere P, Schultz C, et al., Anatomy of the aortic valvar complex and its implications for transcatheter implantation of the aortic valve, *Circ Cardiovasc Interv*, 2008;1:74–81.
- Ng AC, Delgado V, van der Kley F, et al., Comparison of aortic root dimensions and geometries before and after transcatheter aortic valve implantation by 2- and 3 dimensional transoesophageal echocardiography and multislice computed tomography, *Circ Cardiovasc Imaging*, 2010;3:94–102.
- Aggarwal SG, Huq NM, Chan KL, et al., A practical model for Doppler assessment of aortic stenosis severity: Accuracy of a limited window approach, *J Am Soc Echocardiogr*, 1999;12:353.
- Nakai H, Takeuchi M, Yoshitani H, et al., Pitfalls of anatomical aortic valve area measurements using two-dimensional transoesophageal echocardiography and the potential of three-dimensional transoesophageal echocardiography, *Eur J Echocardiogr*, 2010;11:369–76.
- Saito T, Muro T, Takeda H, et al., Prognostic value of aortic valve area index in asymptomatic patients with severe aortic stenosis, *Am J Cardiol*, 2012;110:93–7.
- Jander N, Gohlke-Baerwolf C, Bahlmann E, et al., Indexing aortic valve area by body surface area increases the prevalence of severe aortic stenosis, *Heart*, 2014;100:28–33.
- Minners J, Gohlke-Baerwolf C, Kaufmann BA, et al., Adjusting parameters of aortic valve stenosis severity by body size, *Heart*, 2014;100:1024–30.
- Neelan TG, Pradhan AD, Weyman AE, Derivation of a size-independent variable for scaling of cardiac dimensions in a normal adult population, *J Am Soc Echocardiogr*, 2008;21:779–85.
- Carabello BA, Clinical practice. Aortic stenosis, *N Engl J Med*, 2002;346:677–82.
- Hachicha Z, Dumesnil JG, Bogaty P, Pibarot P, Paradoxical low-flow, low-gradient severe aortic stenosis despite preserved ejection fraction is associated with higher afterload and reduced survival, *Circulation*, 2007;115:2856–64.
- Little SH, Chan KL, Burwash IG, Impact of blood pressure on the Doppler echocardiographic assessment of severity of aortic stenosis, *Heart*, 2007;93:848–55.
- Eleid MF, Sorajja P, Michelena HI, et al., Flow-gradient patterns in severe aortic stenosis with preserved ejection fraction: clinical characteristics and predictors of survival, *Circulation*, 2013;128:1781–9.
- Pibarot P, Dumesnil JG, Aortic stenosis suspected to be severe despite low gradients, *Circ Cardiovasc Imaging*, 2014;7:545–51.
- Mehrotra P, Jansen K, Flynn AW, et al., Differential left ventricular remodeling and longitudinal function distinguishes low flow from normal-flow preserved ejection fraction low-gradient severe aortic stenosis, *Eur Heart J*, 2013;34:1906–14.
- Herrmann S, Störk S, Niemann M, et al., Low-gradient aortic valve stenosis myocardial fibrosis and its influence on function and outcome, *J Am Coll Cardiol*, 2011;58:402–12.
- Adda J, Mielot C, Giorgi R, et al., Low-flow, low-gradient severe aortic stenosis despite normal ejection fraction is associated with severe left ventricular dysfunction as assessed by speckle-tracking echocardiography: a multicenter study, *Circ Cardiovasc Imaging*, 2012;5:27–35.
- Lancellotti P, Magne J, Donal E, et al., Clinical outcome in asymptomatic severe aortic stenosis. Insights from the new proposed aortic stenosis grading classification, *J Am Coll Cardiol*, 2012;59:235–43.
- Shah RG, Novaro GM, Blandon RJ, et al., Aortic valve area: meta-analysis of diagnostic performance of multi-detector computed tomography for aortic valve area measurements as compared to transthoracic echocardiography, *Int J Cardiovasc Imaging*, 2009;25:601–9.
- Saikhrihan N1, Kumar G, Sawaya FJ, et al., Accurate assessment of aortic stenosis: a review of diagnostic modalities and hemodynamics, *Circulation*, 2014;129:244–53.
- Chin CW, Khaw HJ, Luo E, et al., Echocardiography underestimates stroke volume and aortic valve area: implications for patients with small-area low-gradient aortic stenosis, *Can J Cardiol*, 2014;30:1064–72.
- Clavel MA, Messika-Zeitoun D, Pibarot P, et al., The complex nature of discordant severe calcified aortic valve disease grading: new insights from combined Doppler echocardiographic and computed tomographic study, *J Am Coll Cardiol*, 2013;62:2329–38.
- Monin JL, Lancellotti P, Monchi M, et al., Risk score for predicting outcome in patients with asymptomatic aortic stenosis, *Circulation*, 2009;120:69–75.
- Stewart RA, Kerr AJ, Whalley GA, et al., New Zealand Heart Valve Study Investigators. Left ventricular systolic and diastolic function assessed by tissue Doppler imaging and outcome in asymptomatic aortic stenosis, *Eur Heart J*, 2010;31:2216–22.
- Rosenhek R, Zilberszac R, Schemper M, et al., Natural history of very severe aortic stenosis, *Circulation*, 2010;121:151–6.
- Rosenhek R, Klaar U, Schemper M, et al., Mild and moderate aortic stenosis. Natural history and risk stratification by echocardiography, *Eur Heart J*, 2004;25:199–205.
- Malouf J, Le Tourneau T, Pellikka P, et al., Aortic valve stenosis in community medical practice: determinants of outcome and implications for aortic valve replacement, *J Thorac Cardiovasc Surg*, 2012;144:1421–7.
- Clavel MA, Dumesnil JG, Capoulade R, et al., Outcome of patients with aortic stenosis, small valve area, and low-flow, low-gradient despite preserved left ventricular ejection fraction, *J Am Coll Cardiol*, 2012;60:1259–67.
- Ozkan A, Hachamovitch R, Kapadia SR, et al., Impact of aortic valve replacement on outcome of symptomatic patients with severe aortic stenosis with low gradient and preserved left ventricular ejection fraction, *Circulation*, 2013;128:622–31.
- Dumesnil JG, Pibarot P, Carabello B, Paradoxical low flow and/or low gradient severe aortic stenosis despite preserved left ventricular ejection fraction: implication for diagnosis and treatment, *Eur Heart J*, 2010;31:281–9.
- Tarantini G, Covolo E, Razzolini R, et al., Valve replacement for severe aortic stenosis with low transvalvular gradient and left ventricular ejection fraction exceeding 0.50, *Ann Thorac Surg*, 2011;91:1808–15.
- Maor E, Beigel R, Maor E, et al., Survival after intervention in patients with low gradient severe aortic stenosis and preserved left ventricular function, *J Thorac Cardiovasc Surg*, 2014 [Epub ahead of print].
- Eleid MF, Sorajja P, Michelena HI, et al., Survival by stroke volume index in patients with low-gradient normal EF severe aortic stenosis, *Heart*, 2014 [Epub ahead of print].
- Maor E, Beigel R, Grupper A, et al., Relation between stroke volume index to risk of death in patients with low-gradient severe aortic stenosis and preserved left ventricular function, *Am J Cardiol*, 2014;114:449–55.
- Tribouilloy C, Lévy F, Rusinaru D, et al., Outcome after aortic valve replacement for low-flow/low-gradient aortic stenosis without contractile reserve on dobutamine stress echocardiography, *J Am Coll Cardiol*, 2009;53:1865–73.
- Lafitte S, Perlant M, Reant P, et al., Impact of impaired myocardial deformations on exercise tolerance and prognosis in patients with asymptomatic aortic stenosis, *Eur J Echocardiogr*, 2009;10:414–9.
- Lancellotti P, Donal E, Magne J, et al., Risk stratification in asymptomatic moderate to severe aortic stenosis: the importance of the valvular, arterial and ventricular interplay, *Heart*, 2010;96:1364–71.
- Yingchoncharoen T, Gibby C, Rodriguez LL, et al., Association of myocardial deformation with outcome in asymptomatic aortic stenosis with normal ejection fraction, *Circ Cardiovasc Imaging*, 2012;5:719–25.
- Pibarot P, Dumesnil JG, Improving assessment of aortic stenosis, *J Am Coll Cardiol*, 2012;60:169–80.
- Amato MC, Moffa PJ, Werner KE, Ramirez JA, Treatment decision in asymptomatic aortic valve stenosis: role of exercise testing, *Heart*, 2001;86:381–6.
- Das P, Rimgington H, Chambers J, Exercise testing to stratify risk in aortic stenosis, *Eur Heart J*, 2005;26:1309–13.
- Magne J, Lancellotti P, Piérard LA, Exercise testing in asymptomatic severe aortic stenosis, *JACC Cardiovasc Imaging*, 2014;7:188–99.
- Lancellotti P, Lebois F, Simon M, et al., Prognostic importance of quantitative exercise Doppler echocardiography in asymptomatic valvular aortic stenosis, *Circulation*, 2005;112(9 Suppl):1377–82.
- Marechaux S, Hachicha Z, Bellouin A, et al., Usefulness of exercise-stress echocardiography for risk stratification of true asymptomatic patients with aortic valve stenosis, *Eur Heart J*, 2010;31:1390–7.
- Marechaux S, Ennezat PV, LeJemtel TH, et al., Left ventricular response to exercise in aortic stenosis: an exercise echocardiographic study, *Echocardiography*, 2007;24:955–9.
- Donal E, Thebault C, O'Connor K, et al., Impact of aortic stenosis on longitudinal myocardial deformation during exercise, *Eur J Echocardiogr*, 2011;12:235–41.
- McHenry MM, Rice J, Matlof HJ, Flamm MD Jr, Pulmonary hypertension and sudden death in aortic stenosis, *Br Heart J*, 1979;41:463–7.
- Carnero-Alcázar M, Reguillo-Lacruz F, Alsuiw A, et al., Short- and mid-term results for aortic valve replacement in octogenarians, *Interact Cardiovasc Thorac Surg*, 2010;10:549–54.
- Lancellotti P, Magne J, Donal E, et al., Determinants and prognostic significance of exercise pulmonary hypertension in asymptomatic severe aortic stenosis, *Circulation*, 2012;126:851–9.
- Nishimura RA, Grantham A, Connolly HM, et al., Low-output, low-gradient aortic stenosis in patients with depressed left ventricular systolic function. The clinical utility of the

- dobutamine challenge in the catheterization laboratory, *Circulation*, 2002;106:809–13.
70. Monin JL, Quéré JP, Monchi M, et al., Low-gradient aortic stenosis: operative risk stratification and predictors for long-term outcome: a multicenter study using dobutamine stress hemodynamics, *Circulation*, 2003;108:319–24.
71. Clavel MA, Fuchs C, Burwash IG, et al., Predictors of outcomes in low-flow, low gradient aortic stenosis: results of the multicenter TOPAS Study, *Circulation*, 2008;118(14suppl):S234–42.
72. Burwash IG, Thomas DD, Sadahiro M, et al., Dependence of Gorlin formula and continuity equation valve areas on transvalvular volume flow rate in valvular aortic stenosis, *Circulation*, 1994;89:827–35.
73. Burwash IG, Pearlman AS, Kraft CD, et al., Flow dependence of measures of aortic stenosis severity during exercise, *J Am Coll Cardiol*, 1994;24:1342–50.
74. Burwash IG, Hay KM, Chan KL, Hemodynamic stability of valve area, valve resistance, and stroke work loss in aortic stenosis: a comparative analysis, *J Am Soc Echocardiogr*, 2002;15:814–22.
75. Segal J, Lerner DJ, Miller C, et al., When should Doppler determined valve area be better than the Gorlin formula? Variation in hydraulic constants in low flow states, *J Am Coll Cardiol*, 1987;9:1294–305.
76. DeGroff CG, Shandas R, Valdes-Cruz L, Analysis of the effect of flow rate on the Doppler continuity equation for stenotic orifice area calculations: a numerical study, *Circulation*, 1998;97:1597–605.
77. Kadem L, Rieu R, Dumesnil JG, et al., Flow-dependent changes in Doppler derived aortic valve effective orifice area are real and not due to artifact, *J Am Coll Cardiol*, 2006;47:131–7.
78. Montarello JK, Perakis AC, Rosenthal E, et al., Normal and stenotic human aortic valve opening: in vitro assessment of orifice area changes with flow, *Eur Heart J*, 1990;11:484–91.
79. Chambers JB, Spriggs DC, Cochrane T, et al., Continuity equation and Gorlin formula compared with directly observed orifice area in native and prosthetic aortic valves, *Br Heart J*, 1992;67:193–9.
80. deFilippi CR, Willett DL, Brickner ME, et al., Usefulness of dobutamine echocardiography in distinguishing severe from nonsevere valvular aortic stenosis in patients with depressed left ventricular function and low transvalvular gradients, *Am J Cardiol*, 1995;75:191–4.
81. Schwammenthal E, Vered Z, Moshkowitz Y, et al., Dobutamine echocardiography in patients with aortic stenosis and left ventricular dysfunction: predicting outcome as a function of management strategy, *Chest*, 2001;119:1766–77.
82. Blais C, Burwash IG, Mundigler G, et al., Projected valve area at normal flow rate improves the assessment of stenosis severity in patients with low-flow, low-gradient aortic stenosis: the multicenter TOPAS (Truly or Pseudo-Severe Aortic Stenosis) study, *Circulation*, 2006;113:711–21.
83. Clavel MA, Burwash IG, Mundigler G, et al., Validation of conventional and simplified methods to calculate projected valve area at normal flow rate in patients with low flow, low gradient aortic stenosis: the multicenter TOPAS (True or Pseudo Severe Aortic Stenosis) study, *J Am Soc Echocardiogr*, 2010;23:380–6.
84. Cuffe C, Serfaty JM, Cimadevilla C, et al., Measurement of aortic valve calcification using multislice computed tomography: correlation with haemodynamic severity of aortic stenosis and clinical implication for patients with low ejection fraction, *Heart*, 2011;97:721–6.
85. Quere JP, Monin JL, Levy F, et al., Influence of preoperative left ventricular contractile reserve on postoperative ejection fraction in low-gradient aortic stenosis, *Circulation*, 2006;113:1738–44.
86. Bartko PE, Heinze G, Graf S, et al., Two-dimensional strain for the assessment of left ventricular function in low flow-low gradient aortic stenosis, relationship to hemodynamics, and outcome: a substudy of the multicenter TOPAS study, *Circ Cardiovasc Imaging*, 2013;6:268–76.
87. Clavel MA, Ennezat PV, Maréchaux S, et al., Stress echocardiography to assess stenosis severity and predict outcome in patients with paradoxical low-flow, low-gradient aortic stenosis and preserved LVEF, *JACC Cardiovasc Imaging*, 2013;6:175–83.