HHS Public Access

Author manuscript

Arthritis Rheumatol. Author manuscript; available in PMC 2019 October 01.

Published in final edited form as:

Arthritis Rheumatol. 2018 October ; 70(10): 1694. doi:10.1002/art.40530.

Leonine facies presenting as scleromyxedema

Annie Y. Park, MD¹, Lori Lowe, MD², and Dinesh Khanna, MD, MS³

¹Department of Internal Medicine, University of Michigan, Ann Arbor, MI

²Department of Dermatology and Pathology, University of Michigan, Ann Arbor, MI

³University of Michigan Scleroderma Program, University of Michigan, Ann Arbor, MI

References

1. Blum M, Wigley FM, Hummers LK. Scleromyxedema: A Case Series Highlighting Long-Term Outcomes of Treatment With Intravenous Immunoglobulin (IVIG). *Medicine (Baltimore)*. 2008; 87(1):10.doi: 10.1097/MD.0b013e3181630835 [PubMed: 18204366]

Corresponding Author: Dinesh Khanna, MD MS, Professor of Medicine, Scleroderma Program, Division of Rheumatology, Department of Internal Medicine, University of Michigan, 300 North Ingalls Building, Room 7C27, Ann Arbor, MI 48109-5422, Phone 734-936-5561/Fax 734-763-1253, khannad@med.umich.edu.

Financial Disclosures:

Annie Park: No financial disclosures

Lori Lowe: No financial disclosures

Dinesh Khanna: Dr. Khanna was supported by NIH/NIAMS K24 AR063121, and NIH/NIAMS R01 AR070470.

Figure 1.

A 61-year-old man previously in good health was referred to Scleroderma Clinic for a 2-year history of slowly progressive cutaneous eruption involving his dorsal hands, extremities, and central face. On physical examination, the patient had nodular, erythematous indurated lesions of his forehead and erythematous papular lesions on his nose with coalescence of firm erythematous papulonodules, resulting in a leonine facies. On the dorsal hands, arms, and legs were numerous, shiny, firm, closely set, 1–2 mm slightly translucent papules with background erythema (face and hand, shown in Panel A). Skin biopsy demonstrated a spindled fibroblastic proliferation in the dermis with increased mucin and variable fibrosis (Panel B). The clinical and histologic findings are diagnostic of scleromyxedema. Scleromyxedema is a rare disorder of unknown pathogenesis characterized by a generalized lichenoid papular cutaneous eruption resulting in diffuse skin induration that may simulate scleroderma. Rarely, larger exophytic nodules, as seen in this patient, may be present. The vast majority of cases of scleromyxedema occur in association with a monoclonal gammopathy. Our patient was found to have an IgG lambda M-protein spike. The patient did not exhibit any CRAB features (hypercalcemia, renal insufficiency, anemia, and bone lesions) and evaluation with hematology culminated in diagnosis of IgG lambda monogammopathy of unclear significance (MGUS), with plans for ongoing observation. For his scleromyxedema, the patient received intravenous immunoglobulin (IVIG) ¹ at doses up to 2g/kg/month, with significant improvement in appearance of his lesions after 16 months (Panel C). He continues to receive a maintenance dose of 1 gram/kg IVIG every 4 weeks.