Errata

The January 10, 2013, article by Greaves et al, entitled “Expression of FOXP3, CD68, and CD20 at Diagnosis in the Microenvironment of Classical Hodgkin Lymphoma Is Predictive of Outcome” (J Clin Oncol 31:256-262), was published with errors in the Figure 2 parts C and D labels.

In the line color key for both 2C and 2D, the labels are inverted; Low should be High, and High should be Low.

The key order was given as (top to bottom): Low (blue line), Intermediate (yellow line), High (gray line)

Instead, the key should read as (top to bottom): **High** (blue line), Intermediate (yellow line), **Low** (gray line)

The online version has been corrected in departure from the print. *Journal of Clinical Oncology* regrets the error.

DOI: 10.1200/JCO.2017.72.8329; published April 1, 2017

The July 10, 2016, article by Denduluri et al, entitled “Selection of Optimal Adjuvant Chemotherapy Regimens for Human Epidermal Growth Factor Receptor 2 (HER2)-Negative and Adjuvant Targeted Therapy for HER2-Positive Breast Cancers: An American Society of Clinical Oncology Guideline Adaptation of the Cancer Care Ontario Clinical Practice Guideline” (J Clin Oncol 34:2416-2427, 2016), contained errors.

Under Recommendations in the Bottom Line section, based on the findings of the trial conducted by the Gruppo Italiano Mammella investigators, which demonstrated improved disease-free survival and overall survival observed with four cycles of epirubicin 90 mg/m² and cyclophosphamide 600 mg/m² followed by four cycles of paclitaxel 175 mg/m² delivered every 2 weeks (dose-dense EC-P) compared to every 3 weeks, the guideline adaptation should have included the dose-dense EC-P regimen, as provided below, in the list of acceptable adjuvant chemotherapy regimens for patients with higher-risk early breast cancer.

This guideline adaptation should also have been provided in two additional places: under the panel lists “Acceptable Adjuvant Chemotherapy Regimens for Patients With Higher-Risk Early Breast Cancer” and in Table 1 in the row

“Acceptable adjuvant chemotherapy regimens for patients with higher-risk early breast cancer (CCO recommendation 13).” In this table, the text in the right column should now read **Not adapted**.

Dose-dense EC-P = Epirubicin 90mg/m², cyclophosphamide 600mg/m² every 2 weeks 4 cycles followed by paclitaxel 175mg/m² every 2 weeks for 4 cycles.

The regimen **Epirubicin 90mg/m², cyclophosphamide 600mg/m² every 2 weeks 4 cycles followed by paclitaxel 175mg/m² every 2 weeks for 4 cycles** should have been provided under “Acceptable Adjuvant Chemotherapy Regimens for Patients With Higher-Risk Early Breast Cancer.” The sentence “The ASCO Panel modified the CCO list of acceptable adjuvant chemotherapy regimens to remove the “dose-dense, dose-intense epirubicin-cyclophosphamide → paclitaxel” regimen because of higher toxicity and lack of overall survival benefit” under “ASCO Panel rationale and discussion” is now incorrect. The paragraph for “ASCO Panel rationale and discussion” should be removed.

The online version has been corrected in departure from the print. The authors apologize for the errors.

DOI: 10.1200/JCO.2017.72.7198; published April 1, 2017

The December 20, 2016 article by Hofmeister et al, entitled “How to Integrate Elotuzumab and Daratumumab Into Therapy for Multiple Myeloma” (J Clin Oncol 34:4421-4430, 2016), contained errors.

In Figure 1, the infusion-related AEs, as described by Lokhorst et al, for only part 1 of the trial were incorrectly listed as 63% for grades 1-2 and 6% for grades 3-4. The correct values for both parts 1 and 2 of the trial are and **65%** for grade 1-2 and **3%** for grade 3-4 infusion reactions in patients exposed to both 8 mg/kg & 16 mg/kg daratumumab

Of all patients treated at 8mg/kg and 16 mg/kg as described by Lonial et al, the any grade proportion was 63% and

the grade 3-4 proportion was 4%. The grade 1-2 proportion was given as 63% in the figure, but should have been **59%**.

The Palumbo citation in Figure 1 was given as 41, but should be reference **42**.

The last sentence in Figure 1 should have been updated. It was given as, “In the reports of daratumumab by Lokhorst et al⁴⁵ and Lonial et al,^{45a} grade 1 to 2 reactions occurred in the majority of patients, predominately sinorespiratory events; other reactions included bronchospasm, congestion, throat irritation, and cough.”

However, the sentence should have read, “In the reports of daratumumab **from all doses infused** by Lokhorst et al⁴⁵ and

Lonial et al,^{45a} grade 1 to 2 reactions occurred in the majority of patients, predominately sinorespiratory events; other reactions included bronchospasm, congestion, throat irritation, and cough.”

The online version has been corrected in departure from the print. The authors apologize for the errors.

DOI: 10.1200/JCO.2017.72.7586; published April 1, 2017

The February 1, 2017, article by Knight et al, entitled “Group-Wide, Prospective Study of Ototoxicity Assessment in Children Receiving Cisplatin Chemotherapy (ACCL05C1): A Report From the Children’s Oncology Group” (J Clin Oncol 35: 440-445, 2017), was published with an error in the abstract.

Instead, this sentence should read as: One central reviewer also used the **International Society of Pediatric Oncology** Ototoxicity Scale (SIOP).

The online version has been corrected in departure from the print. The authors regret the error.

DOI: 10.1200/JCO.2017.72.7685; published April 1, 2017

The February 20, 2017, article by Finelli et al entitled “Management of Small Renal Masses: American Society of Clinical Oncology Clinical Practice Guideline” (J Clin Oncol 35:668-680) was published with an error.

The online version has been corrected in departure from the print. The authors apologize for the error.

In the author list, Houston R. Thompson should have been **R. Houston Thompson**.

DOI: 10.1200/JCO.2017.72.7867; published April 1, 2017