Published in final edited form as:

Curr Hypertens Rep. 2017 March ; 19(3): 24. doi:10.1007/s11906-017-0724-3.

Defining the relationship between hypertension, cognitive decline, and dementia: a review

Keenan A. Walker¹, Melinda C. Power², and Rebecca F. Gottesman^{1,3}

¹Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

²Department of Epidemiology and Biostatistics, George Washington University Milken Institute School of Public Health, Washington, DC, USA

³Department of Epidemiology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

Abstract

Hypertension is a highly prevalent condition which has been established as a risk factor for cardiovascular and cerebrovascular disease. Although the understanding of the relationship between cardiocirculatory dysfunction and brain health has improved significantly over the last several decades, it is still unclear whether hypertension constitutes a potentially treatable risk factor for cognitive decline and dementia. While it is clear that hypertension can affect brain structure and function, recent findings suggest that the associations between blood pressure and brain health are complex and, in many cases, dependent on factors such as age, hypertension chronicity, and antihypertensive medication use. Whereas large epidemiological studies have demonstrated a consistent association between high midlife BP and late-life cognitive decline and incident dementia, associations between late-life blood pressure and cognition have been less consistent. Recent evidence suggests that hypertension may promote alterations in brain structure and function through a process of cerebral vessel remodeling, which can lead to disruptions in cerebral autoregulation, reductions in cerebral perfusion, and limit the brain's ability to clear potentially harmful proteins such as β -amyloid. The purpose of the current review is to synthesize recent findings from epidemiological, neuroimaging, physiological, genetic, and translational research to provide an overview of what is currently known about the association between blood pressure and cognitive function across the lifespan. In doing so, the current review also discusses the results of recent randomized controlled trials of antihypertensive therapy to reduce cognitive decline, highlights several methodological limitations, and provides recommendations for future clinical trial design.

Corresponding Author: Rebecca F. Gottesman, rgottesm@jhmi.edu, Department of Neurology, Johns Hopkins Hospital, Phipps 446D, 600 North Wolfe St., Baltimore, MD 21287, T: 410-614-2381, F: 410-955-0672.

Conflict of Interest

Drs. Walker, Power, and Gottesman declare no conflict of interest relevant to this manuscript

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

Papers of particular interest have been highlighted as:

* Of importance

** Of great importance

Keywords

Hypertension; Hypotension; Blood Pressure; Cognition; Cognitive Impairment; Dementia

Introduction

Hypertension is a highly prevalent condition, occurring in one-third of the world's adults and two-thirds of adults over age 65 [1,2]. Already an established risk factor for cardiovascular and cerebrovascular disease [3–6], emerging evidence suggests that hypertension may also play an important role in the development of cognitive decline, Alzheimer's disease, and vascular dementia [7–9]. Because hypertension is a modifiable risk factor, it represents a potentially important mechanism through which the prevention or delay of age-related cognitive disorders may be possible. For this reason, understanding hypertension's role in the development and progression of age-related cognitive decline and dementia has been a research priority over the last two decades. Although a great deal has been learned from epidemiological studies, there is still little consensus about the effectiveness of treating hypertension to prevent or slow cognitive decline. What is clear, however, is that the connection between blood pressure (BP) and cognitive function is biologically complex and still not fully understood.

The goal of this review is to provide an overview of the research that has contributed to the understanding of the connection between BP and cognitive function, paying particular attention to recent findings. In doing so, this review will first provide an overview of what is known about the connection between hypertension, cognitive function, Alzheimer's disease, and vascular dementia. Second, the neurobiological changes associated with hypertension will be described, and the research that demonstrates how these biological processes influence neuronal function will be highlighted. Lastly, the findings from clinical trials designed to assess the effectiveness of antihypertensive agents for the prevention or delay of cognitive decline will be summarized. Methodological considerations and specific recommendations for future research will also be discussed. Although this review focuses on the topic of hypertension and cognitive function, the link between low BP and cognition will also be discussed.

Hypertension and cognitive function

Cross-sectional and longitudinal observational studies

Over the last several decades the link between hypertension and cognitive function has been examined across many age groups. Although much of this research has focused on understanding the relationship between BP and cognition in older adults, the group most likely to experience cognitive decline, studies which assess BP starting in middle-age and follow participants forward until they reach older ages have also been especially informative. Multiple epidemiological studies have demonstrated that elevated BP in the 4th and 5th decade of life, particularly untreated hypertension, increases the risk for cognitive impairment 20–30 years later (see Table 1) [10–12]. These findings have been further supported by longitudinal studies which show that high midlife BP is associated with

increased cognitive decline over time [13–15]. Because confounding variables, such as education and socioeconomic status, are less likely to affect cognitive change (compared to baseline cognitive abilities) [16], studies which show an increased rate of cognitive decline over time among hypertensive adults provide especially strong evidence for the deleterious effects of high BP. As will be discussed below, several studies have also identified hypertension duration and the trajectory of BP levels over time as important determinants of cognitive function later in life [17,18].

Hypertension in the 6th and 7th decade has been associated with poorer overall cognitive function and cognitive decline (see Table 2) [19–23]. Hypertension among individuals in their 70s has also been identified as a risk factor for mild cognitive impairment (MCI) – a state of subtle cognitive decline that is believed to precede the onset of dementia [24,25]. In contrast, studies that include individuals in their 8th, 9th, and 10th decade of life have largely either failed to find such an association [26,27] or have found high BP to be protective against cognitive impairment [28,29]. Together, these results suggest that the relationship between cognition and BP in late-life may be age dependent [30]. Inverted U- or J-shaped curves may most accurately represent the relationship between BP and cognition among octogenarians and nonagenarians, as both low BP and extremely high BP (systolic blood pressure (SBP) >160mmHg) have been linked to cognitive impairment in this age group [28,31–33].

While individuals who develop hypertension earlier in life are likely to be subjected to the deleterious neurological effects of hypertension for many decades, this is not the case for individuals who develop hypertension much later. The strong associations found between midlife hypertension and late-life cognitive abilities supports the notion that hypertension duration and chronicity in adulthood may be especially important determinants of cognitive impairment in elderly individuals. Perhaps the strongest support for this hypothesis comes from a longitudinal study which found that a longer duration of time between hypertension initiation and cognitive testing is associated with reduced cognitive abilities independent of age [18]. In particular, longitudinal studies suggest that middle-aged adults with prolonged hypertension and elevated systolic blood pressure (SBP) over a period of 25–30 years are at an exceptionally high risk for cognitive impairment later in life [17,18]. Thus, studies with a longer period between the initiation of BP monitoring and subsequent cognitive assessment may be better able to detect the effects of high BP on neurocognitive outcomes. The trajectory of blood pressure changes from midlife into older age may also be important, as the combination of hypertension in midlife and low diastolic blood pressure (DBP) in late-life has been associated with smaller brain volumes and poorer cognitive outcomes among older adults [34]. Individuals who develop hypertension before middle adulthood may also be at particularly high risk for cognitive impairment, as a number of studies have found associations between high BP, cognitive deficits, and reduced academic functioning in children, adolescents, and young adults [35–40]. Irrespective of age, the cognitive domains that appear most vulnerable to hypertension are executive functioning and information processing speed. Both cognitive processes rely heavily on the integrity of frontal and subcortical brain structures which may be most vulnerable to the effects of hypertension.

Blood pressure variability

BP fluctuates substantially over a 24-hour period as a result of factors such as postural change, circadian rhythm, and general physiologic variability [41,42]. Fluctuations in BP associated with autonomic dysfunction, such as orthostatic hypotension, become more prevalent with increasing age and may be associated with cognitive deficits [41,43,44]. Although a number of studies have demonstrated a connection between orthostatic hypotension and cognitive function, with worse performance in the setting of orthostasis [45–48], others have failed to replicate this finding [49–51]. Ambulatory blood pressure measurement (ABPM) has been used in a number of studies to more accurately capture short-term, daily BP variability, which may reflect autonomic dysfunction or increased arterial stiffness, among other etiologies. Using ABPM, elevated 24-hour mean BP, 24-hour BP variability, and reduced nocturnal dipping (a natural reduction of night-time BP) have each been identified as potential risk factors for cognitive impairment [11,52–54]. Because autonomic dysfunction occurs in the early phase of several neurodegenerative disorders [55], it is difficult to determine whether cognitive deficits found in individuals with potential sequelae of autonomic dysfunction (e.g., BP variability and orthostatic hypotension) are the result of underlying neurodegenerative changes or the direct effect of transient drops in BP.

Genetic factors

Additional insights into the relationship between hypertension and cognition have emerged through genetic studies. A polymorphism in the *ACE* gene, a gene which regulates BP through its effects on angiotensin converting enzyme (ACE) activity [56], has been linked to both cognitive function [57] and the presence of neuroimaging abnormalities [58,59]. Middle-aged and older adults who carry an allele that codes for the high activity variant (D) of the ACE I/D polymorphism show greater levels of cognitive impairment and cognitive decline [57,60–62]. Unexpectedly, other studies have found the low activity allele (I) of the ACE I/D polymorphism to confer increased risk for dementia [63,64]. Polymorphisms in another gene, *AGTR1*, which codes for the angiotensin-II type 1 receptor, also an important part of the regulation of BP, have been associated with reduced prefrontal and hippocampal volume [65], reductions in hippocampal volume over time, and poorer memory in older adults [66]. Additional evidence suggests that specific genetic variants may interact with hypertension to promote or buffer against the effects of elevated BP on cognitive function and brain structural integrity. Two Alzheimer's disease risk genes that have also been associated with cognitive function in nondemented individuals, *Apolipoprotein E* (APOE) and *Clusterin* (CLU), appear to modify the effect of hypertension on cognitive function [67]. For example, multiple studies have found that hypertension is only associated with cognitive deficits in individuals who possess a copy of the $\epsilon 4$ allele of the APOE gene [68,69].

Dementia risk and hypertension

Alzheimer's disease

Several forms of cardiovascular disease have been identified as risk factors for both Alzheimer's disease and vascular dementia [70–73], which together account for the majority of dementia cases worldwide [74,75]. Alzheimer's disease, cerebrovascular disease, and cardiovascular disease have shared genetic contributions [76,77], and approximately 50% of

individuals diagnosed with Alzheimer's disease display significant cerebrovascular pathology on autopsy [78,79]. Together, these findings suggest that cardiovascular disease, Alzheimer's disease, and vascular dementia may have an overlapping pathophysiology [80–82].

Despite significant evidence for the role of cardiovascular disease in the pathogenesis and progression of Alzheimer's disease, the association between hypertension and Alzheimer's disease is still not well understood. Although a consistent relationship between elevated DBP at midlife and incident Alzheimer's disease has been demonstrated [7,83,84], evidence for an association between midlife SBP and incident Alzheimer's disease has been conflicting [84–88]. What is clear is that late-life hypertension does not appear to be a risk factor for incident Alzheimer's disease [72,88–93]. In fact, multiple studies suggest that abnormally low DBP in late-life may increase one's risk for Alzheimer's disease [91,94–98]. Some, but not all, have argued that this inverse relationship between late-life DBP and Alzheimer's disease risk results from a tendency for BP to decline concurrently with the onset and progression of dementia [90,99,100]. Together, previous findings suggest that the combination of high BP in midlife followed by low BP in late-life may place individuals at especially high risk of developing Alzheimer's disease. However, few studies have examined this hypothesis directly [101].

Vascular dementia

Because hypertension is a known risk factor for cerebral small vessel disease [102] and stroke [4], hypertension is often considered a risk factor for vascular dementia, a form of cognitive decline resulting from small- or large-vessel cerebrovascular disease [9,103]. However, only a handful of studies have directly examined the relationship between hypertension and vascular dementia. Although previous research supports the relationship between midlife hypertension and the development of vascular dementia [8,83,85,104–106], it is unclear whether there is an association between late-life hypertension and vascular dementia, as findings have thus far been conflicting [72,89,91,107]. Compared to the associations between midlife hypertension and incident Alzheimer's disease, the associations found between midlife hypertension and incident vascular dementia tend to be more robust and consistent [8,85,108]. However, because patients are more likely to develop mixed Alzheimer's and vascular dementia than pure forms of one or the other, this distinction may not be meaningful.

Pathophysiology of hypertension as it relates to cognitive decline

Evidence from neuroimaging and biomarker studies

Neuroimaging has played a pivotal role in advancing the understanding of how BP influences cognitive function and underlying brain structure. Results from studies that have examined the relationship between BP and brain volume are largely consistent with findings from the BP and cognition studies. High SBP has been associated with smaller regional and total brain volumes [109–113] and reductions in brain volume over time [114]. The relationship between high DBP and brain volume is less consistent, however [110,112,113,115]. In elderly populations, low SBP [116,117] and low DBP [117,118] have

been associated with reduced brain volume and cortical thickness, suggesting that the relationship between BP and brain volume may age-dependent [15,119,120]. A pattern of hypertension in midlife followed by hypotension in late-life appears to be especially harmful [34], particularly in brain regions affected in the earliest phase of Alzheimer's disease [121].

An association between hypertension and the development of Alzheimer's disease has also been supported by research that examines Alzheimer's disease biomarkers directly. Compared to the brains of normotensive individuals, the brains of individuals with a history of hypertension show greater levels of β -amyloid plaques, atrophy, and neurofibrillary tangles [86,122]. Similarly, hypertension has been identified as a risk factor for cortical fibrillar β -amyloid deposits [81,123,124] and reduced glucose metabolism in Alzheimer's disease-specific brain regions [123,125] using positron emission tomography (PET) in the brains of cognitively normal middle-aged and older adults. Consistent with these findings, one study found that individuals with abnormal plasma β -amyloid levels and elevated BP at midlife have an especially high risk of developing Alzheimer's disease later in life [7].

Hypertension has also been associated with several defining features of vascular dementia and cerebral small vessel disease, including WMH volume [15,102], WMH progression [126,127], lacunar infarcts, and cerebral microbleeds [5,128–130]. Supporting the relationship between high BP and white matter pathology, findings from observational studies [127] and clinical trials [113,131] suggest that treatment of hypertension reduces WMH progression. Even before the development of overt neuroimaging abnormalities, hypertension appears to be associated with reduced white matter microstructural integrity in both young and old individuals, suggesting white matter may be especially vulnerable to the deleterious effects of hypertension [132–135].

Hypertension and vascular remodeling

Emerging evidence suggests that sustained elevations in BP may cause cerebral vessel remodeling in a manner which promotes pathological brain changes and subsequent cognitive decline. To preserve the steady low-pressure blood supply to the periphery and protect end organ microcirculation from pulsatile stress associated with hypertension, a rearrangement in vessel wall material in the form of hypertrophic remodeling of the media and vascular smooth muscle cells occurs [136–138]. This enlargement in media size causes a reduction in lumen diameter, leading to increased vascular resistance and vessel wall stiffening [139]. Arterial stiffening, in turn, increases arterial pulse wave velocity and pulsatile pressure, which over time causes rarefaction of downstream capillaries and further inward remodeling of vessel walls [140–142]. Hypertension promotes intracranial atherosclerosis in large intracranial arteries [142,143] and arteriolosclerosis in smaller arterioles supplying subcortical white matter and deep gray matter brain structures [144]. Arteriolosclerosis is a process characterized by a loss of tunica media smooth muscle cells, fibro-hyaline deposits, and thickening of the vessel wall, resulting in increased microvascular resistance. Because the brain requires high levels of continuous perfusion throughout systole and diastole [145], increases in vascular resistance leave cerebral arterioles vulnerable to hypoperfusion when systemic BP is reduced [139,144]. As described

below, hypoperfusion has been associated with several neurovascular changes [82], which together may disrupt cognition [146].

Autoregulation and cerebral perfusion

The brain requires a high volume of consistent blood flow to sustain adequate perfusion. However, the brain's ability to maintain steady low-pressure blood flow in the context of changing systemic BP – a process known as cerebral autoregulation – can be disrupted as a result of chronic hypertension [147,148]. After prolonged exposure to high BP and elevated pulsatility, a shift occurs in the brain's autoregulatory capacity whereby higher systemic BP is required to maintain the same level of cerebral perfusion [149]. Hypertension is believed to alter cerebral autoregulation by inducing changes in arteriole endothelial and vascular smooth muscle cells that diminish cerebrovascular reactivity [150] and increase myogenic tone, respectively [151]. Not only do these vascular changes shift the cerebral autoregulatory curve in a manner which reduces resting cerebral blood flow, but the brain also becomes more susceptible to hypoperfusion during periods of low systemic BP [152] or during periods of normal BP in chronically hypertensive individuals [153]. These hypertension-induced changes to cerebral autoregulation and perfusion may explain why individuals with chronic hypertension in midlife and low BP in late-life show significant reductions in brain volume [34,121] and greater levels of cognitive deficits [101].

While ischemia may occur in some cases, the brain is more likely to be subjected to chronic oligemia (i.e., mild reductions in blood flow) as a result of hypertension. Chronic oligemia may, in turn, lead to endothelial dysfunction, acidosis, oxidative stress, and unmet metabolic energy demands that can impair neuronal function [82,154,155]. Oligemia may also down-regulate the synthesis of proteins necessary for synaptic plasticity and memory formation [155], and promote neuronal tau phosphorylation, β -amyloid oligomerization, and the upregulation of amyloidogenic APP [156–160]. Each of these neurophysiological changes likely contributes to the development of Alzheimer's disease and cerebral amyloid angiopathy (CAA). Evidence suggests that β -amyloid accumulation may also occur as a result of hypertension-induced up-regulation of the receptor for advanced glycation end products (RAGE), which controls the shuttling of β -amyloid from the blood across the endothelial barrier into the brain [161].

Endothelial dysfunction, altered functional hyperemia & Oxidative stress

By promoting endothelial dysfunction, hypertension is also believed to disrupt the coordinated coupling among neurons, glia, and cerebral blood flow in the microvasculature [162]. Uncoupling of this system, known collectively as the neurovascular unit, can impair the homeostatic process of functional hyperemia, whereby increases in CBF occur in coordination with increases in neuronal activity to ensure the delivery of adequate levels oxygen and glucose and facilitate the removal of metabolites [163–165]. Support for these findings comes from animal research, which has demonstrated that hypertension-induced vascular oxidative stress resulting from up-regulation of reactive oxygen species (ROS)-producing enzyme NADPH oxidase impairs the endothelium-dependent expression of vasodilators and vasoconstrictors necessary to maintain neurovascular coupling [150,166,167].

Antihypertensive clinical trials to improve cognition

Given the apparent association between BP and cognitive function, efforts have been made to determine whether improved BP control can be used to delay cognitive decline and reduce dementia risk. To date, evidence from large placebo-controlled, randomized clinical trials (RCTs) has been conflicting [168,169]. A 2009 Cochrane Review of randomized, double-blind, placebo-controlled trials concluded that there is currently no convincing evidence for the protective effects of antihypertensive use in late-life [169]. Although several large placebo-controlled RCTs, such as the Perindopril Protection Against Recurrent Stroke Study (PROGRESS) [170], the Systolic Hypertension in Europe (SYST-EUR study) [171], and the Heart Outcomes Prevention Evaluation (HOPE) study [172] have found antihypertensive agents to be protective against cognitive decline and dementia, just as many trials have failed to replicate this finding [173–177]. Thus, it is unknown whether BP control alone is enough to reduce the risk of cognitive decline. It is possible that the neuroprotective effects of antihypertensive agents may result from drug-specific neurobiological changes as opposed to (or in addition to) BP lowering [178,179]. In support of this idea, a meta-analysis of RCTs which compared the neuroprotective properties of different antihypertensive drug classes found angiotensin receptor blockers (ARBs) to be superior to β -blockers, diuretics, and ACE inhibitors for preventing cognitive decline [180].

The ability to draw conclusions about the effectiveness of BP interventions for the reduction of cognitive decline has been limited by brief study durations and insufficient power to detect effects. Cognitive decline, even in the course of neurodegenerative disease, is a relatively gradual process, and, as described above, elevated BP in midlife may be the most important determinant of risk for subsequent cognitive decline and dementia. Thus, midlife may be the most critical window during which BP control must begin. Extended treatment and follow-up periods and larger sample sizes will likely be needed to reliably detect the effects of BP lowering on cognitive measures. By comparison, neurodegenerative and dementia-specific biomarkers (e.g., hippocampal atrophy and CSF-tau) may be more sensitive to treatment-related effects, but their validity as intermediate endpoints remains a subject of debate [181,182]. Future studies may also benefit from making use of a more comprehensive cognitive battery. The Mini-Mental State Examination (MMSE), which has been used to assess cognitive abilities in the majority of previous trials, is notoriously insensitive to cognitive change, especially in domains of executive functioning and processing speed, making it an especially poor choice for detecting cognitive change in this context [183,184]. Additionally, effect sizes in previous BP lowering trials may have been attenuated because participants receiving antihypertensive medication often saw only minor reductions in BP compared to participants given placebo. This limitation is addressed in an ongoing trial (SPRINT-MIND) to evaluate the neuroprotective effects of reducing BP to below a specific level (i.e., below 120mm Hg) using one or more antihypertensive agent [185]. The parent trial to this study (SPRINT) has already demonstrated improved cardiovascular outcomes in the setting of this tighter blood pressure control [186]. However, the ability of this trial to show benefit in cognitive outcomes will be limited by short follow-up.

Conclusions and future directions

It is clear that hypertension can affect brain structure and function in a manner that increases one's risk of cognitive decline and dementia. Hypertension, high SBP, and high DBP during midlife have been most consistently linked to late-life cognitive decline and incident dementia. However, hypertension has been associated with early-life and midlife cognitive deficits as well. Although the association between late-life hypertension and cognitive function is less clear, particularly among octogenarians and nonagenarians, limited evidence suggests that mildly elevated BP in late life may be protective against cognitive decline, especially for individuals with a history of longstanding hypertension. Hypertension duration may be an especially important determinant of cognitive decline, as evidence suggests that the damaging neurological effects of hypertension may be cumulative. Few studies have assessed BP longitudinally, and even fewer have attempted to retrospectively determine how lifetime duration of hypertension relates to cognitive function. Given the increasing prevalence of hypertension among younger individuals [187], assessing the cumulative effects of elevated BP over the lifespan will be especially important to understanding how BP may influence neurodevelopment and neurodegeneration [188].

Recent advances in neuroimaging and physiologic and hemodynamic monitoring have allowed for an improved understanding of the mechanisms through which hypertension affects neurocognitive function. Hypertension, especially in midlife, has been identified as a risk factor for cerebral atrophy, white matter microstructural damage, and cerebral small vessel disease. Evidence suggests that hypertension contributes to the development and progression of such neurological changes by promoting vessel wall remodeling and endothelial dysfunction, which results in autoregulatory deficits. These changes to the neurovascular unit leave the brain vulnerable to hypoperfusion resulting from drops in systemic BP. Although evidence exists to support this model of hypertension-induced cerebrovascular changes, much is still unknown about how these pathophysiological processes directly influence cognitive function and promote Alzheimer's and vascular dementia in humans.

Additional insights into the role circulatory changes play in cognitive decline will likely come from the study of other markers of vessel function. For example, pulse pressure, a measure of arterial stiffening, which increases with age and exposure to hypertension [145], can be used as an additional method to quantify the effects of vascular pathology resulting from chronic hypertension. Elevations in pulse pressure have been associated with cognitive impairment [189,190], cognitive decline [190], cerebral small vessel disease [127,191], and Alzheimer's disease biomarkers [192]. Compared to BP, pulse pressure is believed to more precisely quantify the exposure of target organs such as the brain to potentially harmful pulsatile energy resulting from arterial stiffening [193].

A more nuanced understanding of the relationship between BP and neural function will likely be needed before antihypertensive therapies can be effectively employed as an intervention to reduce cognitive decline. Given that many individuals who develop hypertension do so before late-life and experience the harmful effects of hypertension for decades, it is unclear whether specific antihypertensive agents will be able to modify the

trajectory of cognitive decline within the span of a multi-year trial. If the effects of hypertension on the brain are cumulative, interindividual differences in the duration and severity of previous hypertension must be considered in future trial design. Because the effects of BP on cognition appear to differ with age, future clinical trials may also benefit from limiting enrollment to specific age groups. Other factors such as race, sex, genetics, and the presence of cerebrovascular morbidity have each been identified as effect modifiers in observational studies and should, therefore, be considered when designing future antihypertensive trials.

Acknowledgments

Acknowledgments: Keenan A. Walker was supported by the NIA (T32 AG027668).

References

1. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, et al. Global disparities of hypertension prevalence and control. *Circulation*. 2016;134:441–50. [PubMed: 27502908]
2. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: Analysis of worldwide data. *Lancet*. 2005;365:217–23. [PubMed: 15652604]
3. Fanning JP, Wong AA, Fraser JF. The epidemiology of silent brain infarction: a systematic review of population-based cohorts. *BMC Med* 2014;12:119. [PubMed: 25012298]
4. Knopman DS, Penman AD, Catellier DJ, Coker LH, Shibata DK, Sharrett AR, et al. Vascular risk factors and longitudinal changes on brain MRI: the ARIC study. *Neurology*. American Academy of Neurology; 2011;76:1879–85.
5. Bezerra DC, Sharrett AR, Matsushita K, Gottesman RF, Shibata D, Mosley TH, et al. Risk factors for lacune subtypes in the Atherosclerosis Risk in Communities (ARIC) study. *Neurology*. 2012;78:102–8. [PubMed: 22170882]
6. MacMahon S, Peto R, Collins R, Godwin J, MacMahon S, Cutler J, et al. Blood pressure, stroke, and coronary heart disease. Part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet*. 1990;335:765–74. [PubMed: 1969518]
7. Shah NS, Vidal JS, Masaki K, Petrovitch H, Ross GW, Tilley C, et al. Midlife blood pressure, plasma beta-amyloid, and the risk for alzheimer disease: The honolulu asia aging study. *Hypertension*. 2012;59:780–6. [PubMed: 22392902]
8. Rönnemaa E, Zethelius B, Lannfelt L, Kilander L. Vascular risk factors and dementia: 40-year follow-up of a population-based cohort. *Dement. Geriatr. Cogn. Disord* 2011;31:460–6. [PubMed: 21791923]
9. Kennelly SP, Lawlor BA, Kenny RA. Blood pressure and dementia - a comprehensive review. *Ther. Adv. Neurol. Disord* 2009;2:241–60. [PubMed: 21179532]
10. Kilander L, Nyman H, Boberg M, Lithell H. The association between low diastolic blood pressure in middle age and cognitive function in old age. A population-based study. *Age Ageing*. 2000;29:243–8. [PubMed: 10855907]
11. Kilander L, Nyman H, Boberg M, Hansson L, Lithell H. Hypertension is related to cognitive impairment: a 20-year follow-up of 999 men. *Hypertension*. 1998;31:780–6. [PubMed: 9495261]
12. ** Launer LJ, Masaki K, Petrovitch H, Foley D, Havlik RJ. The association between midlife blood pressure levels and late-life cognitive function. The Honolulu-Asia Aging Study. *JAMA*. 1995;274:1846–51. [PubMed: 7500533] One of the earliest prospective observational studies to demonstrate a link between midlife systolic blood pressure and reduced cognitive function later in life
13. * Gottesman RF, Schneider ALC, Albert M, Alonso A, Bandeen-Roche K, Coker L, et al. Midlife Hypertension and 20-Year Cognitive Change. *JAMA Neurol* 2014;21287:1–10. This large

longitudinal study found that hypertension and high blood pressure during midlife were associated with greater cognitive decline over a 20-year follow-up period

14. Knopman DS, Mosley TH, Catellier DJ, Coker LH, Atherosclerosis Risk in Communities Study Brain MRI Study. Fourteen-year longitudinal study of vascular risk factors, APOE genotype, and cognition: the ARIC MRI Study. *Alzheimers. Dement* 2009;5:207–14. [PubMed: 19362884]
15. Swan GE, DeCarli C, Miller BL, Reed T, Wolf PA, Jack LM, et al. Association of midlife blood pressure to late-life cognitive decline and brain morphology. *Neurology*. 1998;51:986–93. [PubMed: 9781518]
16. Schneider ALC, Sharrett AR, Patel MD, Alonso A, Coresh J, Mosley T, et al. Education and cognitive change over 15 years: The atherosclerosis risk in communities study. *J. Am. Geriatr. Soc* 2012;60:1847–53. [PubMed: 23013064]
17. * Swan GE, Carmelli D, Larue A. Systolic blood pressure tracking over 25 to 30 years and cognitive performance in older adults. *Stroke*. 1998;29:2334–40. [PubMed: 9804644] This is one of the few longitudinal studies that has examined how the trajectory of blood pressure over multiple decades relates to cognition later in life
18. ** Power MC, Tchetgen EJT, Sparrow D, Schwartz J, Weisskopf MG. Blood pressure and cognition: factors that may account for their inconsistent association. *Epidemiology*. 2013;24:886–93. [PubMed: 24030502] This longitudinal study demonstrated the importance of examining hypertension duration; a greater duration of time since the onset of hypertension was associated with lower cognitive function later in life
19. Kuo H-K, Sorond F, Iloputaife I, Gagnon M, Milberg W, Lipsitz LA. Effect of blood pressure on cognitive functions in elderly persons. *J. Gerontol. A. Biol. Sci. Med. Sci* 2004;59:1191–4. [PubMed: 15602074]
20. Budge MM, De Jager C, Hogervorst E, Smith AD. Total plasma homocysteine, age, systolic blood pressure, and cognitive performance in older people. *J. Am. Geriatr. Soc* 2002;50:2014–8. [PubMed: 12473014]
21. Vinyoles E, De la Figuera M, Gonzalez-Segura D. Cognitive function and blood pressure control in hypertensive patients over 60 years of age: COGNIPRES study. *Curr. Med. Res. Opin* 2008;24:3331–9. [PubMed: 18954496]
22. Tsivgoulis G, Alexandrov A V, Wadley VG, Unverzagt FW, Go RCP, Moy CS, et al. Association of higher diastolic blood pressure levels with cognitive impairment. *Neurology*. 2009;73:589–95. [PubMed: 19704077]
23. Elias MF, Wolf PA, D'Agostino RB, Cobb J, White LR. Untreated blood pressure level is inversely related to cognitive functioning: the Framingham Study. *Am. J. Epidemiol* 1993;138:353–64. [PubMed: 8213741]
24. Cherbuin N, Reglade-Meslin C, Kumar R, Jacomb P, Eastale S, Christensen H, et al. Risk factors of transition from normal cognition to mild cognitive disorder: The PATH through life study. *Dement. Geriatr. Cogn. Disord* 2009;28:47–55. [PubMed: 19628940]
25. Reitz C, Tang M-X, Manly J, Mayeux R, Luchsinger JA. Hypertension and the risk of mild cognitive impairment. *Arch. Neurol* 2007;64:1734–40. [PubMed: 18071036]
26. Scherr PA, Hebert LE, Smith LA, Evans DA. Relation of blood pressure to cognitive function in the elderly. *Am. J. Epidemiol* 1991;134:1303–15. [PubMed: 1755444]
27. Pandav R, Dodge HH, DeKosky ST, Ganguli M. Blood pressure and cognitive impairment in India and the United States: a cross-national epidemiological study. *Arch. Neurol* 2003;60:1123–8. [PubMed: 12925370]
28. Morris MC, Scherr PA, Hebert LE, Bennett DA, Wilson RS, Glynn RJ, et al. Association between blood pressure and cognitive function in a biracial community population of older persons. *Neuroepidemiology*. 2002;21:123–30. [PubMed: 12006775]
29. Waldstein SR, Giggey PP, Thayer JF, Zonderman AB. Nonlinear relations of blood pressure to cognitive function: The Baltimore longitudinal study of aging. *Hypertension*. 2005;45:374–9. [PubMed: 15699446]
30. Shang S, Li P, Deng M, Jiang Y, Chen C, Qu Q. The Age-Dependent Relationship between Blood Pressure and Cognitive Impairment: A Cross-Sectional Study in a Rural Area of Xi'an, China. *PLoS One*. 2016;11:e0159485. [PubMed: 27438476]

31. Gorelick PB, Nyenhuis D. Blood pressure and treatment of persons with hypertension as it relates to cognitive outcomes including executive function. *J. Am. Soc. Hypertens* 2012;6:309–15. [PubMed: 22995799]
32. Glynn RJ, Beckett LA, Hebert LE, Morris MC, Scherr PA, Evans DA. Current and remote blood pressure and cognitive decline. *JAMA*. 1999;281:438–45. [PubMed: 9952204]
33. Bohannon AD, Fillenbaum GG, Pieper CF, Hanlon JT, Blazer DG. Relationship of race/ethnicity and blood pressure to change in cognitive function. *J. Am. Geriatr. Soc* 2002;50:424–9. [PubMed: 11943035]
34. ** Muller M, Sigurdsson S, Kjartansson O, Aspelund T, Lopez OL, Jonnson PV, et al. Joint effect of mid- and late-life blood pressure on the brain: The AGES-Reykjavik Study. *Neurology*. 2014;82:2187–95. [PubMed: 24898928] This large longitudinal study demonstrated that the association between late-life blood pressure, cognition, and brain volume was dependent on midlife blood pressure; a combination of midlife hypertension and low late-life diastolic blood pressure was associated with worse outcomes
35. Suhr JA, Stewart JC, France CR. The relationship between blood pressure and cognitive performance in the Third National Health and Nutrition Examination Survey (NHANES III). *Psychosom. Med* 2004;66:291–7. [PubMed: 15184686]
36. Adams HR, Szilagyi PG, Gebhardt L, Lande MB. Learning and attention problems among children with pediatric primary hypertension. *Pediatrics*. 2010;126:e1425–9. [PubMed: 21059718]
37. Lande MB, Batsisky DL, Kupferman JC, Samuels J, Hooper SR, Falkner B, et al. Neurocognitive Function in Children with Primary Hypertension. *J. Pediatr* 2016;
38. Hill CM, Hogan AM, Onugha N, Harrison D, Cooper S, McGrigor VJ, et al. Increased cerebral blood flow velocity in children with mild sleep-disordered breathing: a possible association with abnormal neuropsychological function. *Pediatrics*. 2006;118:e1100–8. [PubMed: 17015501]
39. Ditto B, Séguin JR, Tremblay RE. Neuropsychological characteristics of adolescent boys differing in risk for high blood pressure. *Ann. Behav. Med* 2006;31:231–7. [PubMed: 16700636]
40. Lande MB, Kaczorowski JM, Auinger P, Schwartz GJ, Weitzman M. Elevated blood pressure and decreased cognitive function among school-age children and adolescents in the United States. *J. Pediatr* 2003 p. 720–4. [PubMed: 14657815]
41. O'Callaghan S, Kenny RA. Neurocardiovascular Instability and Cognition. *Yale J. Biol. Med. Yale Journal of Biology and Medicine*; 2016;89:59–71.
42. O'Brien E, Coats A, Owens P, Petrie J, Padfield PL, Littler WA, et al. Use and interpretation of ambulatory blood pressure monitoring: recommendations of the British hypertension society. *BMJ*. 2000;320:1128–34. [PubMed: 10775227]
43. Rose KM, Tyroler HA, Nardo CJ, Arnett DK, Light KC, Rosamond W, et al. Orthostatic hypotension and the incidence of coronary heart disease: the Atherosclerosis Risk in Communities study. *Am. J. Hypertens* 2000;13:571–8. [PubMed: 10912737]
44. Rose KM, Holme I, Light KC, Sharrett AR, Tyroler HA, Heiss G. Association between the blood pressure response to a change in posture and the 6-year incidence of hypertension: prospective findings from the ARIC study. *J. Hum. Hypertens* 2002;16:771–7. [PubMed: 12444538]
45. Czajkowska J, Ozhog S, Smith E, Perlmutter LC. Cognition and hopelessness in association with subsyndromal orthostatic hypotension. *J. Gerontol. A. Biol. Sci. Med. Sci* 2010;65:873–9. [PubMed: 20478907]
46. Elmståhl S, Widerström E. Orthostatic intolerance predicts mild cognitive impairment: incidence of mild cognitive impairment and dementia from the Swedish general population cohort Good Aging in Skåne Clin. Interv. Aging. Dove Press; 2014;9:1993–2002.
47. Frewen J, Savva GM, Boyle G, Finucane C, Kenny RA. Cognitive performance in orthostatic hypotension: findings from a nationally representative sample. *J. Am. Geriatr. Soc* 2014;62:117–22. [PubMed: 25180380]
48. Frewen J, Finucane C, Savva GM, Boyle G, Kenny RA. Orthostatic hypotension is associated with lower cognitive performance in adults aged 50 plus with supine hypertension. *J. Gerontol. A. Biol. Sci. Med. Sci* 2014;69:878–85. [PubMed: 24214492]

49. Schoon Y, Lagro J, Verhoeven Y, Rikkert MO, Claassen J. Hypotensive syndromes are not associated with cognitive impairment in geriatric patients. *Am. J. Alzheimers. Dis. Other Dement* 2013;28:47–53. [PubMed: 23242123]
50. Rose KM, Couper D, Eigenbrodt ML, Mosley TH, Sharrett AR, Gottesman RF. Orthostatic hypotension and cognitive function: the Atherosclerosis Risk in Communities Study. *Neuroepidemiology*. 2010;34:1–7. [PubMed: 19893322]
51. Viramo P, Luukinen H, Koski K, Laippala P, Sulkava R, Kivelä SL. Orthostatic hypotension and cognitive decline in older people. *J. Am. Geriatr. Soc* 1999;47:600–4. [PubMed: 10323655]
52. Kanemaru A, Kanemaru K, Kuwajima I. The effects of short-term blood pressure variability and nighttime blood pressure levels on cognitive function. *Hypertens. Res* 2001;24:19–24. [PubMed: 11213025]
53. Bellelli G, GB F, Lucchi E, Guerini F, Geroldi C, Magnifico F, et al. Blunted reduction in night-time blood pressure is associated with cognitive deterioration in subjects with long-standing hypertension. *Blood Press. Monit* 2004;9:71–76 pp. [PubMed: 15096903]
54. Sakakura K, Ishikawa J, Okuno M, Shimada K, Kario K. Exaggerated Ambulatory Blood Pressure Variability Is Associated with Cognitive Dysfunction in the Very Elderly and Quality of Life in the Younger Elderly. *Am. J. Hypertens* 2007;20:720–7. [PubMed: 17586405]
55. Allan LM, Ballard CG, Allen J, Murray A, Davidson AW, McKeith IG, et al. Autonomic dysfunction in dementia. *J. Neurol. Neurosurg. Psychiatry. BMJ Group*; 2007;78:671–7.
56. Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, Soubrier F. An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J. Clin. Invest* 1990;86:1343–6. [PubMed: 1976655]
57. Schuch JB, Constantin PC, da Silva VK, Korb C, Bamberg DP, da Rocha TJ, et al. ACE polymorphism and use of ACE inhibitors: effects on memory performance. *Age (Dordr)* 2014;36:9646. [PubMed: 24696269]
58. Saidi S, Zammiti W, Slamia LB, Ammou SB, Almawi WY, Mahjoub T. Interaction of angiotensin-converting enzyme and apolipoprotein E gene polymorphisms in ischemic stroke involving large-vessel disease. *J. Thromb. Thrombolysis* 2009;27:68–74. [PubMed: 18027070]
59. Hassan A, Lansbury A, Catto AJ, Guthrie A, Spencer J, Craven C, et al. Angiotensin converting enzyme insertion/deletion genotype is associated with leukoaraiosis in lacunar syndromes. *J. Neurol. Neurosurg. Psychiatry*. 2002;72:343–6. [PubMed: 11861692]
60. Bartrés-Faz D, Junqué C, Clemente IC, López-Alomar A, Valveny N, López-Guillén A, et al. Angiotensin I converting enzyme polymorphism in humans with age-associated memory impairment: relationship with cognitive performance. *Neurosci Lett* 2000;290:177–80. [PubMed: 10963892]
61. * Richard F, Berr C, Amant C, Helbecque N, Amouyel P, Alperovitch A. Effect of the angiotensin I-converting enzyme I/D polymorphism on cognitive decline. The EVA Study Group. *Neurobiol Aging* 2000;21:75–80. One of the first large prospective studies to demonstrate a relationship between the angiotensin I-converting enzyme I/D polymorphism and cognitive decline in the elderly
62. Raz N, Dahle CL, Rodrigue KM, Kennedy KM, Land S. Effects of age, genes, and pulse pressure on executive functions in healthy adults. *Neurobiol. Aging*. 2011;32:1124–37. [PubMed: 19559505]
63. Wang B, Jin F, Yang Z, Lu Z, Kan R, Li S, et al. The insertion polymorphism in angiotensin-converting enzyme gene associated with the APOE epsilon 4 allele increases the risk of late-onset Alzheimer disease. *J. Mol. Neurosci* 2006;30:267–71. [PubMed: 17401152]
64. Lehmann DJ, Cortina-Borja M, Warden DR, Smith AD, Sleegers K, Prince JA, et al. Large meta-analysis establishes the ACE insertion-deletion polymorphism as a marker of Alzheimer's disease. *Am. J. Epidemiol* 2005;162:305–17. [PubMed: 16033878]
65. Taylor WD, Benjamin S, McQuoid DR, Payne ME, Krishnan RR, MacFall JR, et al. AGTR1 gene variation: Association with depression and frontotemporal morphology. *Psychiatry Res. - Neuroimaging* 2012;202:104–9.

66. Zannas AS, McQuoid DR, Payne ME, Macfall JR, Ashley-Koch A, Steffens DC, et al. Association of gene variants of the renin-angiotensin system with accelerated hippocampal volume loss and cognitive decline in old age. *Am. J. Psychiatry.* 2014;171:1214–21. [PubMed: 25124854]
67. McFall GP, Sapkota S, McDermott KL, Dixon RA. Risk-reducing Apolipoprotein E and Clusterin genotypes protect against the consequences of poor vascular health on executive function performance and change in nondemented older adults. *Neurobiol. Aging* 2016;42:91–100. [PubMed: 27143425]
68. Andrews S, Das D, Anstey KJ, Eastel S. Interactive effect of APOE genotype and blood pressure on cognitive decline: The PATH through life study. *J. Alzheimer's Dis* 2015;44:1087–98. [PubMed: 25672766]
69. de Frias CM, Warner SK, Willis SL. Hypertension moderates the effect of APOE on 21-year cognitive trajectories. *Psychol. Aging* 2014;29:431–9. [PubMed: 24956008]
70. Li W, Wang T, Xiao S. Type 2 diabetes mellitus might be a risk factor for mild cognitive impairment progressing to Alzheimer's disease. *Neuropsychiatr. Dis. Treat* 2016;12:2489–95. [PubMed: 27729793]
71. Xu W, Qiu C, Gatz M, Pedersen NL, Johansson B, Fratiglioni L. Mid- And late-life diabetes in relation to the risk of dementia: A population-based twin study. *Diabetes.* 2009;58:71–7. [PubMed: 18952836]
72. Raffaitin C, Gin H, Empana J-P, Helmer C, Berr C, Tzourio C, et al. Metabolic syndrome and risk for incident Alzheimer's disease or vascular dementia: the Three-City Study. *Diabetes Care.* 2009;32:169–74. [PubMed: 18945929]
73. Rosendorff C, Beeri MS, Silverman JM. Cardiovascular risk factors for Alzheimer's disease. *Am. J. Geriatr. Cardiol* 2007 p. 143–9. [PubMed: 17483665]
74. Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W, Ferri CP. The global prevalence of dementia: a systematic review and metaanalysis. *Alzheimer's Dement* 2013;9:63–75. [PubMed: 23305823]
75. Gorelick PB, Scuteri A, Black SE, Decarli C, Greenberg SM, Iadecola C, et al. Vascular contributions to cognitive impairment and dementia: A statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2011 p. 2672–713. [PubMed: 21778438]
76. Lopez MF, Krastins B, Ning M. The role of apolipoprotein E in neurodegeneration and cardiovascular disease. *Expert Rev. Proteomics.* 2014;11:371–81. [PubMed: 24754513]
77. Traylor M, Adib-Samii P, Harold D, Alzheimer's Disease Neuroimaging Initiative, International Stroke Genetics Consortium (ISGC), UK Young Lacunar Stroke DNA resource, Dichgans M, et al. Shared genetic contribution to Ischaemic Stroke and Alzheimer's Disease. *Ann. Neurol* 2016;
78. Schneider JA, Arvanitakis Z, Bang W, Bennett DA. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology.* 2007;69:2197–204. [PubMed: 17568013]
79. Sonnen JA, Larson EB, Crane PK, Haneuse S, Li G, Schellenberg GD, et al. Pathological correlates of dementia in a longitudinal, population-based sample of aging. *Ann. Neurol* 2007;62:406–13. [PubMed: 17879383]
80. Vascular Iadecola C. and Metabolic Factors in Alzheimer's Disease and Related Dementias: Introduction. *Cell. Mol. Neurobiol* 2016 3;36:151–4. [PubMed: 26898551]
81. Rodrigue KM, Rieck JR, Kennedy KM, Devous MD, Diaz-Arrastia R, Park DC. Risk factors for β -amyloid deposition in healthy aging: vascular and genetic effects. *JAMA Neurol* 2013;70:600–6. [PubMed: 23553344]
82. Zlokovic BV Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat. Rev. Neurosci* 2011;12:723–38. [PubMed: 22048062]
83. * Launer LJ, Ross GW, Petrovitch H, Masaki K, Foley D, White LR, et al. Midlife blood pressure and dementia: The Honolulu-Asia aging study. *Neurobiol. Aging* 2000;21:49–55. [PubMed: 10794848] This prospective observational study provided early evidence that treatment of blood pressure reduced the risk for dementia associated with midlife hypertension
84. Kivipelto M, Helkala EL, Laakso MP, Hänninen T, Hallikainen M, Alhainen K, et al. Apolipoprotein e4 allele, elevated midlife total cholesterol level, and high midlife systolic blood

pressure are independent risk factors for late-life Alzheimer disease. *Ann. Intern. Med.* 2002;137:149–55. [PubMed: 12160362]

85. Yamada M, Kasagi F, Sasaki H, Masunari N, Mimori Y, Suzuki G. Association between dementia and midlife risk factors: the Radiation Effects Research Foundation Adult Health Study. *J. Am. Geriatr. Soc* 2003;51:410–4. [PubMed: 12588587]
86. Petrovitch H, White LR, Izmirlian G, Ross GW, Havlik RJ, Markesbery W, et al. Midlife blood pressure and neuritic plaques, neurofibrillary tangles, and brain weight at death: the HAAS. *Neurobiol. Aging* 2000;21:57–62. [PubMed: 10794849]
87. Morris MC, Scherr PA, Hebert LE, Glynn RJ, Bennett DA, Evans DA. Association of incident Alzheimer disease and blood pressure measured from 13 years before to 2 years after diagnosis in a large community study. *Arch. Neurol* 2001;58:1640–6. [PubMed: 11594923]
88. Power MC, Weuve J, Gagne JJ, McQueen MB, Viswanathan A, Blacker D. The association between blood pressure and incident Alzheimer disease: a systematic review and meta-analysis. *Epidemiology.* 2011;22:646–59. [PubMed: 21705906]
89. Forti P, Pisacane N, Rietti E, Lucicesare A, Olivelli V, Mariani E, et al. Metabolic syndrome and risk of dementia in older adults. *J. Am. Geriatr. Soc* 2010;58:487–92. [PubMed: 20398117]
90. * Li G, Rhew IC, Shofer JB, Kukull WA, Breitner JCS, Peskind E, et al. Age-varying association between blood pressure and risk of dementia in those aged 65 and older: a community-based prospective cohort study. *J. Am. Geriatr. Soc* 2007;55:1161–7. [PubMed: 17661953] This prospective study found strong evidence for an age-dependent association between systolic blood pressure and dementia risk: the association between high systolic blood pressure and dementia risk decreased with increasing age
91. Verghese J, Lipton RB, Hall CB, Kuslansky G, Katz MJ. Low blood pressure and the risk of dementia in very old individuals. *Neurology.* 2003;61:1667–72. [PubMed: 14694027]
92. Shah RC, Wilson RS, Bienias JL, Arvanitakis Z, Evans DA, Bennett DA. Relation of blood pressure to risk of incident Alzheimer's disease and change in global cognitive function in older persons. *Neuroepidemiology.* 2005;26:30–6. [PubMed: 16254451]
93. Muller M, Tang M-X, Schupf N, Manly JJ, Mayeux R, Luchsinger JA. Metabolic syndrome and dementia risk in a multiethnic elderly cohort. *Dement. Geriatr. Cogn. Disord* 2007;24:185–92. [PubMed: 17641531]
94. Qiu C, Winblad B, Fastbom J, Fratiglioni L. Combined effects of APOE genotype, blood pressure, and antihypertensive drug use on incident AD. *Neurology.* 2003;61:655–60. [PubMed: 12963757]
95. Petitti DB, Crooks VC, Buckwalter JG, Chiu V. Blood pressure levels before dementia. *Arch. Neurol* 2005;62:112–6. [PubMed: 15642857]
96. Ruitenberg A, Skoog I, Ott A, Aevarsson O, Witteman JCM, Lernfelt B, et al. Blood pressure and risk of dementia: Results from the Rotterdam study and the Gothenburg H-70 study. *Dement. Geriatr. Cogn. Disord* 2001;12:33–9. [PubMed: 11125239]
97. Qiu C, Winblad B, Marengoni A, Klarin I, Fastbom J, Fratiglioni L. Heart failure and risk of dementia and Alzheimer disease: a population-based cohort study. *Arch. Intern. Med* 2006;166:1003–8. [PubMed: 16682574]
98. Nilsson SE, Read S, Berg S, Johansson B, Melander A, Lindblad U. Low systolic blood pressure is associated with impaired cognitive function in the oldest old: longitudinal observations in a population-based sample 80 years and older. *Aging Clin. Exp. Res* 2007;19:41–7. [PubMed: 17332720]
99. Qiu C, Von Strauss E, Winblad B, Fratiglioni L. Decline in blood pressure over time and risk of dementia: A longitudinal study from the Kungsholmen project. *Stroke.* 2004;35:1810–5. [PubMed: 15232128]
100. Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, et al. 15-Year Longitudinal Study of Blood Pressure and Dementia. *Lancet* (London, England). 1996;347:1141–5.
101. Glodzik L, Rusinek H, Pirraglia E, McHugh P, Tsui W, Williams S, et al. Blood pressure decrease correlates with tau pathology and memory decline in hypertensive elderly. *Neurobiol. Aging* 2014;35:64–71. [PubMed: 23969178]

102. Raz N, Rodrigue KM, Kennedy KM, Acker JD. Vascular health and longitudinal changes in brain and cognition in middle-aged and older adults. *Neuropsychology*. 2007;21:149–57. [PubMed: 17402815]
103. Ying H, Jianping C, Jianqing Y, Shanquan Z. Cognitive variations among vascular dementia subtypes caused by small-, large-, or mixed-vessel disease. *Arch. Med. Sci* 2016;12:747–53. [PubMed: 27478455]
104. Ninomiya T, Ohara T, Hirakawa Y, Yoshida D, Doi Y, Hata J, et al. Midlife and late-life blood pressure and dementia in Japanese elderly: The hisayama study. *Hypertension*. 2011;58:22–8. [PubMed: 21555680]
105. Kimm H, Lee PH, Shin YJ, Park KS, Jo J, Lee Y, et al. Mid-life and late-life vascular risk factors and dementia in Korean men and women. *Arch. Gerontol. Geriatr* 2011;52:e117–22. [PubMed: 20932588]
106. Yamada M, Mimori Y, Kasagi F, Miyachi T, Ohshita T, Sasaki H. Incidence and risks of dementia in Japanese women: Radiation Effects Research Foundation Adult Health Study. *J. Neurol. Sci* 2009;283:57–61. [PubMed: 19268313]
107. Posner HB, Tang M-X, Luchsinger J, Lantigua R, Stern Y, Mayeux R. The relationship of hypertension in the elderly to AD, vascular dementia, and cognitive function. *Neurology*. 2002;58:1175–81. [PubMed: 11971083]
108. Ninomiya T, Ohara T, Hirakawa Y, Yoshida D, Doi Y, Hata J, et al. Midlife and late-life blood pressure and dementia in Japanese elderly: The hisayama study. *Hypertension*. 2011;58:22–8. [PubMed: 21555680]
109. Leritz EC, Salat DH, Williams VJ, Schnyer DM, Rudolph JL, Lipsitz L, et al. Thickness of the human cerebral cortex is associated with metrics of cerebrovascular health in a normative sample of community dwelling older adults. *Neuroimage*. 2011;54:2659–71. [PubMed: 21035552]
110. Nagai M, Hoshida S, Ishikawa J, Shimada K, Kario K. Ambulatory blood pressure as an independent determinant of brain atrophy and cognitive function in elderly hypertension. *J. Hypertens* 2008;26:1636–41. [PubMed: 18622243]
111. Glodzik L, Mosconi L, Tsui W, de Santi S, Zinkowski R, Pirraglia E, et al. Alzheimer's disease markers, hypertension, and gray matter damage in normal elderly. *Neurobiol. Aging* 2012;33:1215–27. [PubMed: 21530003]
112. Gianaros PJ, Greer PJ, Ryan CM, Jennings JR. Higher blood pressure predicts lower regional grey matter volume: Consequences on short-term information processing. *Neuroimage*. 2006;31:754–65. [PubMed: 16488626]
113. Fjorbank MJ, Wiseman RM, Burton EJ, Saxby BK, O'Brien JT, Ford GA. Brain atrophy and white matter hyperintensity change in older adults and relationship to blood pressure. *Brain atrophy, WMH change and blood pressure*. *J. Neurol* 2007;254:713–21. [PubMed: 17446997]
114. Jennings JR, Mendelson DN, Muldoon MF, Ryan CM, Gianaros PJ, Raz N, et al. Regional grey matter shrinks in hypertensive individuals despite successful lowering of blood pressure. *J. Hum. Hypertens* 2012;26:295–305. [PubMed: 21490622]
115. Harris P, Alcantara DA, Amenta N, Lopez OL, Eiriksdóttir G, Sigurdsson S, et al. Localized measures of callosal atrophy are associated with late-life hypertension: AGES-Reykjavik Study. *Neuroimage*. 2008;43:489–96. [PubMed: 18692143]
116. Muller M, van der Graaf Y, Visseren FL, Vlek ALM, Mali WPTM, Geerlings MI, et al. Blood pressure, cerebral blood flow, and brain volumes. The SMART-MR study. *J. Hypertens* 2010;28:1498–505. [PubMed: 20453669]
117. Foster-Dingley JC, Van Der Grond J, Moonen JEF, Van Den Berg-Huijsmans AA, De Ruijter W, Van Buchem MA, et al. Lower Blood Pressure Is Associated With Smaller Subcortical Brain Volumes in Older Persons. *Am. J. Hypertens* 2015;28:1127–33. [PubMed: 25714132]
118. van Velsen EFS, Vernooij MW, Vrooman HA, van der Lugt A, Breteler MMB, Hofman A, et al. Brain cortical thickness in the general elderly population: The Rotterdam Scan Study. *Neurosci. Lett* 2013;550:189–94. [PubMed: 23831346]
119. DeCarli C, Miller BL, Swan GE, Reed T, Wolf PA, Garner J, et al. Predictors of brain morphology for the men of the NHLBI twin study. *Stroke*. 1999;30:529–36. [PubMed: 10066847]

120. Korf ESC, White LR, Scheltens P, Launer LJ. Midlife blood pressure and the risk of hippocampal atrophy: the Honolulu Asia Aging Study. *Hypertens. (Dallas, Tex. 1979)*. 2004;44:29–34.
121. * Power MC, Schneider ALC, Wruck L, Griswold M, Coker LH, Alonso A, et al. Life-course blood pressure in relation to brain volumes. *Alzheimer's Dement* 2016 8;890–9. [PubMed: 27139841] This large longitudinal study found that increasing systolic blood pressure and sustained hypertension over a 10-year period during midlife, but not late-life blood pressure, was associated with smaller late-life regional brain volumes
122. Ashby EL, Miners JS, Kehoe PG, Love S. Effects of Hypertension and Anti-Hypertensive Treatment on Amyloid- β Plaque Load and A β -Synthesizing and A β -Degrading Enzymes in Frontal Cortex. *J. Alzheimer's Dis* 2016;50:1191–203. [PubMed: 26836178]
123. Langbaum JBS, Chen K, Launer LJ, Fleisher AS, Lee W, Liu X, et al. Blood pressure is associated with higher brain amyloid burden and lower glucose metabolism in healthy late middle-age persons. *Neurobiol. Aging* 2012;33:827.e11–9.
124. Toledo JB, Toledo E, Weiner MW, Jack CR, Jagust W, Lee VMY, et al. Cardiovascular risk factors, cortisol, and amyloid- β deposition in Alzheimer's Disease Neuroimaging Initiative. *Alzheimers. Dement* 2012;8:483–9. [PubMed: 23102118]
125. Beason-Held LL, Moghekar A, Zonderman AB, Kraut MA, Resnick SM. Longitudinal changes in cerebral blood flow in the older hypertensive brain. *Stroke*. 2007;38:1766–73. [PubMed: 17510458]
126. * Gottesman RF, Coresh J, Catellier DJ, Sharrett AR, Rose KM, Coker LH, et al. Blood pressure and white-matter disease progression in a biethnic cohort: Atherosclerosis risk in communities (ARIC) study. *Stroke*. 2010;41:3–8. [PubMed: 19926835] This large community-based study found that cumulative systolic blood pressure measured over a period of 5 visits was a strong predictor white matter hyperintensity progression over a 10-year period
127. Verhaaren BFJ, Vernooij MW, De Boer R, Hofman A, Niessen WJ, Van Der Lugt A, et al. High blood pressure and cerebral white matter lesion progression in the general population. *Hypertension*. 2013;61:1354–9. [PubMed: 23529163]
128. Shams S, Martola J, Granberg T, Li X, Shams M, Fereshtehnejad SM, et al. Cerebral microbleeds: Different prevalence, topography, and risk factors depending on dementia diagnosis-the Karolinska imaging dementia study. *Am. J. Neuroradiol* 2015;36:661–6. [PubMed: 25523590]
129. Poels MMF, Zaccai K, Verwoert GC, Vernooij MW, Hofman A, Van Der Lugt A, et al. Arterial stiffness and cerebral small vessel disease: The rotterdam scan study. *Stroke*. 2012;43:2637–42. [PubMed: 22879099]
130. van Sloten TT, Protogerou AD, Henry RMA, Schram MT, Launer LJ, Stehouwer CDA. Association between arterial stiffness, cerebral small vessel disease and cognitive impairment: A systematic review and meta-analysis. *Neurosci. Biobehav. Rev* 2015 p. 121–30.
131. Dufouil C, Chalmers J, Coskun O, Besançon V, Bousser MG, Guillon P, et al. Effects of blood pressure lowering on cerebral white matter hyperintensities in patients with stroke: The PROGRESS (Perindopril Protection Against Recurrent Stroke Study) Magnetic Resonance Imaging Substudy. *Circulation*. 2005;112:1644–50. [PubMed: 16145004]
132. Launer LJ, Lewis CE, Schreiner PJ, Sidney S, Battapady H, Jacobs DR, et al. Vascular factors and multiple measures of early brain health: CARDIA brain MRI study. *PLoS One*. 2015;10:e0122138. [PubMed: 25812012]
133. Burgmans S, van Boxtel MPI, Gronenschild EHBM, Vuurman EFPM, Hofman P, Uylings HBM, et al. Multiple indicators of age-related differences in cerebral white matter and the modifying effects of hypertension. *Neuroimage*. 2010;49:2083–93. [PubMed: 19850136]
134. Maillard P, Seshadri S, Beiser A, Himali JJ, Au R, Fletcher E, et al. Effects of systolic blood pressure on white-matter integrity in young adults in the Framingham Heart Study: A cross-sectional study. *Lancet Neurol* 2012;11:1039–47. [PubMed: 23122892]
135. Gons RAR, De Laat KF, Van Norden AGW, Van Oudheusden LJB, Van Uden IWM, Norris DG, et al. Hypertension and cerebral diffusion tensor imaging in small vessel disease. *Stroke*. 2010;41:2801–6. [PubMed: 21030696]

136. Martinez-Lemus LA, Hill MA, Meininger GA. The plastic nature of the vascular wall: a continuum of remodeling events contributing to control of arteriolar diameter and structure. *Physiology*. 2009;24:45–57. [PubMed: 19196651]
137. Heagerty AM, Aalkjaer C, Bund SJ, Korsgaard N, Mulvany MJ. Small artery structure in hypertension. Dual processes of remodeling and growth. *Hypertension*. 1993;21:391–7. [PubMed: 8458640]
138. Rizzoni D, Porteri E, Castellano M, Bettoni G, Muiesan ML, Muiesan P, et al. Vascular hypertrophy and remodeling in secondary hypertension. *Hypertension*. 1996;28:785–90. [PubMed: 8901824]
139. Faraco G, Iadecola C. Hypertension: A harbinger of stroke and dementia. *Hypertension*. 2013 p. 810–7. [PubMed: 23980072]
140. Castorena-Gonzalez JA, Staiculescu MC, Foote C, Martinez-Lemus LA. Mechanisms of the inward remodeling process in resistance vessels: Is the actin cytoskeleton involved? *Microcirculation*. 2014;21:219–29. [PubMed: 24635509]
141. Scuteri A, Nilsson PM, Tzourio C, Redon J, Laurent S. Microvascular brain damage with aging and hypertension: pathophysiological consideration and clinical implications. *J. Hypertens* 2011;29:1469–77. [PubMed: 21577138]
142. Pedrinelli R, Penno G, Omo GD, Bandinelli S, Giorgi D, Bello V Di, et al. Microalbuminuria and Transcapillary Albumin Leakage in Essential Hypertension. *Hypertension*. 1999;34:491–6. [PubMed: 10489399]
143. Holmstedt CA, Turan TN, Chimowitz MI. Atherosclerotic intracranial arterial stenosis: risk factors, diagnosis, and treatment. *Lancet. Neurol* 2013;12:1106–14. [PubMed: 24135208]
144. Pantoni L Cerebral small vessel disease: from pathogenesis and clinical characteristics to therapeutic challenges. *Lancet. Neurol* 2010;9:689–701. [PubMed: 20610345]
145. Scuteri A, Morrell CH, Orrù M, Strait JB, Tarasov K V, Ferreli LAP, et al. Longitudinal perspective on the conundrum of central arterial stiffness, blood pressure, and aging. *Hypertens. (Dallas, Tex. 1979)*. 2014;64:1219–27.
146. Duschek S, Schandry R. Reduced brain perfusion and cognitive performance due to constitutional hypotension. *Clin. Auton. Res* 2007;17:69–76. [PubMed: 17106628]
147. Lenzi GL, Frackowiak RS, Jones T. Cerebral oxygen metabolism and blood flow in human cerebral ischemic infarction. *J. Cereb. Blood Flow Metab* 1982;2:321–35. [PubMed: 6980224]
148. * Muller M, Van Der Graaf Y, Visseren FL, Mali WPTM, Geerlings MI. Hypertension and longitudinal changes in cerebral blood flow: The SMART-MR study. *Ann. Neurol* 2012;71:825–33. [PubMed: 22447734] The first longitudinal study to demonstrate a relationship between hypertension, high blood pressure, and declines in cerebral blood flow using magnetic resonance angiography
149. Wong LJ, Kupferman JC, Prohovnik I, Kirkham FJ, Goodman S, Paterno K, et al. Hypertension impairs vascular reactivity in the pediatric brain. *Stroke*. 2011;42:1834–8. [PubMed: 21617149]
150. Capone C, Faraco G, Peterson JR, Coleman C, Anrather J, Milner TA, et al. Central cardiovascular circuits contribute to the neurovascular dysfunction in angiotensin II hypertension. *J. Neurosci* 2012;32:4878–86. [PubMed: 22492044]
151. Pires PW, Jackson WF, Dorrance AM. Regulation of myogenic tone and structure of parenchymal arterioles by hypertension and the mineralocorticoid receptor. *Am. J. Physiol. - Hear. Circ. Physiol* 2015;309:H127–36.
152. Matsushita K, Kuriyama Y, Nagatsuka K, Nakamura M, Sawada T, Omae T. Periventricular white matter lucency and cerebral blood flow autoregulation in hypertensive patients. *Hypertension*. 1994;23:565–8. [PubMed: 8175162]
153. Wang T, Li Y, Guo X, Huang D, Ma L, Wang DJJ, et al. Reduced perfusion in normal-appearing white matter in mild to moderate hypertension as revealed by 3D pseudocontinuous arterial spin labeling. *J. Magn. Reson. Imaging*. 2016;43:635–43. [PubMed: 26256700]
154. Heiss W-D. The Ischemic Penumbra: Correlates in Imaging and Implications for Treatment of Ischemic Stroke. *Cerebrovasc. Dis* 2011;32:307–20. [PubMed: 21921593]
155. Iadecola C Neurovascular regulation in the normal brain and in Alzheimer's disease. *Nat. Rev. Neurosci* 2004;5:347–60. [PubMed: 15100718]

156. Koike MA, Green KN, Blurton-Jones M, Laferla FM. Oligemic hypoperfusion differentially affects tau and amyloid- β . *Am. J. Pathol* 2010;177:300–10. [PubMed: 20472896]
157. Gentile MT, Poulet R, Pardo A Di, Cifelli G, Maffei A, Vecchione C, et al. Beta-Amyloid deposition in brain is enhanced in mouse models of arterial hypertension. *Neurobiol. Aging* 2009;30:222–8. [PubMed: 17673335]
158. Wang X, Xing A, Xu C, Cai Q, Liu H, Li L. Cerebrovascular hypoperfusion induces spatial memory impairment, synaptic changes, and amyloid- β oligomerization in rats. *J. Alzheimer's Dis* 2010;21:813–22. [PubMed: 20634588]
159. Walsh DM, Klyubin I, Fadeeva J V, Cullen WK, Anwyl R, Wolfe MS, et al. Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature*. 2002;416:535–9. [PubMed: 11932745]
160. Li L, Zhang X, Yang D, Luo G, Chen S, Le W. Hypoxia increases A β generation by altering beta - and gamma-cleavage of APP. *Neurobiol. Aging* 2009;30:1091–8. [PubMed: 18063223]
161. * Carnevale D, Mascio G, D'Andrea I, Fardella V, Bell RD, Branchi I, et al. Hypertension induces brain β -amyloid accumulation, cognitive impairment, and memory deterioration through activation of receptor for advanced glycation end products in brain vasculature. *Hypertens*. 2012;60:188–97. This study provides convincing evidence for a link between hypertension and increased Alzheimer's disease pathology that occurs as a result of an upregulation of receptor for advanced glycation end products (RAGE) in cerebral vessels
162. Girouard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease. *J. Appl. Physiol* 2006;100:328–35. [PubMed: 16357086]
163. Kazama K, Wang G, Frys K, Anrather J, Iadecola C. Angiotensin II attenuates functional hyperemia in the mouse somatosensory cortex. *Am. J. Physiol. Heart Circ. Physiol*. 2003;285:H1890–9. [PubMed: 12907423]
164. Xia W, Rao H, Spaeth AM, Huang R, Tian S, Cai R, et al. Blood Pressure is Associated With Cerebral Blood Flow Alterations in Patients With T2DM as Revealed by Perfusion Functional MRI. *Medicine (Baltimore)*. 2015;94:e2231. [PubMed: 26632913]
165. Jennings JR, Muldoon MF, Ryan C, Price JC, Greer P, Sutton-Tyrrell K, et al. Reduced cerebral blood flow response and compensation among patients with untreated hypertension. *Neurology*. 2005;64:1358–65. [PubMed: 15851723]
166. Capone C, Faraco G, Park L, Cao X, Davisson RL, Iadecola C. The cerebrovascular dysfunction induced by slow pressor doses of angiotensin II precedes the development of hypertension. *Am. J. Physiol. Heart Circ. Physiol* 2011;300:H397–407. [PubMed: 20971763]
167. Laplante MA, Wu R, Moreau P, De Champlain J. Endothelin mediates superoxide production in angiotensin II-induced hypertension in rats. *Free Radic. Biol. Med* 2005;38:589–96. [PubMed: 15683715]
168. Duron E, Hanon O. Antihypertensive treatments, cognitive decline, and dementia. *J. Alzheimer's Dis* 2010 p. 903–14.
169. ** McGuinness B, Todd S, Passmore P, Bullock R. Blood pressure lowering in patients without prior cerebrovascular disease for prevention of cognitive impairment and dementia. *Cochrane Database Syst. Rev* 2009. p. CD004034. This is a 2009 Cochrane review of the effectiveness of blood pressure lowering for prevention of cognitive impairment and dementia
170. Tzourio C, Anderson C, Chapman N, Woodward M, Neal B, MacMahon S, et al. Effects of blood pressure lowering with perindopril and indapamide therapy on dementia and cognitive decline in patients with cerebrovascular disease. *Arch. Intern. Med* 2003;163:1069–75. [PubMed: 12742805]
171. Forette F, Seux ML, Staessen JA, Thijs L, Birkenhäger WH, Babarskiene MR, et al. Prevention of dementia in randomised double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) trial. *Lancet*. 1998;352:1347–51. [PubMed: 9802273]
172. Bosch J, Yusuf S, Pogue J, Sleight P, Lonn E, Rangoonwala B, et al. Use of ramipril in preventing stroke: double blind randomised trial. *BMJ*. 2002;324:699. [PubMed: 11909785]
173. Anderson C, Teo K, Gao P, Arima H, Dans A, Unger T, et al. Renin-angiotensin system blockade and cognitive function in patients at high risk of cardiovascular disease: Analysis of data from the ONTARGET and TRANSCEND studies. *Lancet Neurol* 2011;10:43–53. [PubMed: 20980201]

174. SHEP. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. Final results of the Systolic Hypertension in the Elderly Program (SHEP). SHEP Cooperative Research Group. *JAMA* 1991;265:3255–64. [PubMed: 2046107]
175. Lithell H, Hansson L, Skoog I, Elmfeldt D, Hofman A, Olofsson B, et al. The Study on Cognition and Prognosis in the Elderly (SCOPE): principal results of a randomized double-blind intervention trial. *J. Hypertens* 2003;21:875–86. [PubMed: 12714861]
176. Peters R, Beckett N, Forette F, Tuomilehto J, Clarke R, Ritchie C, et al. Incident dementia and blood pressure lowering in the Hypertension in the Very Elderly Trial cognitive function assessment (HYVET-COG): a double-blind, placebo controlled trial. *Lancet Neurol* 2008;7:683–9. [PubMed: 18614402]
177. Prince MJ, Bird AS, Blizard RA, Mann AH. Is the cognitive function of older patients affected by antihypertensive treatment? Results from 54 months of the Medical Research Council's trial of hypertension in older adults. *BMJ*. 1996;312:801–5. [PubMed: 8608285]
178. Wright JW, Stubley L, Pederson ES, Kramár EA, Hanesworth JM, Harding JW. Contributions of the brain angiotensin IV-AT4 receptor subtype system to spatial learning. *J. Neurosci* 1999;19:3952–61. [PubMed: 10234025]
179. Takeda S, Sato N, Takeuchi D, Kurinami H, Shinohara M, Niisato K, et al. Angiotensin receptor blocker prevented beta-amyloid-induced cognitive impairment associated with recovery of neurovascular coupling. *Hypertension*. 2009;54:1345–52. [PubMed: 19805638]
180. Marpillat NL, Macquin-Mavier I, Tropeano A-I, Bachoud-Levi A-C, Maison P. Antihypertensive classes, cognitive decline and incidence of dementia: a network meta-analysis. *J. Hypertens* 2013;31:1073–82. [PubMed: 23552124]
181. Wharton W, Stein JH, Korcarz C, Sachs J, Olson SR, Zetterberg H, et al. The effects of ramipril in individuals at risk for Alzheimer's disease: Results of a pilot clinical trial. *J. Alzheimer's Dis* 2012;32:147–56. [PubMed: 22776970]
182. Hughes TM, Sink KM. Hypertension and Its Role in Cognitive Function: Current Evidence and Challenges for the Future. *Am. J. Hypertens* 2016;29:149–57. [PubMed: 26563965]
183. Bossers WJR, van der Woude LHV, Boersma F, Scherder EJA, van Heuvelen MJG. Recommended measures for the assessment of cognitive and physical performance in older patients with dementia: a systematic review. *Dement. Geriatr. Cogn. Dis. Extra* 2012;2:589–609. [PubMed: 23341825]
184. Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: a Frontal Assessment Battery at bedside. *Neurology*. 2000;55:1621–6. [PubMed: 11113214]
185. * Ambrosius WT, Sink KM, Foy CG, Berlowitz DR, Cheung AK, Cushman WC, et al. The design and rationale of a multicenter clinical trial comparing two strategies for control of systolic blood pressure: the Systolic Blood Pressure Intervention Trial (SPRINT). *Clin. Trials* 2014;11:532–46. [PubMed: 24902920] Outlines the design and rationale for the parent study of the recent Systolic Blood Pressure Intervention Trial: Memory and Cognition in Decreased Hypertension (SPRINT-MIND) trial
186. SPRINT Research Group, Wright JT, Williamson JD, Whelton PK, Snyder JK, Sink KM, et al. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N. Engl. J. Med* 2015;373:2103–16. [PubMed: 26551272]
187. McNiece KL, Poffenbarger TS, Turner JL, Franco KD, Sorof JM, Portman RJ. Prevalence of Hypertension and Pre-Hypertension among Adolescents. *J. Pediatr* 2007;150:640–644.e1. [PubMed: 17517252]
188. Lande MB, Kupferman JC. Cognitive function in hypertensive children. *Curr. Hypertens. Rep* 2015;17:508. [PubMed: 25432900]
189. Tsao CW, Himali JJ, Beiser AS, Larson MG, DeCarli C, Vasan RS, et al. Association of arterial stiffness with progression of subclinical brain and cognitive disease. *Neurology*. 2016;86:619–26. [PubMed: 26791155]
190. Waldstein SR, Rice SC, Thayer JF, Najjar SS, Scuteri A, Zonderman AB. Pulse pressure and pulse wave velocity are related to cognitive decline in the Baltimore Longitudinal Study of Aging. *Hypertens. (Dallas, Tex. 1979)*. 2008;51:99–104.

191. Liao D, Cooper L, Cai J, Toole J, Bryan N, Burke G, et al. The prevalence and severity of white matter lesions, their relationship with age, ethnicity, gender, and cardiovascular disease risk factors: the ARIC Study. *Neuroepidemiology*. 1997;16:149–62. [PubMed: 9159770]
192. Nation DA, Edmonds EC, Bangen KJ, Delano-Wood L, Scanlon BK, Han SD, et al. Pulse Pressure in Relation to Tau-Mediated Neurodegeneration, Cerebral Amyloidosis, and Progression to Dementia in Very Old Adults. *JAMA Neurol* 2015;72:546. [PubMed: 25822631]
193. Webb AJS, Simoni M, Mazzucco S, Kuker W, Schulz U, Rothwell PM. Increased cerebral arterial pulsatility in patients with leukoaraiosis: Arterial stiffness enhances transmission of aortic pulsatility. *Stroke*. 2012;43:2631–6. [PubMed: 22923446]
194. Swan GE, DeCarli C, Miller BL, Reed T, Wolf PA, Jack LM, et al. Association of midlife blood pressure to late-life cognitive decline and brain morphology. *Neurology*. 1998;51:986–93. [PubMed: 9781518]
195. Singh-Manoux A, Marmot M. High blood pressure was associated with cognitive function in middle-age in the Whitehall II study. *J. Clin. Epidemiol* 2005;58:1308–15. [PubMed: 16291476]
196. Gillett SR, Thacker EL, Letter AJ, McClure LA, Wadley VG, Unverzagt FW, et al. Correlates of Incident Cognitive Impairment in the REasons for Geographic and Racial Differences in Stroke (REGARDS) Study. *Clin. Neuropsychol* 2015;29:466–86. [PubMed: 25978342]
197. Wolf PA, Beiser A, Elias MF, Au R, Vasan RS, Seshadri S. Relation of obesity to cognitive function: importance of central obesity and synergistic influence of concomitant hypertension. The Framingham Heart Study. *Curr. Alzheimer Res* 2007;4:111–6. [PubMed: 17430232]
198. Pavlik VN, Hyman DJ, Doody R. Cardiovascular risk factors and cognitive function in adults 30–59 years of age (NHANES III). *Neuroepidemiology*. 2005;24:42–50. [PubMed: 15459509]
199. Tzourio C, Dufouil C, Ducimetiere P, Alperovitch A. Cognitive decline in individuals with high blood pressure: A longitudinal study in the elderly. *Neurology*. 1999;53:1948–52. [PubMed: 10599763]
200. Knopman D, Boland L, Mosley T, Howard G, Liao D, Szklo M, et al. Cardiovascular risk factors and cognitive decline in middle-aged adults. *Neurology*. 2001;56:42–8. [PubMed: 11148234]
201. Farmer ME, Kittner SJ, Abbott RD, Wolz MM, Wolf PA, White LR. Longitudinally measured blood pressure, antihypertensive medication use, and cognitive performance: the Framingham Study. *J. Clin. Epidemiol* 1990;43:475–80. [PubMed: 2324788]
202. Cerhan JR, Folsom AR, Mortimer JA, Shahar E, Knopman DS, McGovern PG, et al. Correlates of cognitive function in middle-aged adults. Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Gerontology*. 1998;44:95–105. [PubMed: 9523221]
203. Guo Z, Fratiglioni L, Winblad B, Viitanen M. Blood pressure and performance on the Mini-Mental State Examination in the very old. Cross-sectional and longitudinal data from the Kungsholmen Project. *Am. J. Epidemiol*. 1997;145:1106–13. [PubMed: 9199540]
204. Wadley VG, McClure LA, Howard VJ, Unverzagt FW, Go RC, Moy CS, et al. Cognitive status, stroke symptom reports, and modifiable risk factors among individuals with no diagnosis of stroke or transient ischemic attack in the REasons for Geographic and Racial Differences in Stroke (REGARDS) study. *Stroke*. 2007;38:1143–7. [PubMed: 17322077]
205. Elias MF, Elias PK, Sullivan LM, Wolf PA, D'Agostino RB. Lower cognitive function in the presence of obesity and hypertension: the Framingham heart study. *Int. J. Obes. Relat. Metab. Disord* 2003;27:260–8. [PubMed: 12587008]
206. Cacciatore F, Abete P, Ferrara N, Paolisso G, Amato L, Canonico S, et al. The role of blood pressure in cognitive impairment in an elderly population. *Osservatorio Geriatrico Campano Group. J. Hypertens* 1997;15:135–42. [PubMed: 9469788]
207. Hebert LE, Scherr PA, Bennett DA, Bienias JL, Wilson RS, Morris MC, et al. Blood pressure and late-life cognitive function change: a biracial longitudinal population study. *Neurology*. 2004;62:2021–4. [PubMed: 15184608]
208. Di Carlo A, Baldereschi M, Amaducci L, Maggi S, Grigoletto F, Scarlato G, et al. Cognitive impairment without dementia in older people: prevalence, vascular risk factors, impact on disability. The Italian Longitudinal Study on Aging. *J. Am. Geriatr. Soc* 2000;48:775–82. [PubMed: 10894316]

209. André-Petersson L, Hagberg B, Janzon L, Steen G. A Comparison of Cognitive Ability in Normotensive and Hypertensive 68-YearOld Men: Results from Population Study “Men Born in 1914,” in Malmö, Sweden. *Exp. Aging Res* 2001;27:319–40. [PubMed: 11681196]
210. Haan MN, Shemanski L, Jagust WJ, Manolio TA, Kuller L. The role of APOE epsilon4 in modulating effects of other risk factors for cognitive decline in elderly persons. *JAMA*. 1999;282:40–6. [PubMed: 10404910]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

Epidemiologic studies of early- and midlife hypertension and cognition

Study	Study Design	Duration	Age at BP Assessment	BP Measure	Age at Cognitive Assessment	Cognitive Domains Assessed	Major Findings
NHANES III, USA [40]	Cross-sectional	<i>n</i> = 5,077	6-16	SBP, DBP	6-16	Arithmetic Reading Visuospatial Working memory	Elevated SBP, but not DBP, was associated with poorer working memory.
Normative Aging Study, USA [18]	Cross-temporal ^a	29 years <i>n</i> = 758	37 (5) (BP measured every 3-5 years for up to 44 years)	HTN	66 (5)	Multi-domain composite score	HTN at any point during follow-up and greater duration since onset of HTN were associated with lower cognitive functioning later in life, independent of age at onset.
NHLBI Twin Study, USA [194]	Cross-temporal ^a	25 years Longitudinal 10 years <i>n</i> = 392	43-53	SBP, DBP	T1: 59-69 T2: 68-79	MMSE Processing speed Verbal fluency Visual memory	Elevated SBP at baseline was associated with declines in processing speed over a period of 10 years.
Honolulu-Asia Aging Study, USA [12]	Cross-temporal ^a	25 years <i>n</i> = 3,735	53 (5)	SBP, DBP	78 (5)	Cognitive Abilities Screening Instrument (CASI)	Elevated SBP at baseline was associated with poorer cognitive functioning in late-life.
Xi'an, China [30]	Cross-sectional	<i>n</i> = 1,799	40-85	SBP, DBP, MAP	40-85	MMSE	Elevated SBP, DBP, and MAP were associated with cognitive impairment among 40-60 year-olds, but this was not the case for older participants.
Whitehall II, UK [195]	Cross-temporal ^a	12 years <i>n</i> = 5,838	44 (6)	SBP, DBP	56	Memory Reasoning Verbal fluency Vocabulary	Elevated SBP at baseline was associated with poorer baseline memory and reduced verbal fluency at follow-up, especially among women.
REGARDS, USA [196]	Cross-temporal ^a	3 years <i>n</i> = 17,630	64 (9)	HTN	67	Multi-domain composite score	HTN was not associated with the development of cognitive impairment over a span of 40 months.
Framingham Heart Study, USA [197]	Cross-temporal ^a	12 years <i>n</i> = 1,814	40-69	HTN	T1: 52 (8) T2: 64	Executive function Memory Visuospatial	HTN was associated with worse performance on measures of executive functioning and visual memory.
NHANES III, USA [198]	Cross-sectional	<i>n</i> = 3,270	30-59	HTN	30-59	Processing speed Reaction time Working memory	HTN and DM, but not HTN alone, was associated with worse reaction time, processing speed, and working memory.
EVA Study Group, France [199]	Longitudinal	4 years <i>n</i> = 1,373	59-71	HTN	T1: 59-71 T2: 63-75	MMSE	Baseline HTN was associated greater MMSE declines over 4 years. This relationship was stronger among

Study	Study Design Duration <i>n</i>	Age at BP Assessment	BP Measure	Age at Cognitive Assessment	Cognitive Domains Assessed	Major Findings
ARIC, USA [200]	Longitudinal 6 years <i>n</i> = 10,963	47-70	HTN	T1: 47-70 T2: 53-76	Memory Processing speed Verbal fluency	participants who were untreated for hypertension. Baseline HTN was associated with greater decline in processing speed over a 6-year period.
ARIC, USA [14]	Longitudinal 14 years <i>n</i> = 12,702	59 (4)	HTN	T1: 59 (4) T2: 62 T3: 65 T4: 73	Memory Processing speed Verbal fluency	Baseline HTN was associated with greater declines in verbal fluency over a 14-year period.
ARIC, USA [13]	Longitudinal 20 years <i>n</i> = 13,476	45-64	Pre-HTN, HTN	T1: 48-67 T2: 54-73 T3: 70-89	Processing speed Verbal fluency Verbal memory	Baseline HTN was associated with greater declines in processing speed, verbal fluency, and a global composite score of cognitive functioning over a 20-year period.
Western Collaborative Group Study, USA [17]	Cross-temporal ^a 25-30 years <i>n</i> = 717	39-59 (BP measured approximately 10 times over 30 years)	SBP	75 (4)	Executive function Memory Processing speed	Compared to participants who maintained a normal SBP over 30 years, participants who had persistently high SBP had poorer verbal memory in late-life. Participants who displayed a significant decrease in SBP over 30 years performed more poorly on measure of processing speed.
Male Cohort in Uppsala, Sweden [11]	Cross-temporal ^a 20 years <i>n</i> = 999	50	SBP, DBP	72	Multi-domain composite score	Elevated DBP at baseline was associated with reduced cognitive functioning in late-life. This relationship was stronger among participants who were untreated for hypertension.
Male Cohort in Uppsala, Sweden [10]	Cross-temporal ^a 20 years <i>n</i> = 502	50	SBP, DBP	72 (1)	Memory Processing speed Verbal fluency Visuospatial Working memory	Elevated DBP at baseline was associated with poorer performance on measures of working memory, processing speed, and verbal fluency in late-life. Participants with DBP 70mg Hg at baseline demonstrated highest levels cognitive functioning in late-life.
Framingham, USA [201]	Cross-temporal ^a 28 years <i>n</i> = 1,993	27-87 (BP measured biennially over 28 years)	HTN, SBP, DBP	55-89	Attention Language Memory Visuospatial	Among participants untreated for hypertension, the proportion of visits during which HTN was present and the average SBP and DBP were inversely associated with cognitive functioning.
ARIC, USA [202]	Cross-sectional <i>n</i> = 13,840	45-69	HTN	45-69	Processing speed Verbal fluency Verbal memory	HTN among women, but not men, was associated with poorer performance on all cognitive measures.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

ARIC = Atherosclerosis Risk in Communities; BP = blood pressure; DBP = diastolic blood pressure; EVA = Epidemiology of Vascular Aging; HTN = hypertension; MAP = mean arterial pressure; MMSE = Mini Mental Status Exam; NHANES III = National Health and Nutrition Examination Survey; NHLBI = National Heart, Lung, and Blood Institute; REGARDS = Reasons for Geographic and Racial Differences in Stroke; SBP = systolic blood pressure

^aCross temporal: a study design in which the exposure variable (e.g., hypertension) and the outcome variable (e.g., cognition) are measured distinct time points.

Table 2.

Epidemiologic studies of late-life hypertension and cognition

Study	Study Design Duration <i>n</i>	Age at BP Assessment	BP Measure	Age at Cognitive Assessment	Cognitive Domains Assessed	Major Findings
COGNIPRES, Spain [21]	Cross-sectional <i>n</i> = 1,579	71 (7)	HTN, SBP, DBP, PP, MAP	71 (7)	MMSE	HTN and antihypertensive medication noncompliance were associated with lower MMSE scores.
Kungsholmen Project, Sweden [203]	Longitudinal 3 years <i>n</i> = 1,736	75-101	SBP, DBP	T1: 75-101 T2: 78-104	MMSE	Higher SBP and DBP at baseline were associated with better MMSE score at baseline and 3-year follow-up. Baseline SBP < 130 mmHg was associated with increased risk of cognitive impairment at follow-up.
REGARDS, USA [204]	Cross-sectional <i>n</i> = 14,566	65 (9)	HTN	65 (9)	Six-item Screener	HTN was not associated with risk of cognitive impairment.
Framingham Heart Study, USA [23]	Cross-temporal ^a 13 years <i>n</i> = 1,702	67 (8)	HTN	80	Abstract reasoning Language Memory Verbal fluency Working memory	Baseline HTN was associated with poorer working memory, visual memory, and verbal memory at follow-up among participants not on antihypertensive medication.
Framingham Heart Study, USA [205]	Cross-temporal ^a 4-6 years <i>n</i> = 1,423	66 (7)	HTN	71	Abstract reasoning Executive function Memory	Baseline HTN was associated with greater memory impairment 4-6 years later in men, but not in women.
Northern Manhattan Study, USA [25]	Longitudinal 6 years <i>n</i> = 4,337	76 (6)	HTN	T1: 78 T2: 80 T3: 81	Executive function Language Memory	Baseline HTN was associated with declines in executive functioning, but not memory or language.
REGARDS USA [22]	Cross-sectional <i>n</i> = 19,836	65 (10)	SBP, DBP, PP	65 (10)	Six-item Screener	Higher DBP was associated with greater risk of cognitive impairment.
Indo-US Cross National Dementia Epidemiology Study, India/USA [27]	Cross-sectional <i>n</i> = 4810 <i>n</i> = 636	67 (7) 82 (4)	SBP, DBP	67 (7) 82 (4)	MMSE	Higher SBP and DBP were associated with increased risk for cognitive impairment in the younger Indian cohort. No association between BP and cognitive impairment was found in the older American cohort.
Osservatorio Geriatrico Regione Campania, Italy [206]	Cross-sectional <i>n</i> = 1,229	74 (6)	SBP, DBP	74 (6)	MMSE	Higher DBP, but not SBP, was associated with cognitive impairment in participants > 75, but not in participants 65-74 years of age.
Baltimore Longitudinal Study of Ageing, USA [29]	Longitudinal 11 years <i>n</i> = 847	71 (9)	SBP, DBP	82	Attention Executive function Memory Naming Processing speed	High SBP was associated with memory declines among older participants. Cross-sectional analyses demonstrated that both high and low diastolic BP were associated with poorer executive functioning, processing speed, and naming among participant groups.

Study	Study Design	Duration	Age at BP Assessment	BP Measure	Age at Cognitive Assessment	Cognitive Domains Assessed	Major Findings
Chicago Health and Aging, USA [28]	Cross-sectional	$n = 5,816$	65-104	SBP, DBP	65-104	MMSE	Participants with SBP < 100 mm Hg and SBP > 140 mm Hg had lower MMSE scores.
Chicago Health and Aging, USA [207]	Longitudinal 6 years	$n = 4,284$	74 (6)	SBP, DBP	80	MMSE Memory Processing speed	BP was not associated with cognitive change over the span of 6 years.
The Italian Longitudinal Study on Aging, Italy [208]	Cross-sectional	$n = 3,425$	65-84	HTN	65-84	MMSE	Hypertension was not associated with MMSE score.
Men Born in 1914, Sweden [209]	Cross-sectional	$n = 500$	68	HTN	68	Processing speed Verbal abilities Visual memory Visuospatial	HTN (SBP 140-159 mm Hg) was associated with better visuospatial and verbal abilities. Severe HTN (SBP 180 mm Hg) was associated with poorer performance on measures of memory and processing speed.
Cardiovascular Health Study, USA [210]	Longitudinal 7 years	$n = 5,888$	65	SBP	72	Modified MMSE Processing speed	SBP Elevated SBP was associated with a decline in MMSE and processing speed over a period of 7 years.
East Boston cohort study, USA [32]	Longitudinal 9 years	$n = 3,657$	74 (6)	SBP, DBP	83	Memory SPMSQ	A U-shaped relationship between SBP and cognition was found whereby SBP < 130mm Hg or 160mm Hg was associated with a higher rate of errors on a mental status questionnaire (SPMSQ).
Duke Population Studies of the Elderly, USA [33]	Longitudinal 3 years	$n = 3,202$	73 (6)	SBP, DBP	76	SPMSQ	Among white participants, a U-shaped relationship between SBP cognitive decline was found whereby SBP < 110 mm Hg and SBP > 165 mm Hg was associated with 3-year cognitive decline. No association between BP and cognition was found in black participants.
East Boston Study, USA [26]	Cross-sectional	$n = 3,627$	65	SBP, DBP, HTN	65	Attention Memory	BP was not associated with cognitive functioning.

BP = blood pressure; COGNIPRES = Cognitive function and blood pressure control; DBP = diastolic blood pressure; HTN = hypertension; MAP = mean arterial pressure; MMSE = Mini Mental Status Exam; PP = pulse pressure; REGARDS = Reasons for Geographic and Racial Differences in Stroke; SBP = systolic blood pressure; SPMSQ = Short Portable Mental Status Questionnaire

^aCross temporal: a study design in which the exposure variable (e.g., hypertension) and the outcome variable (e.g., cognition) are measured distinct time points.