Published in final edited form as:

Osteoporos Int. 2018 October ; 29(10): 2275–2281. doi:10.1007/s00198-018-4626-2.

Breech presentation is associated with lower bone mass and area: findings from the Southampton Women's Survey

Alex Ireland¹, Sarah R Crozier², Alexander E P Heazell³, Kate Ward², Keith M Godfrey^{2,4}, Hazel M Inskip^{2,4}, Cyrus Cooper^{2,4,5}, and Nicholas C Harvey^{2,4}

¹School of Healthcare Science, Manchester Metropolitan University, Manchester, UK

²MRC Lifecourse Epidemiology Unit, University of Southampton, UK

³Division of Developmental Biology and Medicine, Faculty of Biology, Medicine and Health, University of Manchester, UK

⁴NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

⁵NIHR Oxford Biomedical Research Centre, University of Oxford, UK

Abstract

Purpose—Breech presentation is associated with altered joint shape and hip dysplasias, but effects on bone mineral content (BMC), area (BA) and density (BMD) are unknown.

Methods—In the prospective Southampton Women's Survey mother-offspring cohort, whole-body bone outcomes were measured using dual-energy X-ray absorptiometry (DXA) in 1430 offspring, as neonates (mean age 6 days, $n=965$, 39 with a breech presentation at birth) and/or at age 4.1 years ($n=999$, 39 breech). Hip and spine bone outcomes were also measured at age 4 years.

Results—Neonates with breech presentation had 4.2g lower whole-body BMC (95%CI -7.4g to -0.9g, $P=0.012$), 5.9cm² lower BA (-10.8 cm² to -1.0 cm², $P=0.019$), but BMD was similar between groups (mean difference -0.007g/cm², -0.016g/cm² to 0.002g/cm², $P=0.146$) adjusting for sex, maternal smoking, gestational diabetes, mode of delivery, social class, parity, ethnicity, age at scan, birthweight, gestational age and crown-heel length. There were no associations between breech presentation and whole-body outcomes at age 4 years, but, in similarly-adjusted models, regional DXA (not available in infants) showed that breech presentation was associated with lower hip BMC (-0.51g, -0.98g to -0.04g, $P=0.034$) and BA (-0.67cm², -1.28cm² to -0.07cm², $P=0.03$) but not with BMD (-0.009g, -0.029g to 0.012g, $P=0.408$), or spine outcomes.

Conclusions—These results suggest that breech presentation is associated with lower neonatal whole-body BMC and BA, which may relate to altered prenatal loading in babies occupying a breech position; these differences did not persist into later childhood. Modest differences in 4-year hip BMC and BA require further investigation.

Corresponding Author: Professor Nicholas Harvey, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, SO16 6YD, UK, +44 23 8077 7624, nch@mrc.soton.ac.uk.

Author Disclosures

Alex Ireland, Sarah Crozier, Alexander Heazell, Kate Ward, Keith M Godfrey, Hazel M Inskip, Cyrus Cooper and Nicholas Harvey declare that they have no conflict of interest.

Keywords

Osteoporosis; epidemiology; fetal growth; bone mass; pregnancy

Introduction

The prenatal period has a strong and persisting influence on skeletal development and health. Maternal factors such as smoking, body composition, parity and activity influence accrual of bone mass during pregnancy [1, 2]. In addition, factors intrinsic to or influenced by pregnancy including birthweight and gestational age [2] and placental size [3] are also determinants of neonatal bone mass. Birthweight in particular remains predictive of bone mass throughout childhood [4] and into older age [5], whilst preeclampsia has been associated with reduced hip BMD in adolescence [6]. Therefore, it is important to identify predictors of fetal bone mineral accrual during pregnancy to develop preventative strategies against future osteoporosis and fractures [7]. However, whilst mechanical loading via physical activity and exercise has a dominating influence on bone area/mass throughout early childhood [8], adolescence [9–11] and into old age [12–14], the influence of fetal presentation on bone development remains little explored.

Breech presentation (where the fetus is lying bottom-down in pregnancy), may be considered as a natural model of altered fetal loading and is associated with restricted fetal movement [15]. Breech presentation is common throughout pregnancy, evident in ~50% of fetuses at 25 weeks' gestation [16] falling to approximately ~5% at birth [17]. Breech presentation results in restriction of lower limb movements in the third trimester, particularly at the hip joint [15]. This is particularly important as the mechanical loading of the skeleton caused by fetal movements is greatest in late pregnancy [18]. However, it has been proposed that poor fetal movement may also be responsible for the failure to attain a cephalic position [19] *i.e.* reduced movements in breech fetuses may not solely be attributable to fetal position [20]. This is supported by a higher frequency of breech presentation in children with conditions affecting motor development such as cerebral palsy [21]; the incidence increases with disease severity [19]. Given the importance of fetal skeletal loading as demonstrated in animal [22] and *in silico* [23] models, it is not surprising that breech presentation has consequences for skeletal development. Breech presentation at birth is associated with a ten-fold higher incidence of developmental hip dysplasias [24], whilst femoral anteversion is 10° higher (equivalent to 1.25SD) in breech rather than cephalic presentation [25]. However, the effects of breech presentation on bone mass accrual during gestation and early life have not been explored. Tibial bone ultrasound speed of sound (an indirect indicator of bone quality) was lower in breech than cephalic presentation [26], but clinical measures of bone mass, area and density have not previously been assessed – nor have longer-term associations been examined.

In this study we compared bone mass, area and density at birth and age four years in children born with breech or cephalic presentation in a large prospective mother-offspring cohort – the Southampton Women's Survey (SWS). It was hypothesised that breech presentation would be associated with deficits in bone outcomes at birth and early

childhood, which would not be explained by lower gestational age or birthweight in this group.

Methods

The SWS recruited 12,583 women living within the city of Southampton in southern England when they were not pregnant. Women in the cohort who subsequently became pregnant were invited to complete a lifestyle assessment questionnaire at 11 and 34 weeks' gestation, from which details of maternal smoking, ethnicity, parity, gestational diabetes and social class were extracted. Only singleton pregnancies, and the first pregnancy per study mother were included. At birth, gestational age, mode of delivery and fetal presentation were recorded, as were crown-heel length measured using a neonatometer (CMS Ltd, UK) and birthweight using calibrated digital scales (Seca, UK). Within two weeks of birth and following maternal consent, total body dual-energy X-ray absorptiometry (DXA) scans were taken using a Lunar DPX-L scanner (GE Corporation, Madison, WA, USA) using pediatric software (pediatric small scan mode v4.7c). During the neonatal scan visit infants were fed if necessary, undressed completely and then swaddled in a standard towel. They were placed in a standard position on a waterproof sheet, and kept in position using rice bags placed over the bottom end of the towel. At age 4 years, height was measured using a calibrated stadiometer and weight using calibrated digital scales (Seca, UK); total body and dedicated hip and spine scans were performed using a Discovery DXA scanner (Hologic Inc., Bedford, MA, USA). The scan machines underwent a daily quality assessment, alongside a weekly calibration against a water phantom. To encourage compliance, the scanner bed was covered with a sheet with appropriate coloured cartoons was laid on the couch prior to scan positioning. To help reduce movement artefact, children were shown a suitable cartoon on DVD. From the neonatal total body scan, measurements of bone mineral content (BMC), bone area (BA) and bone mineral density (BMD) for the whole body including the head were obtained. Size-adjusted or SA BMC (BMC adjusted for height, weight and BA) was also assessed to give an estimate of volumetric BMD. At age 4 years, BMC, BA, BMD were recorded for whole body (less head), hip and spine scans, and SA BMC was assessed. Scans were assessed by two independent operators; 31 neonatal scans and 32 four year scans were excluded from analysis due to movement artefacts. Babies with a known congenital anomaly ($n=7$), and children with a fetal position other than breech or cephalic *e.g.* transverse ($n=14$) were excluded from analysis. The study was approved by Southampton and Southwest Hampshire Local Research Ethics Committee (approval numbers 267/97, 307/97, 153/99 and 005/03/t) and parental informed consent was given at both scan timepoints.

Statistical analysis

A complete case analysis was performed with no imputation, resulting in a cohort of 1430 participants (745 male). The neonatal and 4 year cohorts partially overlapped such that 534 individuals had complete data at both timepoints, and the maximum number of participants was used at either time point to optimise statistical power. 965 infants (498 male) had complete data at birth and were included in neonatal analyses; 999 children (515 male) had complete data at age four years. In a sensitivity analysis, the cohort was limited to the offspring with data at both time points. Crown-heel length and birthweight were converted to

Z-scores using WHO Child Growth Standards [27]. Differences in cohort characteristics by presentation type (conventional/breech) were assessed by Fisher's exact test, χ^2 tests and t-tests for binary, categorical and continuous variables respectively. Associations between presentation type and bone outcomes were assessed with multiple linear regression models using the R statistical environment (version 3.1.2, www.r-project.org). There was no evidence of sex interactions in relation to presentation mode and its associations with bone outcomes, so separate analyses were not conducted by sex. A minimal Model 1 was adjusted only for sex. In Model 2, potential confounders known to be associated with both breech presentation and bone outcomes were included. In Model 3, potential mediating factors relating to length of gestation and body size were included, in order to identify the direct association between breech presentation and bone outcomes independent of these known differences between breech and cephalic births. The structure of model predictors was as follows:

Model 1 (Minimal): Presentation + Sex (four-year scans additionally adjusted for height)

Model 2 (Confounders): Model 1 + Maternal Smoking + Gestational Diabetes + Delivery Mode + Social Class + Parity + Ethnicity + Age at Scan

Model 3 (Mediators): Model 2 + Birthweight + Gestational Age + Crown-Heel Length (neonatal outcomes) or Height at Time of Scan (four-year outcomes)

Model residuals were checked for heteroscedasticity, and variance inflation factors (VIFs) were used to check for multicollinearity. Outcomes are presented as means and 95% confidence intervals, unless otherwise stated.

Results

Of the 965 participants with neonatal scans, 39 (19 male) had a breech presentation at birth; 39 (20 male) of the 999 children with four year scans had a breech presentation at birth (Table 1). Breech presentation was associated with lower birthweight, and shorter crown-heel length and gestational age, whilst primiparity and caesarean section delivery were more common in children with breech presentation. There were no differences in height, body mass or body composition between breech and cephalic presentation children at age 4 years (data not shown).

Breech presentation was associated with lower neonatal total body BMC and BA, but not BMD or SA BMC (Figure 1). After adjustment for potential confounders in Model 2, children with breech presentation had 25% lower BMC, 22% lower BA and 3% lower BMD than those with cephalic presentation (equivalent to 0.9 SD, 0.9 SD and 0.5 SD deficits respectively). After adjustment for gestational age, birthweight, and length in Model 3 these associations were substantially attenuated, although lower BMC (7% or 0.26 SD) and BA (5% or 0.22 SD) were still evident in breech presentation.

There were no group differences evident in outcomes from total body (less head) or spine scans at age 4 years (Supplementary Figure 1). However, in unadjusted Model 1, breech presentation was associated with lower hip BMC and BA (but not BMD or SA BMC)

(Figure 2). There was little effect of adjustment in further models, such that in fully-adjusted analyses BMC was 7% (0.33 SD) and BA 5% (0.35 SD) lower in children who had a breech presentation at delivery than in those with a cephalic presentation.

Sensitivity/additional analyses

Similar results were observed when using total body measures without subtraction of head segment (values not shown). Head circumference at birth was similar in breech (35.0 ± 1.4 cm) and cephalic (35.0 ± 1.1 cm) births ($P = 0.77$), and additional adjustment for head circumference did not attenuate associations between breech presentation and bone outcomes. Restricting analyses only to those individuals with neonatal and 4-year scans resulted in similar regression coefficients (Supplementary Table 1). VIF analysis indicated moderate multicollinearity ($3 < \text{VIF} < 3.5$) for birthweight and birth length in neonatal outcome models, but not 4-year outcomes where $\text{VIF} < 2$ in all cases. However, these variables were strong independent predictors of neonatal outcomes, and removal of either variable in additional sensitivity analyses did not substantially affect regression coefficients for other variables including breech presentation.

Discussion

In this large mother-offspring cohort study, breech presentation at delivery was associated with lower bone mass, area and density at birth. These associations were substantially attenuated by adjustment for birthweight, crown-heel length and gestational age, although residual associations were observed. These associations did not persist at age 4 years, but here breech presentation was modestly associated with lower hip bone mass and area (but not density).

Comparison with previous findings

A previous study found lower tibia bone ultrasound speed of sound measurements (an indicator of bone quality) in breech versus cephalic-presenting babies at birth [26]. To the best of our knowledge, this is the first study to examine bone outcomes using DXA in breech and cephalic presentations. In addition, this study extends previous work by investigating whether deficits in bone outcomes persist into later childhood. Whilst deficits in total body outcomes were not evident at age 4 years, localised deficits in hip bone mass and area were observed even after consideration of potential confounding factors. This site corresponds with the location of dysplasias [24] and joint instability [28] which are more commonly found in breech births, and with altered femoral geometry [25].

Potential explanations of findings

Whilst large deficits in neonatal bone mass and area were observed in breech births, a large proportion of this difference was explained by shorter gestation and lower birthweight in these infants. Fetal growth has previously been shown to be impaired in children who present as breech at birth even accounting for differences in gestational age [29], although, in this cohort, sex and gestation-adjusted birthweight was not affected by presentation. However, when mediating factors such as birthweight and gestational age were considered, residual deficits in neonatal bone mass and area were still evident in breech presentation.

A potential cause of skeletal deficits associated with breech presentation independent of body size and length of pregnancy is the restriction of normal frequency and type of fetal movements, particularly later in pregnancy. However, the exact cause of reduced fetal movements is unknown and could relate to maternal or fetal factors. Whilst maternal mechanical factors such as oligohydramnios or uterine malformations likely contribute to reduced fetal movement, they are only evident in a small percentage of breech presentation births [29]. Breech presentation may be a result of poor neuromuscular development, leading to poor movement and an inability to attain a cephalic position. Evidence for this includes the higher breech incidence in children with conditions affecting motor development such as cerebral palsy [21], and correlation between breech incidence and severity of impairments either between or within clinical conditions (*e.g.* thoracic rather than lumbar level meningocele) [19]. In addition, umbilical cord length correlates with fetal movement [30] (and in turn bone ultrasound measures at birth [31]), and cords are shorter in breech than cephalic presentation [32]. Regardless of the cause, it seems likely that reduced fetal movements contribute to the size-independent deficits in bone outcomes in breech presentation evident in this study.

Significance and implications

Individuals born in the breech position are a substantial group when considered at the population level (approximately 22,500 p.a. in the UK). Therefore, any health deficits identified in this group have implications for a large number of individuals. Childhood is an important time for the attainment of a high peak bone mass, which is protective against fractures in later life [7]. Total body and regional deficits in bone mass identified at birth and in early childhood may therefore have implications for osteoporosis and fracture risk in later life. Studies examining bone outcomes in adult or elderly individuals whose mode of presentation at birth is known would be a valuable development of this study. Effective interventions to reverse breech presentation are available [33]. However, whilst they substantially reduce the incidence of hip dysplasias in breech presentation residual excess risk still exists [34]. This may be due to the lack of information on time sensitivity of exposure to breech position during pregnancy. From a 50% incidence of breech presentation at 25 weeks' gestation, there is a roughly linear decline to only ~5% incidence at term [17]. Therefore, studies combining regular observations of fetal position with neonatal bone mass and broader skeletal outcomes might reveal periods during which cephalic version must be maintained to ensure healthy skeletal development.

Strengths and limitations

The strengths of the study are the use of a large, representative cohort and consideration of a number of potential maternal and offspring confounders, in addition to exclusion of foetuses with known congenital malformations. However, as this was an observational study, causality cannot be attributed. Longitudinal measurements of offspring bone outcomes were only available in a limited sample, although associations between breech presentation and bone outcomes observed in this sub-cohort were similar to those found in larger groups of children with complete data at only one timepoint. In addition, fetal position was only recorded at birth with no information on position throughout pregnancy. Around 45% of the individuals with cephalic presentation at birth will have occupied a breech position around

25 weeks [35]. A decreasing percentage of these children will have occupied the breech position from 25-36 weeks, although the likelihood of version to a cephalic position decreases with length of gestation [36]. Conversely, very few babies move from cephalic to breech position in the later stages of pregnancy. Therefore, whilst we can be confident that individuals presenting as breech at birth will have occupied that position for the later stages of pregnancy, many individuals with cephalic presentation will have occupied a large portion of their prenatal life in a breech position. As a result we may have underestimated the effects of a continual breech position compared to a continual cephalic position throughout the third trimester, when mechanical loading of the skeleton is greatest [18].

The breech-presenting cohort was small and we may not have been powered to detect minor effects of breech position on total bone measures in early childhood. Whilst hip DXA scans have been used to examine bone strength in children of similar age in this and other cohorts [37–39], regional bone scans are not commonly used in clinical practice at this age, and are likely to yield a less reliable measure than those at the whole body or spine. Similar studies in older children would reveal whether hip bone mass deficits persist at a stage where they become clinically relevant.

Conclusions

Breech presentation is associated with lower neonatal total body bone mass and area, differences which, although attenuated, remained statistically significant after adjustment for gestation, birthweight and length. They did not persist at 4 years, although modest deficits in hip bone mass and area were observed at this age. The findings of this study extend previous work reporting altered neonatal hip geometry and higher risk of joint instability and dysplasias at birth in breech presentation, but confirmation of a localised effect in later childhood is required. Future studies examining fetal position throughout pregnancy, and neonatal skeletal health could give valuable information on key mechanosensitive periods for the developing skeleton.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

KMG is supported by the UK Medical Research Council (MC_UU_12011/4), the National Institute for Health Research (as an NIHR Senior Investigator (NF-SI-0515-10042) and through the NIHR Southampton Biomedical Research Centre) and the European Union's Erasmus+ Capacity-Building ENeASEA Project and Seventh Framework Programme (FP7/2007-2013), projects EarlyNutrition and ODIN under grant agreement numbers 289346 and 613977; and the ALPHABET project [an award made through the ERA-Net on Biomarkers for Nutrition and Health (ERA HDHL)], Horizon 2020 grant agreement number 696295 (UK component: BBSRC: BB/P028179/1). HMI is supported by the UK Medical Research Council (MC_UU_12011.4). This work was supported by grants from the Medical Research Council, British Heart Foundation, Arthritis Research UK, National Institute for Health Research (NIHR) Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Nestec and NIHR Biomedical Research Centre, University of Oxford.

References

1. Harvey NC, Javaid MK, Arden NK, et al. Maternal predictors of neonatal bone size and geometry: the Southampton Women's Survey. *J Dev Orig Health Dis.* 2010; 1:35–41. [PubMed: 23750315]
2. Godfrey K, Walker-Bone K, Robinson S, Taylor P, Shore S, Wheeler T, Cooper C. Neonatal bone mass: influence of parental birthweight, maternal smoking, body composition, and activity during pregnancy. *J Bone Miner Res.* 2001; 16:1694–1703. [PubMed: 11547840]
3. Holroyd CR, Osmond C, Barker DJ, Ring SM, Lawlor DA, Tobias JH, Smith GD, Cooper C, Harvey NC. Placental Size Is Associated Differentially With Postnatal Bone Size and Density. *J Bone Miner Res.* 2016; 31:1855–1864. [PubMed: 26999363]
4. Rogers IS, Ness AR, Steer CD, Wells JC, Emmett PM, Reilly JR, Tobias J, Smith GD. Associations of size at birth and dual-energy X-ray absorptiometry measures of lean and fat mass at 9 to 10 y of age. *Am J Clin Nutr.* 2006; 84:739–747. [PubMed: 17023699]
5. Dennison EM, Syddall HE, Sayer AA, Gilbody HJ, Cooper C. Birth weight and weight at 1 year are independent determinants of bone mass in the seventh decade: the Hertfordshire cohort study. *Pediatr Res.* 2005; 57:582–586. [PubMed: 15695596]
6. Hannam K, Lawlor DA, Tobias JH. Maternal Preeclampsia Is Associated With Reduced Adolescent Offspring Hip BMD in a UK Population-Based Birth Cohort. *J Bone Miner Res.* 2015; 30:1684–1691. [PubMed: 25761963]
7. Cooper C, Westlake S, Harvey N, Javaid K, Dennison E, Hanson M. Review: developmental origins of osteoporotic fracture. *Osteoporos Int.* 2006; 17:337–347. [PubMed: 16331359]
8. Ireland A, Rittweger J, Schönauf E, Lamberg-Allardt C, Viljakainen H. Time Since Onset of Walking Predicts Tibial Bone Strength in Early Childhood. *Bone.* 2014; 68:76–84. [PubMed: 25132490]
9. Ireland A, Maden-Wilkinson T, McPhee J, Cooke K, Narici M, Degens H, Rittweger J. Upper Limb Muscle-Bone Asymmetries and Bone Adaptation in Elite Youth Tennis Players. *Med Sci Sports Exerc.* 2013; 45:1749–1758. [PubMed: 23475166]
10. Sayers A, Deere K, Tobias JH. The effect of vigorous physical activity and body composition on cortical bone mass in adolescence. *J Bone Miner Res.* 2015; 30:584. [PubMed: 25380296]
11. Ireland A, Sayers A, Deere KC, Emond A, Tobias JH. Motor Competence in Early Childhood Is Positively Associated With Bone Strength in Late Adolescence. *J Bone Miner Res.* 2016; 31:1089–1098. [PubMed: 26713753]
12. Ireland A, Muthuri S, Rittweger J, Adams JE, Ward KA, Kuh D, Cooper R. Later Age at Onset of Independent Walking Is Associated with Lower Bone Strength at Fracture-Prone Sites in Older Men. *J Bone Miner Res.* 2017
13. Hannam K, Deere KC, Hartley A, Al-Sari UA, Clark EM, Fraser WD, Tobias JH. Habitual levels of higher, but not medium or low, impact physical activity are positively related to lower limb bone strength in older women: findings from a population-based study using accelerometers to classify impact magnitude. *Osteoporos Int.* 2016
14. Johansson J, Nordström A, Nordström P. Objectively measured physical activity is associated with parameters of bone in 70-year-old men and women. *Bone.* 2015; 81:72–79. [PubMed: 26151120]
15. Sival DA, Prechtl HF, Sonder GH, Touwen BC. The effect of intra-uterine breech position on postnatal motor functions of the lower limbs. *Early Hum Dev.* 1993; 32:161–176. [PubMed: 8486118]
16. Miller E, Kouam L. Zur Häufigkeit von Beckenendlagen im Verlauf Der Schwangerschaft und zum Zeitpunkt der Geburt. *Zentralbl Gynakol.* 1981; 103:105–109. [PubMed: 7234203]
17. Cammu H, Dony N, Martens G, Colman R. Common determinants of breech presentation at birth in singletons: a population-based study. *Eur J Obstet Gynecol Reprod Biol.* 2014; 177:106–109. [PubMed: 24784711]
18. Verbruggen SW, Kainz B, Shelmerdine SC, Hajnal JV, Rutherford MA, Arthurs OJ, Phillips AT, Nowlan NC. Stresses and strains on the human fetal skeleton during development. *Journal of The Royal Society Interface.* 2018
19. Bartlett D, Okun N. Breech presentation: a random event or an explainable phenomenon? *Dev Med Child Neurol.* 1994; 36:833–838. [PubMed: 7926333]

20. Fong BF, Savelsbergh GJ, de Vries JI. Fetal leg posture in uncomplicated breech and cephalic pregnancies. *Eur J Pediatr.* 2009; 168:443–447. [PubMed: 18661151]
21. Andersen GL, Irgens LM, Skranes J, Salvesen KA, Meberg A, Vik T. Is breech presentation a risk factor for cerebral palsy? A Norwegian birth cohort study. *Dev Med Child Neurol.* 2009; 51:860–865. [PubMed: 19469792]
22. Nowlan NC, Sharpe J, Roddy KA, Prendergast PJ, Murphy P. Mechanobiology of embryonic skeletal development: Insights from animal models. *Birth Defects Res C Embryo Today.* 2010; 90:203–213. [PubMed: 20860060]
23. Giorgi M, Carriero A, Shefelbine SJ, Nowlan NC. Effects of normal and abnormal loading conditions on morphogenesis of the prenatal hip joint: application to hip dysplasia. *J Biomech.* 2015; 48:3390–3397. [PubMed: 26163754]
24. Chan A, McCaul KA, Cundy PJ, Haan EA, Byron-Scott R. Perinatal risk factors for developmental dysplasia of the hip. *Arch Dis Child Fetal Neonatal Ed.* 1997; 76:F94–100. [PubMed: 9135287]
25. Hinderaker T, Uden A, Reikerås O. Direct ultrasonographic measurement of femoral anteversion in newborns. *Skeletal Radiol.* 1994; 23:133–135. [PubMed: 8191298]
26. Tshorny M, Mimouni FB, Littner Y, Alper A, Mandel D. Decreased neonatal tibial bone ultrasound velocity in term infants born after breech presentation. *J Perinatol.* 2007; 27:693–696. [PubMed: 17703182]
27. World Health Organisation. WHO child growth standards: length/height for age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age, methods and development. World Health Organization; Geneva, Switzerland: 2006.
28. Hinderaker T, Daltveit AK, Irgens LM, Udén A, Reikerås O. The impact of intra-uterine factors on neonatal hip instability. An analysis of 1,059,479 children in Norway. *Acta Orthop Scand.* 1994; 65:239–242. [PubMed: 8042471]
29. Luterkort M, Persson PH, Weldner BM. Maternal and fetal factors in breech presentation. *Obstet Gynecol.* 1984; 64:55–59. [PubMed: 6738947]
30. Moessinger AC, Blanc WA, Marone PA, Polsen DC. Umbilical cord length as an index of fetal activity: experimental study and clinical implications. *Pediatr Res.* 1982; 16:109–112. [PubMed: 7036074]
31. Wright D, Chan GM. Fetal bone strength and umbilical cord length. *J Perinatol.* 2009; 29:603–605. [PubMed: 19474797]
32. Soernes T, Bakke T. The length of the human umbilical cord in vertex and breech presentations. *Am J Obstet Gynecol.* 1986; 154:1086–1087. [PubMed: 3706433]
33. Hofmeyr GJ, Kulier R, West HM. External cephalic version for breech presentation at term. *Cochrane Database Syst Rev.* 2015 CD000083.
34. Lambeek AF, De Hundt M, Vlemmix F, Akerboom BM, Bais JM, Papatsonis DN, Mol BW, Kok M. Risk of developmental dysplasia of the hip in breech presentation: the effect of successful external cephalic version. *BJOG.* 2013; 120:607–612. [PubMed: 23145903]
35. Suzuki S, Yamamuro T. Fetal movement and fetal presentation. *Early Hum Dev.* 1985; 11:255–263. [PubMed: 4054045]
36. Witkop CT, Zhang J, Sun W, Troendle J. Natural history of fetal position during pregnancy and risk of nonvertex delivery. *Obstet Gynecol.* 2008; 111:875–880. [PubMed: 18378746]
37. Harvey NC, Cole ZA, Crozier SR, et al. Physical activity, calcium intake and childhood bone mineral: a population-based cross-sectional study. *Osteoporos Int.* 2012; 23:121–130. [PubMed: 21562877]
38. Janz KF, Burns TL, Torner JC, Levy SM, Paulos R, Willing MC, Warren JJ. Physical activity and bone measures in young children: the Iowa bone development study. *Pediatrics.* 2001; 107:1387–1393. [PubMed: 11389262]
39. Johannsen N, Binkley T, Englert V, Neiderauer G, Specker B. Bone response to jumping is site-specific in children: a randomized trial. *Bone.* 2003; 33:533–539. [PubMed: 14555256]

Mini Abstract

We compared bone outcomes in children with breech and cephalic presentation at delivery. Neonatal whole-body bone mineral content (BMC) and area were lower in children with breech presentation. At four years, no differences in whole-body or spine measures were found, but hip BMC and area were lower after breech presentation.

Figure 1.

Neonatal bone outcomes in breech presentation, shown as regression coefficients and 95% CI. Data adjusted as follows:

Model 1 - Presentation + Sex,

Model 2 - Model 1 + Maternal Smoking + Social Class + Birthweight + Parity + Ethnicity + Age at Scan,

Model 3 - Model 2 + Birthweight + Gestational Age + Crown-Heel Length

Figure 2.

Hip bone outcomes at four years of age in breech presentation, shown as regression coefficients and 95% CI. Data adjusted as follows:

Model 1 - Presentation + Sex

Model 2- Model 1 + Maternal Smoking + Social Class + Birthweight + Parity + Ethnicity + Age at Scan

Model 3 - Model 2 + Birthweight + Gestational Age + Height at Scan

Table 1

Cohort characteristics separated by presentation.

Variable		Presentation at birth			
		Cephalic		Breech	
n (male/female)		1372 (717/655)		58 (28/30)	
		Mean	SD	Mean	SD
Gestational age (weeks)		40	1.6	38.8	0.9
Crown-heel length	cm	50	2.1	48.5	1.8
	Z-score	-0.28	0.82	-0.59	0.89
Birthweight	g	3492	512	3239	436
	Z-score	0.06	0.95	-0.05	0.92
Age at time of scan (days)		6.4	4.9	4.9	4.3
Maternal age (years)		30.8	3.8	30.8	3.8
		n	%	n	%
Maternal Social Class	I	164	12	6	10.3
	II	622	45.2	30	51.7
	IIINM	374	27.3	11	19
	IIIM	122	8.9	5	8.6
	IV	87	6.3	6	10.3
	V	3	0.2	0	0
Ethnicity (white)		1322	96.4	58	100
Maternal Smoking		204	14.9	10	17.2
Parity (primiparous)		693	50.5	40	69
Gestational diabetes		12	0.9	0	0
Mode of Delivery	Spontaneous	887	64.7	2	3.4
	Forceps	123	9	0	0
	Caesarean section	264	19.2	56	96.6
	Ventouse	98	7.1	0	0

Table 2

Bone outcomes by presentation type

Variable		Presentation at birth			
		Cephalic		Breech	
		Mean	SD	Mean	SD
Neonatal DXA (n=965)					
Total Body	BMC (g)	63	26	52	12
	BA (cm ²)	118	26	98	21
	BMD g/(cm ²)	0.53	0.03	0.52	0.02
DXA at four years (n=999)					
Total Body	BMC (g)	371	44	367	51
	BA (cm ²)	754	48	751	56
	BMD (g/cm ²)	0.49	0.04	0.49	0.04
Hip	BMC (g)	7.2	1.6	6.7	1.6
	BA (cm ²)	12.6	1.9	11.9	2.2
	BMD (g/cm ²)	0.57	0.06	0.56	0.06
Spine	BMC (g)	13.1	2	12.9	1.8
	BA (cm ²)	27.2	2.7	26.7	2.8
	BMD (g/cm ²)	0.48	0.05	0.48	0.05