Evidence of Subclavian Vein Thrombosis on First-Pass ^{18}F FDG PET in a Patient with Relapsing Upper Mediastinum Lymphoma

Fayçal Ben Bouallègue^{1,2,3} · Fabien Vauchot¹ · Denis Mariano-Goulart^{1,2}

Received: 4 July 2018 / Revised: 7 August 2018 / Accepted: 8 August 2018 / Published online: 20 August 2018
© Korean Society of Nuclear Medicine 2018

Abstract

Baseline ^{18}F -FDG PET was performed in a 74-year-old patient with relapsing upper mediastinum lymphoma. Left subclavian thrombosis was suspected on prior contrast-enhanced CT. Dynamic PET imaging was achieved during 3 min after IV injection of ^{18}F -FDG to the left arm in order to further assess left subclavian vein permeability. The 20-s dynamic frame at 1 min after injection confirmed the absence of flow in the left subclavian vein and evidenced the derivation of ^{18}F -FDG through left axillary, then superficial, then right internal mammary collaterals to the superior vena cava, hence confirming the subclavian thrombosis.

Keywords FDG PET · First-pass · Deep vein thrombosis · Lymphoma

✉ Fayçal Ben Bouallègue
faycal.ben-bouallegue@umontpellier.fr

¹ Nuclear Medicine Department, Montpellier University Hospital, Montpellier, France

² PhyMedExp, INSERM, CNRS, Montpellier University, Montpellier, France

³ Nuclear medicine department, Lapeyronie University Hospital, Avenue du Doyen Giraud, 34295 Montpellier Cedex 5, France

◀ **Fig. 1** Baseline ^{18}F -FDG PET was performed in a 74-year-old woman with suspected relapse of diffuse large B cell lymphoma of the upper mediastinum. Prior contrast-enhanced CT (**a**) showed upper left mediastinum adenomegalies compressing the left brachiocephalic vein with anterior thoracic collaterality (red arrowheads) and possible left subclavian thrombosis (green arrowheads). First-pass ^{18}F -FDG PET-CT was performed during 3 min after IV injection to the left arm of 178 MBq in order to assess left subclavian vein permeability. The 20-s dynamic frame at 1 min after injection (**b**) confirmed the absence of flow in the left subclavian vein (yellow arrows) and evidenced the derivation of ^{18}F -FDG through left axillary, then superficial, then right internal mammary collaterals to the superior vena cava (green arrows). Static acquisition at 60 min post-injection (**c**) showed hypermetabolic foci centred on upper left mediastinum lymph nodes (red arrows) and no significant tracer activity in the left subclavian vein (yellow arrow). Left anterior oblique maximum intensity projection images (**d**) at first-pass (left, green arrows show venous collateral deviation of the tracer) and equilibrium (right, red arrow points the hypermetabolic lymph nodes) are provided as a synthetic view. Venous thromboembolism is frequent in hematological malignancies, particularly in aggressive lymphomas with an incidence of about 10% [1], and is associated with substantial morbidity and mortality. Standard diagnosis relies on ultrasonography, or contrast-enhanced CT in case of abdominopelvic involvement [2, 3]. Alternative diagnostic explorations relying on metabolic imaging have been proposed, including radionuclide venography using macroaggregated albumin, labeled platelets scintigraphy, and fibrin or plasmin targeting peptide imaging [4, 5]. Recent preliminary studies have reported on the potential value of ^{18}F -FDG PET in the diagnostic evaluation of acute deep vein thrombosis [6, 7]. However, other research has highlighted the critical lack of sensitivity and specificity of both visual assessment and SUV measurements in routine clinical setting [8, 9]. Dynamic ^{18}F -FDG PET imaging during tracer first-pass allows direct visualization of functional vasculature and assessment of venous permeability. It may offer additional diagnostic information in hematological patients with clinically or radiologically suspected deep vein thrombosis, without any modification in injected dose or protocol duration

Compliance with Ethical Standards

Conflict of Interest Fayçal Ben Bouallègue, Fabien Vauchot, and Denis Mariano-Goulart declare that they have no conflict of interest.

Ethical Approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in the study.

References

1. Falanga A, Marchetti M. Venous thromboembolism in the hematologic malignancies. *J Clin Oncol*. 2009;27:4848–57.
2. Karande GY, Hedgire SS, Sanchez Y, Baliyan V, Mishra V, Ganguli S, et al. Advanced imaging in acute and chronic deep vein thrombosis. *Cardiovasc Diagn Ther*. 2016;6:493–507.
3. Khosa F, Otero HJ, Prevedello LM, Rybicki FJ, Di Salvo DN. Imaging presentation of venous thrombosis in patients with cancer. *AJR Am J Roentgenol*. 2010;194:1099–108.
4. Seabold JE. Radionuclide venography and labeled platelets in deep venous thrombosis. *Semin Nucl Med*. 2001;31:124–8.
5. Houshmand S, Salavati A, Hess S, Ravina M, Alavi A. The role of molecular imaging in diagnosis of deep vein thrombosis. *Am J Nucl Med Mol Imaging*. 2014;4:406–25.
6. Rondina MT, Lam UT, Pendleton RC, Kraiss LW, Wanner N, Zimmerman GA, et al. (18)F-FDG PET in the evaluation of acuity of deep vein thrombosis. *Clin Nucl Med*. 2012;37:1139–45.
7. Hess S, Madsen PH, Iversen ED, Frifelt JJ, Hoiland-Carlsen PF, Alavi A. Efficacy of FDG PET/CT imaging for venous thromboembolic disorders: preliminary results from a prospective, observational pilot study. *Clin Nucl Med*. 2015;40:e23–6.
8. Le Roux PY, Robin P, Delluc A, Tardy B, Abgral R, Couturaud F, et al. Performance of 18F fluoro-2-désoxy-D-glucose positron emission tomography/computed tomography for the diagnosis of venous thromboembolism. *Thromb Res*. 2015;135:31–5.
9. Zhu HJ, Hess S, Rubello D, Goris ML, Alavi A. The strong but nonspecific relationship between 18F-FDG uptake in the lower-extremity veins and venous thromboembolism. *Nucl Med Commun*. 2016;37:322–8.