REVIEW

For reprint orders, please contact: reprints@futuremedicine.com

Surgical management of high-grade glioma: a standard of care

Colin Watts*

Practice Points

- The most important priority for the neurosurgical oncologist is patient safety.
- The role of the neurosurgeon begins long before the patient is taken to the operating room and requires them to function as part of a multidisciplinary team.
- Integration of multimodal datasets is an integral part of preoperative surgical planning in the modern era. The use of MRI rather than computed tomography imaging should be routine at every stage of the patient journey.
- Advanced technologies are no substitute for detailed knowledge of neuro-anatomy and neuro-physiology.
- Meticulous neurosurgical technique is essential when operating on patients with high-grade glioma, recognizing that the extent of resection is a vital component of overall patient management.
- Age alone should not be a contraindication to radical surgery. The performance status and intercurrent morbidity inform surgical decision-making together with the location and extent of disease.
- A multidisciplinary approach is essential in managing patients with recurrent high-grade glioma.

SUMMARY Neurosurgical oncology for intrinsic glioma is evolving rapidly. It must be patient-centered, consultant-led and research-orientated. The value of specialist neurosurgical engagement is becoming more widely recognized. Detailed evaluation tailored to each patient is essential before the surgical admission, in conjunction with clinical oncology input. Medical optimization, collation of magnetic resonance datasets for preoperative planning and providing an informed explanation of the proposed management and its alternatives are all part of the neurosurgeon's remit. Meticulous microsurgical technique during surgery utilizing modern neuronavigation and physiological monitoring are integral components of the specialist armamentarium. A clear understanding of the rationale for surgical intervention, including its place alongside radiotherapy and chemotherapy, informs surgical decision-making. Recognition and understanding of these issues are driving the evolution of neurosurgical management of high-grade glioma. New challenges are emerging and need to be critically evaluated in robustly designed clinical trials.

*University of Cambridge Department of Clinical Neurosciences, Division of Neurosurgery, Box 167 Addenbrookes Hospital, Hills Road, Cambridge, CB2 0QQ, UK; cw209@cam.ac.uk

The role of the surgeon in the management of patients with high-grade glioma (HGG) has never been more important. The key to ensuring the success of the role is to recognize that the surgeon must be part of a multidisciplinary team (MDT) whose philosophy eschews therapeutic nihilism [1]. Clinical service delivery must be patient-centered, consultant-led and research-orientated. The research must be ethically rigorous and peer reviewed to ensure that there is no potential antagonism between benefit for the patient now versus future benefit for the discipline. Since any one unit will only ever see a limited number of patients with HGG, it is essential that the process of care is as evidence-based as possible and replicated across multiple units, forming a national network of neuro-oncology centers. This is the only way sufficient numbers of patients can be recruited into randomized clinical trials that will generate the evidence base needed to move the discipline forward. This review will examine this philosophy from the neurosurgical perspective.

Role of the surgeon in the pathway of care

Surgical management begins long before the patient gets to the operating room. The neurosurgeon is often the first point of contact for specialist care of patients with HGG. It is essential that a minimum imaging dataset is obtained that will provide sufficient qualitative structural information about the tumor and surrounding anatomy to enable a radiological diagnosis of HGG to be established and surgical planning to be developed. These should be volumetric magnetic resonance (MR) datasets including T₁ pre- and post-contrast, T₂ and fluid attenuation inversion recovery. Diffusion-weighted imaging is also useful in excluding a cerebral abscess that has restricted diffusion [2].

Armed with this information together with an accurate clinical history, a preoperative MDT review involving neuro-oncologists, neuro-radiologists and neurosurgeons is the next step. This allows a comprehensive management plan to be determined and the patient to be stabilized on steroids and anticonvulsants as required [3]. It also provides opportunity to discuss potential clinical trials for which each patient might be suitable and to consider tissue collection and banking as an integrated component of the process of consent [3]. In many centers, the MDT does not occur until a pathologic diagnosis is available. This limits specialist input prior to

surgery, which can compromise medical optimization. An alternative often proposed is a 'pre-operative protocol', determined by the MDT. However, this does not allow management to be tailored to suit the individual patient. It also limits its opportunity to discuss tissue collection and research trials, a major problem in many centers.

Many patients are referred as 'emergencies' and often languish in their referring hospital until a neurosurgical bed is available, whereupon they are transferred, often for surgery the following day. We have established a care pathway that allows the patient to be sent home wherever possible and to attend a dedicated neurosurgical oncology clinic as an outpatient (Figure 1). At the outpatient review the surgical plan can be finalized, the imaging discussed and a date set for surgery. The process of consent is initiated and potential research studies discussed. Patient information and specialist nurse contact details can be given. Thus, the patient leaves clinic with a clear plan of action. This 'urgent' rather than 'emergent' outpatient-based approach to the surgical management of HGG is both clinically efficient and cost effective [3].

The scale of the problem

WHO classification of tumors of the CNS has established a grading system for astrocytic tumors. HGG comprise tumors with cytological atypia, anaplasia and mitoses (grade III), plus necrosis (grade IV). This malignancy scale includes diverse morphological entities, which are reviewed in detail by Louis *et al.* [4]. For pragmatic surgical purposes these include glioblastoma multiforme (GBM), anaplastic astrocytoma (AA), anaplastic oligodendroglioma (AOD) and anaplastic oligoastrocytoma (AOA). With an incidence of 3.55/100,000 population per year, GBM is the commonest histological subtype, accounting for 69% of astrocytic and oligodendroglial tumors, followed by AA (0.25/100,000), AOD (0.07/100,000) and AOA (0.06/100,000) [5]. In optimally managed patients with GBM who have a good performance status, a good resection and radical radiotherapy, plus concomitant and adjuvant temozolomide chemotherapy, median survival is approximately 14 months [6]. However, much of the survival benefit derived from temozolomide is limited to those patients with promoter methylation of *MGMT* [7]. In patients with secondary GBM, improved prognosis is also strongly associated with mutation of *IDH1* [8].

Figure 1. The pathway of care. The consultant neurosurgeon checks scans of referred patients to ensure surgical emergencies such as cerebral abscess are not missed. The referring team pass patient information to the MDT. Following review, the patient is seen in the surgical glioma clinic. There is a postoperative MDT review of diagnosis, followed by a same-day consultation led by the consultant neurosurgeon to give diagnosis. This is followed by clinical oncology review to consider further management.

MDT: Multidisciplinary team.

Adapted with permission from [3].

Survival for optimally managed patients with AA is approximately 3 years [9], compared with 10 years for AOD. In AOD patients with 1p/19q codeletion treated with radiotherapy plus procarbazine, lomustine and vincristine, median survival exceeds 12 years [VAN DEN BENT MJ, UNPUBLISHED DATA].

However, many patients do not have favorable prognostic factors and/or are not suitable for radical therapy. Population-based survival statistics are rare. In one study [5], the median survival for patients with AOD was 3.5 years, for AA was 1.6 years and for GBM just 4.9 months, compared with 5.6 years for astrocytoma and long-term survival for surgically resected pilocytic astrocytomas (WHO grade I astrocytomas) (Figure 2). These poor survival statistics are reflected in the average years of life lost by individual patients: 20.1 years for brain cancer,

compared with a mean of 12.6 years for all cancers [10]. The high levels of premature morbidity and mortality, together with a lack of therapeutic options, make these cancers an area of unmet clinical need.

The rationale for surgery

■ Cytoreduction & palliation of symptoms

Although our understanding of the biological diversity of glioma has increased dramatically [11], diagnosis is still made primarily on morphological criteria by the neuropathologist, and they share the common clinical phenotype of diffusely infiltrating tumors [12]. Consequently, aggressive cytoreduction rather than radical excision is the primary objective of surgery. This strategy encompasses more than a rapid log-kill of tumor cells [13]. Palliation of symptoms related to raised intracerebral pressure and

Figure 2. Population-based survival data for patients with astrocytomas. Only those with pilocytic astrocytomas have prolonged survival. Patients with high-grade astrocytomas (anaplastic astrocytoma and glioblastoma) have very poor survival.

MST: Median survival time.

Reproduced with permission from [89].

peritumoral edema has been established in large surgical series with associated delay in deterioration of neurological function [14–16]. However, in a study of 30 elderly (>65 years old) patients with radiological HGG randomized to either biopsy or craniotomy, improved overall survival was demonstrated in the craniotomy group, but no difference in time to neurological deterioration [17]. This study was too small to draw any definitive conclusions, but emphasizes the need for careful patient selection when considering aggressive surgery in HGG.

■ Improving progression-free & overall survival

There is growing recognition that in certain carefully selected patients surgery can play an important role in improving progression-free and overall survival. Numerous retrospective studies have evaluated the impact of surgery on overall survival, but interpretation

has been confounded by the fact that younger patients with high performance status tend to get more aggressive resections [16,18]. More recently, *post hoc* analysis of three large prospective randomized controlled trials has provided additional evidence for the benefit of surgery.

Patients with GBM recruited to the European Organisation for Research Trials in Cancer (EORTC) 26981-22981 National Cancer Institute of Canada clinical trials groups CE3 randomized Phase III trial have recently been analyzed for long-term outcome [19]. Putative prognostic and predictive factors were also analyzed. Overall 5-year survival was improved in patients receiving combined chemoradiotherapy. Those patients who received a complete resection survived longer than those with a partial resection [19]. However, early postoperative MRI was not mandated in this study, and estimates of completeness of resection were subjective and likely to be inaccurate [18].

In another prospective randomized controlled trial of 240 patients evaluating the survival benefit of intra-operative biodegradable carmustine (BCNU) wafer implantation [20], postoperative radiological data was used to determine extent of resection. Complete resection was defined as removal of >90% of tumor tissue on postoperative images obtained within 48–72 h of surgery. In both the BCNU group and in the placebo wafer group, a survival advantage was observed in patients receiving a complete resection compared with an incomplete resection [20,21]. However, there was no central radiology reporting, and volume and/or area analyses of MR data and the use of computed tomography data was allowed.

A cohort of 243 patients, from 322 randomized to receive 5-aminolevulinic acid (ALA) as an adjunct to surgical resection [16], was analyzed to compare patients with complete and incomplete resections. Those with complete resections survived longer (16.7 vs 11.8 months; $p < 0.0001$). Multivariate analysis of putative prognostic factors showed that only residual tumor, age and performance status were significant [22].

In a subsequent prospective noninterventional cohort study of 166 GBM patients, the completeness of resection was determined using MR data obtained within 72 h of surgery [23]. The absence of visible contrast-enhancing disease was associated with better progression-free survival and overall survival (Figure 3). In patients with no residual enhancing disease the

median overall survival exceeded the follow-up period of 24 months (mean: 23.6 months; range: 21.4–25.8 months) [23]. Together, these data provide strong evidence (level 2b) of the benefit of surgery in HGG, but it must be remembered that the patients in these studies represent a small fraction of patients with diffusely infiltrating HGG [5].

■ Control of seizure activity

Seizures are a common presenting symptom and cause of morbidity in patients with HGG. In a retrospective study of 648 patients with AA or GBM, multivariate logistic regression analysis identified AA pathology, temporal lobe involvement and cortical location as factors associated with preoperative seizures [24]. In those patients with preoperative seizures, the postoperative outcomes were seizure freedom (Engel class I) in 77%; rare seizures (Engel class II) in 12% and meaningful improvement (Engel class III) in 6%. In only 5% was there no improvement. Seizures are less common in primary HGG compared with low-grade glioma and secondary GBM, but are more difficult to control [25], and patients should be counseled accordingly.

■ Facilitating local therapies

The concept of delivering genotoxic technologies to the residual tumor at the end of cytoreductive surgery is an attractive concept. Several strategies have been attempted, including chemotherapy, immunotherapy [26], gene therapy and

Figure 3. Correlation between the extent of residual disease as determined on an early (within 72 h) postoperative MRI scan and survival. (A) Progression-free and (B) overall survival.

Reproduced with permission from [23].

brachytherapy [27,28]. However, only BCNU wafers and gene therapy have progressed to Phase III studies. Early Phase III data of HSV-tk gene therapy in GBM showed no benefit, but the poor results were attributed to poor gene-delivery technology and crude surgical techniques for vector delivery [29]. After initially promising Phase I/II data of HSV-tk gene therapy using an adenoviral vector [30], a subsequent Phase II/III study showed no benefit [31,32]. This poor translation of promising preclinical data is likely due to a combination of factors, including poor models of the disease, poor surgical delivery technologies and variable activity of the constructs [31]. In contrast, the use of intra-operative BCNU wafers has now progressed into mainstream practice. A prospective randomized placebo-controlled clinical trial of 240 patients with HGG [20] showed a modest improvement in survival in patients receiving BCNU wafers (13.9 months), compared with the placebo group (11.7 months; $p = 0.03$). Of the 240 randomized patients, 207 had GBM. The remainder were AA ($n = 2$), AOD ($n = 9$), AOA ($n = 10$), metastasis ($n = 3$) and 'other' ($n = 9$). When the GBM cohort was analyzed (101 in the treatment arm and 106 in the placebo arm), the survival advantage of the BCNU was lost with a trend to improved survival (13.5 months vs 11.4 months) that did not reach significance. However, when a Cox proportional hazards method was applied to correct for potential imbalances in prognostic factors, a modest survival advantage for BCNU wafers was demonstrated in the GBM subgroup (stratified log-rank statistic $p = 0.04$ with a 31% risk reduction; 95% CI: 3–51%). For these reasons, the uptake of this technology is far from universal, and the data suggest further prospective studies are needed.

■ Facilitating adjuvant treatment

When the clinical impact of resection was evaluated in a *post hoc* analysis of survival by extent of resection in the EORTC–National Cancer Institute of Canada trial cohort, the data suggested that the more complete the resection the longer the survival [33]. The median gain in survival for patients receiving radiotherapy plus temozolomide compared with radiotherapy alone was +4.1 months in patients with a complete resection, compared with +1.8 months in patients with incomplete resections and +1.5 months in those who were biopsied. Although this did not reach significance, Kaplan–Meier survival data

corresponded to a risk reduction for chemoradiotherapy versus radiotherapy alone of 43% in patients completely resected ($p = 0.0001$). Patients who were partially resected showed a risk reduction of 35% ($p = 0.0001$), whereas patients biopsied had a nonsignificant risk reduction of 31% ($p = 0.088$). Adjuvant temozolomide appeared to have the best survival impact in patients who were completely resected [34].

The concept that optimizing cytoreduction can enhance the efficacy of adjuvant therapies is novel [34]. Possible mechanisms include enhanced drug penetration, alterations in interstitial fluid pressure and modulation of hypoxic factors [34]. High-quality data on drug penetration into the residual tumor bed is limited. One study established the tissue to blood ratio of temozolomide using microdialysis catheters implanted into the tumor bed at the end of surgery [35]. The results confirmed limited tumor penetration with a ratio of 17.8% and a peak concentration of 0.6 $\mu\text{g/ml}$. Data using methyl- ^{11}C temozolomide PET predicted intra-tumor drug concentrations of 2.6–6.7 $\mu\text{g/ml}$ [36]. A possible explanation for the discrepancy is that the microdialysis technique only samples interstitial fluid. Defects in the blood–brain barrier in glioma result in raised interstitial fluid pressure (IFP), which may compromise drug penetration or preferentially distribute it to low-pressure areas such as necrotic regions [37]. Following surgery, cerebral edema in the early postoperative period increases IFP in the tumor bed. As the edema resolves the IFP declines, generating a positive pressure gradient allowing fluid to penetrate the tumor bed [38]. This may enhance drug penetration into the tumor margin [39,40].

Expression of various hypoxia inducible factors such as HIF2 α is associated with poor prognosis in glioblastoma, possibly by regulating tumor competency of gliomagenic cell populations [41,42]. However, clinical validation of antiangiogenic strategies in large-scale clinical trials has yet to be established [43,44]. Tumor margins are areas in which angiogenesis plays a significant role in the oncogenic process [45], and contain cell populations resistant to high-dose conformal radiation, thereby facilitating local recurrence [46]. Objective and reproducible surgical targeting of tumor margins is now possible [47], paving the way for improving our understanding of the biology of this tumor compartment, and hence how we can characterize and target the residual disease there.

Maximizing patient safety & the impact of postoperative neurological deficits

Internationally agreed management guidelines for both HGG [1] and low-grade glioma [48] both advocate maximal surgical resection when it is safe to do so. The diffusely infiltrative nature of glioma makes this extremely challenging to achieve, and close proximity to eloquent brain further compromises the resection of detectable disease [16,18,49]. Variability in the location of cortical function, particularly language, memory and other higher cognitive functions [50–53], adds a further level of complexity. It is important for the surgeon to recognize that the postoperative neurological deficits compromise patient survival. Surgically acquired motor or language deficits are independently associated with reduced median survival compared with patients with no new postoperative deficits [15,54]. The reason for this is unclear, but neurological deficits compromise subsequent radio- and chemo-therapy and this will impact on patient survival.

For these reasons, traditional neurosurgical techniques based on conventional microscopy, computed tomography imaging datasets and anatomical knowledge alone are insufficient for the neurosurgical management of any intrinsic neoplasm. Intra-operative neuronavigation has become a mainstream component of the neurosurgical armamentarium. Its use improves surgical confidence, accurate bone flap placement, reduced craniotomy size and anatomical orientation [55,56]. However, detailed accurate intra-operative navigation is compromised by brainshift [57]. Recent studies suggest that this problem can be abrogated by the integration of additional MR datasets such as diffusion tensor imaging-based fiber tracking [58–60]. Blood oxygen level-dependent functional MRI can be used to identify the location of eloquent brain to spare during surgery. Motor tapping yields robust identification of the primary motor cortex, and visual checkerboard stimuli identify primary visual cortex. Language-processing areas in frontal and temporal regions are best identified with a panel of paradigms that target receptive and expressive aspects of language, and also provide redundancy to assure reproducibility of findings [61]. These datasets can be incorporated into preoperative planning strategies [62,63]; however, concordance with physiological data obtained from direct cortical stimulation remains controversial and awaits robust prospective evaluation [63].

A third component of the modern neurosurgical toolbox is the use of awake craniotomies with intra-operative monitoring and mapping. This represents the gold standard for preserving function in glioma surgery [64]. The identification of cortically active areas can be used to refine the surgical approach [65], while stimulation of deep white matter tracts helps to define the deep margins of a resection and can establish a physiological end point for resection. This is particularly important in diffusely infiltrating disease. The importance of good neuro-anesthesia is essential to optimize functional outcome and minimize peri-operative complications [66].

Technologies allowing navigational datasets to be updated during surgery include intra-operative ultrasound and intra-operative MRI. Intra-operative ultrasound can be used to update the navigational datasets in real time [56,67]. It is relatively cheap and can be used repeatedly during the operation. It is most useful in surgery for low-grade glioma, but edema and hemorrhage at the operative site can limit its use in HGG surgery [68]. Intra-operative MRI (iMRI) can provide high-quality updated imaging datasets [69–71], leading to improved resection [59]. A recent prospective randomized trial of iMRI has been reported [72]. This recruited 58 patients who were randomized to surgery plus iMRI or standard neuronavigation-based resection. Postoperative volumetric datasets were used to establish a complete resection (defined as a residual volume $<0.175 \text{ cm}^3$). The addition of iMRI resulted in a complete resection in 23/24 patients (96%), compared with 17/25 (68%) in the control group. Enhanced resection was not associated with additional neurological deficits, indicating that iMRI has a role to play in optimizing both surgical resection and patient safety. It will be interesting to learn if this benefit is confined to the more expensive high-field machines or could be achieved with less expensive low-field technologies.

In many European centers, an alternative approach for patients with HGG is the use of intra-operative fluorescence (**Figure 4**), which provides real-time feedback for the surgeon, establishes an objective surgical end point and is cost effective. It is the only surgical adjunct to be validated in a randomized clinical trial [16].

It must be stressed that these technologies must be applied appropriately and to their full potential to have clinical utility. The surgeon must recognize the strengths and limitations of

Figure 4. Fluorescence-guided tumor resection. Visible fluorescence provides real-time feedback helping the surgeon to identify disease with >95% accuracy. Tumor extending through the cortical mantle can be visualized at the surface. Fluorescence reveals a greater extent of disease than can be visualized under standard operating conditions.

each and understand that these adjuncts complement, but cannot replace, detailed knowledge of neuroanatomy and meticulous microsurgical technique. The clinical data overwhelmingly support the concept that the basis for successful neurosurgical oncology is to prioritize patient safety.

A multidisciplinary approach is essential in managing patients with recurrent HGG

The role of surgery in the management of patients with recurrent HGG remains controversial. There is insufficient data to support specific practice guidelines [73], a situation that has not changed in over 10 years [74]. Validated criteria for patient selection have not been established, but focal recurrence, good performance status and a multidisciplinary rationale (such as potential for further chemotherapy) constitute good clinical practice. Unfortunately, the number of appropriate patients remains limited [75]. For patients progressing after previous chemotherapy there is no established chemotherapy regimen and patients should be entered into clinical trials wherever possible. Patients with recurrent anaplastic glioma may benefit from salvage chemotherapy [76–78]. Evidence of clinical benefit from intra-operative chemotherapy

wafers suggests modest benefit. In a trial of 222 patients with recurrent malignant glioma randomized to chemotherapy wafers, or placebo wafers median survival was 31 weeks compared with 23 weeks (hazard ratio: 0.67; $p = 0.006$, after controlling for prognostic factors) [79]. However, the use of chemotherapy wafers has failed to gain wide acceptance. The majority of patients are recruited into chemotherapy trials that have yet to demonstrate significant clinical efficacy [1] although modest survival benefit can be obtained [80].

Fluorescence has a high predictive value for detecting tumor in recurrent glioma. In a multicenter prospective single-arm study, the positive predictive value of visible 5-ALA fluorescence was over 97% in areas that appeared neoplastic. In areas that did not appear to be involved in recurrent disease, the positive predictive value was 93% [81]. These data support the use of 5-ALA in routine clinical practice in recurrent HGG.

Patients with recurrent HGG must be managed in a multidisciplinary environment incorporating the highest standards of neurosurgical practice. Patients with a good performance status and favorable prognostic biomarkers should be considered for further surgical intervention, preferably in the context of a clinical trial.

Managing the elderly patient

Population-based data confirm that the incidence of GBM is greatest in the elderly. The mean age is 61.3 ± 14.0 years, with nearly 60% of cases occurring in people over 60 years old [5]. Older people are traditionally subject to therapy bias linked to the perception that older patients might suffer more side effects and fewer benefits from standard treatment. Thus, older patients are usually under-represented in clinical trials [82,83], and brain cancer is no exception [80]. For example, the median age in the EORTC 26981 trial was 56 years [6], and in the BCNU trial 53 years [20]. The elderly also receive less aggressive treatment, beginning with a diagnostic biopsy rather than respective surgery [84], and extending to radiotherapy and chemotherapy [85–87].

The median survival of 83 patients aged over 60 years recruited into EORTC 26981 was 10.9 months [6]. In this study, the surgical management was not specified and data regarding the extent of resection were not available. More recent data incorporating modern

surgical techniques, including the use of fluorescence-guided cytorreduction, is available for 130 GBM patients with a median age of 68 years. The median survival for those who went on to receive radiotherapy and chemotherapy was 16.3 months (range: 12–17.2 months), compared with 11.2 months for those receiving radiotherapy alone [87]. These prospective data are supported by a retrospective analysis [88] and suggest that elderly patients with HGG are able to benefit from aggressive management and should not be excluded from current therapies on the basis of age alone.

Conclusion & future perspective

The modern management of patients with HGG must be patient-centered, led by specialist consultants and research-orientated. Patients should be reviewed by a multidisciplinary team before surgery. The use of neuronavigation, volumetric multimodality imaging, awake surgery, cortical mapping and 5-ALA fluorescence should all be part of routine specialist practice. Modern technology is not a substitute for detailed knowledge of neuroanatomy and neurophysiology, and meticulous microsurgical technique. Neurosurgical specialization will improve patient management and recruitment into clinical trials.

In the medium term, the process of care for patients with HGG should be standardized so that comparable national networks of

neuro-oncology centers can be developed to facilitate clinical care and research. A more challenging problem will be the early detection of HGG because the disease is well advanced at the time of clinical presentation. How early detection could be achieved with current technologies is unclear, because any neurological deficit is a poor predictor of CNS malignancy. This will make any imaging-based screening strategies expensive, time-consuming and relatively insensitive.

In the longer term, greater neurosurgical involvement and meticulous tissue collection is likely to drive improvement in our understanding of the biology of HGG from which new therapeutic strategies will emerge. From a surgical perspective, technologies to improve local delivery of targeted therapeutic agents will become a major component of the role of the neurosurgical oncologist.

Financial & competing interests disclosure

The author has no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as:

- of interest
 - of considerable interest
- 1 Stupp R, Tonn JC, Brada M, Pentheroudakis G. High-grade malignant glioma: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* 21(Suppl. 5), v190–v193 (2010).
 - 2 Waldman AD, Jackson A, Price SJ *et al.* Quantitative imaging biomarkers in neuro-oncology. *Nat. Rev. Clin. Oncol.* 6(8), 445–454 (2009).
 - 3 Guilfoyle MR, Weerakkody RA, Oswal A *et al.* Implementation of neuro-oncology service reconfiguration in accordance with NICE guidance provides enhanced clinical care for patients with glioblastoma multiforme. *Br. J. Cancer* 104(12), 1810–1815 (2011).
 - Details a clinically validated process of care, emphasizing a multidisciplinary approach and the role of the neurosurgeon.
 - 4 Louis DN, Ohgaki H, Wiestler OD *et al.* The 2007 WHO classification of tumors of the central nervous system. *Acta Neuropathol. (Berl.)* 114(2), 97–109 (2007).
 - 5 Ohgaki H, Kleihues P. Population-based studies on incidence, survival rates, and genetic alterations in astrocytic and oligodendroglial gliomas. *J. Neuropathol. Exp. Neurol.* 64(6), 479–489 (2005).
 - Highlights the grim reality for many patients who are not suitable for aggressive treatment.
 - 6 Stupp R, Mason WP, van den Bent MJ *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N. Engl. J. Med.* 352(10), 987–996 (2005).
 - 7 Hegi ME, Diserens AC, Gorlia T *et al.* MGMT gene silencing and benefit from temozolomide in glioblastoma. *N. Engl. J. Med.* 352(10), 997–1003 (2005).
 - 8 Hartmann C, Hentschel B, Wick W *et al.* Patients with *IDH1* wild type anaplastic astrocytomas exhibit worse prognosis than *IDH1*-mutated glioblastomas, and *IDH1* mutation status accounts for the unfavorable prognostic effect of higher age: implications for classification of gliomas. *Acta Neuropathol.* 120(6), 707–718 (2010).
 - Emphasizes the value of biomarkers when interpreting clinical outcome data.
 - 9 Wick W, Hartmann C, Engel C *et al.* NOA-04 randomized Phase III trial of sequential radiochemotherapy of anaplastic glioma with procarbazine, lomustine, and vincristine or temozolomide. *J. Clin. Oncol.* 27(35), 5874–5880 (2009).
 - 10 Burnet NG, Jefferies SJ, Benson RJ, Hunt DP, Treasure FP. Years of life lost (YLL) from cancer is an important measure of population burden – and should be considered when allocating research funds. *Br. J. Cancer* 92(2), 241–245 (2005).
 - 11 Huse JT, Phillips HS, Brennan CW. Molecular subclassification of diffuse gliomas: seeing order in the chaos. *Glia* 59(8), 1190–1199 (2011).

- 12 Louis DN. *WHO Classification of Tumors of the Central Nervous System (4th Edition)*. International Agency for Research on Cancer, Lyon, France (2007).
- 13 McCarter MD, Fong Y. Role for surgical cytoreduction in multimodality treatments for cancer. *Ann. Surg. Oncol.* 8(1), 38–43 (2001).
- 14 Sawaya R, Hammoud M, Schoppa D *et al.* Neurosurgical outcomes in a modern series of 400 craniotomies for treatment of parenchymal tumors. *Neurosurgery* 42(5), 1044–1055; discussion 1055–1056 (1998).
- 15 Stummer W, Tonn JC, Mehdorn HM *et al.* Counterbalancing risks and gains from extended resections in malignant glioma surgery: a supplemental analysis from the randomized 5-aminolevulinic acid glioma resection study. Clinical article. *J. Neurosurg.* 114(3), 613–623 (2011).
- 16 Stummer W, Pichlmeier U, Meinel T, Wiestler OD, Zanella F, Reulen HJ. Fluorescence-guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised controlled multicentre Phase III trial. *Lancet Oncol.* 7(5), 392–401 (2006).
- **Definitive study reporting the value of fluorescence-guided surgery.**
- 17 Vuorinen V, Hinkka S, Farkkila M, Jaaskelainen J. Debulking or biopsy of malignant glioma in elderly people – a randomised study. *Acta Neurochirurgica* 145(1), 5–10 (2003).
- 18 Albert FK, Forsting M, Sartor K, Adams HP, Kunze S. Early postoperative magnetic resonance imaging after resection of malignant glioma: objective evaluation of residual tumor and its influence on regrowth and prognosis. *Neurosurgery* 34(1), 45–60 (1994).
- 19 Stupp R, Hegi ME, Mason WP *et al.* Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised Phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol.* 10(5), 459–466 (2009).
- 20 Westphal M, Hilt DC, Bortey E *et al.* A Phase 3 trial of local chemotherapy with biodegradable carmustine (BCNU) wafers (Gliadel wafers) in patients with primary malignant glioma. *Neuro Oncol.* 5(2), 79–88 (2003).
- 21 Westphal M, Ram Z, Riddle V, Hilt D, Bortey E. Gliadel wafer in initial surgery for malignant glioma: long-term follow-up of a multicenter controlled trial. *Acta Neurochir. (Wien)* 148(3), 269–275 (2006).
- 22 Stummer W, Reulen HJ, Meinel T *et al.* Extent of resection and survival in glioblastoma multiforme: identification of and adjustment for bias. *Neurosurgery* 62(3), 564–576 (2008).
- 23 Stummer W, Meinel T, Ewelt C *et al.* Prospective cohort study of radiotherapy with concomitant and adjuvant temozolomide chemotherapy for glioblastoma patients with no or minimal residual enhancing tumor load after surgery. *J. Neurooncol.* 108(1), 89–97 (2012).
- **Important evidence in the debate over the value of good resections.**
- 24 Chaichana KL, Parker SL, Olivi A, Quinones-Hinojosa A. Long-term seizure outcomes in adult patients undergoing primary resection of malignant brain astrocytomas. Clinical article. *J. Neurosurg.* 111(2), 282–292 (2009).
- 25 Rosati A, Tomassini A, Pollo B *et al.* Epilepsy in cerebral glioma: timing of appearance and histological correlations. *J. Neurooncol.* 93(3), 395–400 (2009).
- 26 Vauleon E, Avril T, Collet B, Mosser J, Quillien V. Overview of cellular immunotherapy for patients with glioblastoma. *Clin. Dev. Immunol.* pii, 689171 (2010).
- 27 Welsh J, Sanan A, Gabayan AJ *et al.* GliSite brachytherapy boost as part of initial treatment of glioblastoma multiforme: a retrospective multi-institutional pilot study. *Int. J. Radiat. Oncol. Biol. Phys.* 68(1), 159–165 (2007).
- 28 Wernicke AG, Sherr DL, Schwartz TH *et al.* Feasibility and safety of GliSite brachytherapy in treatment of CNS tumors following neurosurgical resection. *J. Cancer Res. Ther.* 6(1), 65–74 (2010).
- 29 Rainov NG. A Phase III clinical evaluation of herpes simplex virus type 1 thymidine kinase and ganciclovir gene therapy as an adjuvant to surgical resection and radiation in adults with previously untreated glioblastoma multiforme. *Hum. Gene Ther.* 11(17), 2389–2401 (2000).
- 30 Immonen A, Vapalahti M, Tyynela K *et al.* AdvHSV-tk gene therapy with intravenous ganciclovir improves survival in human malignant glioma: a randomised, controlled study. *Mol. Ther.* 10(5), 967–972 (2004).
- 31 Castro MG, Candolfi M, Kroeger K *et al.* Gene therapy and targeted toxins for glioma. *Curr. Gene Ther.* 11(3), 155–180 (2011).
- 32 Mitchell P. Ark's gene therapy stumbles at the finish line. *Nat. Biotechnol.* 28(3), 183–184 (2010).
- 33 van den Bent M, Stupp R, Mason W, Mirimanoff RO, Lacombe D, Gorlia T. Impact of extent of resection on overall survival in newly-diagnosed glioblastoma after chemo-irradiation with temozolomide: further analysis of EORTC study 26981. *Eur. J. Cancer* 2, Abstract (2005).
- 34 Stummer W, van den Bent MJ, Westphal M. Cytoreductive surgery of glioblastoma as the key to successful adjuvant therapies: new arguments in an old discussion. *Acta Neurochirurgica* 153(6), 1211–1218 (2011).
- 35 Portnow J, Badie B, Chen M, Liu A, Blanchard S, Synold TW. The neuropharmacokinetics of temozolomide in patients with resectable brain tumors: potential implications for the current approach to chemoradiation. *Clin. Cancer Res.* 15(22), 7092–7098 (2009).
- 36 Rosso L, Brock CS, Gallo JM *et al.* A new model for prediction of drug distribution in tumor and normal tissues: pharmacokinetics of temozolomide in glioma patients. *Cancer Res.* 69(1), 120–127 (2009).
- 37 Navalitloha Y, Schwartz ES, Groothuis EN, Allen CV, Levy RM, Groothuis DR. Therapeutic implications of tumor interstitial fluid pressure in subcutaneous RG-2 tumors. *Neuro Oncol.* 8(3), 227–233 (2006).
- 38 Giese A, Bjerkvig R, Berens ME, Westphal M. Cost of migration: invasion of malignant gliomas and implications for treatment. *J. Clin. Oncol.* 21(8), 1624–1636 (2003).
- 39 Fung LK, Ewend MG, Sills A *et al.* Pharmacokinetics of interstitial delivery of carmustine, 4-hydroperoxycyclophosphamide, and paclitaxel from a biodegradable polymer implant in the monkey brain. *Cancer Res.* 58(4), 672–684 (1998).
- 40 Fleming AB, Saltzman WM. Pharmacokinetics of the carmustine implant. *Clin. Pharmacokinet.* 41(6), 403–419 (2002).
- 41 Li Z, Bao S, Wu Q *et al.* Hypoxia-inducible factors regulate tumorigenic capacity of glioma stem cells. *Cancer Cell* 15(6), 501–513 (2009).
- 42 Seidel S, Garvalov BK, Wirta V *et al.* A hypoxic niche regulates glioblastoma stem cells through hypoxia inducible factor 2 α . *Brain* 133(Pt 4), 983–995 (2010).
- 43 Sathornsumetee S, Cao Y, Marcello JE *et al.* Tumor angiogenic and hypoxic profiles predict radiographic response and survival in malignant astrocytoma patients treated with bevacizumab and irinotecan. *J. Clin. Oncol.* 26(2), 271–278 (2008).

- 44 Wick W, Weller M, van den Bent M, Stupp R. Bevacizumab and recurrent malignant gliomas: a European perspective. *J. Clin. Oncol.* 28(12), e188–e189 (2010).
- 45 Holash J, Maisonpierre PC, Compton D *et al.* Vessel cooption, regression, and growth in tumors mediated by angiopoietins and VEGF. *Science* 284(5422), 1994–1998 (1999).
- 46 Chan JL, Lee SW, Fraass BA *et al.* Survival and failure patterns of high-grade gliomas after three-dimensional conformal radiotherapy. *J. Clin. Oncol.* 20(6), 1635–1642 (2002).
- 47 Roberts DW, Valdes PA, Harris BT *et al.* Coregistered fluorescence-enhanced tumor resection of malignant glioma: relationships between delta-aminolevulinic acid-induced protoporphyrin IX fluorescence, magnetic resonance imaging enhancement, and neuropathological parameters. Clinical article. *J. Neurosurg.* 114(3), 595–603 (2011).
- 48 Soffietti R, Baumert BG, Bello L *et al.* Guidelines on management of low-grade gliomas: report of an EFNS-EANO task force. *Eur. J. Neurol.* 17(9), 1124–1133 (2010).
- 49 Vogelbaum MA, Jost S, Aghi MK *et al.* Application of novel response/progression measures for surgically delivered therapies for gliomas. *Neurosurgery* 70(1), 234–243 (2012).
- **Essential guidelines for defining outcome measures.**
- 50 Teixidor P, Gatignol P, Leroy M, Masuet-Aumatell C, Capelle L, Duffau H. Assessment of verbal working memory before and after surgery for low-grade glioma. *J. Neurooncol.* 81(3), 305–313 (2007).
- 51 Klein M, Duffau H, De Witt Hamer PC. Cognition and resective surgery for diffuse infiltrative glioma: an overview. *J. Neurooncol.* 108(2), 309–318 (2012).
- 52 De Witt Hamer PC, Gil Robles S, Zwinderman AH, Duffau H, Berger MS. Impact of intraoperative stimulation brain mapping on glioma surgery outcome: a meta-analysis. *J. Clin. Oncol.* 30(20), 2559–2565 (2012).
- 53 Ojemann G, Ojemann J, Lettich E, Berger M. Cortical language localization in left, dominant hemisphere. An electrical stimulation mapping investigation in 117 patients. *J. Neurosurg.* 71(3), 316–326 (1989).
- 54 McGirt MJ, Mukherjee D, Chaichana KL, Than KD, Weingart JD, Quinones-Hinojosa A. Association of surgically acquired motor and language deficits on overall survival after resection of glioblastoma multiforme. *Neurosurgery* 65(3), 463–469; discussion 469–470 (2009).
- 55 Wadley J, Dorward N, Kitchen N, Thomas D. Pre-operative planning and intra-operative guidance in modern neurosurgery: a review of 300 cases. *Ann. R. Coll. Surg. Engl.* 81(4), 217–225 (1999).
- 56 Rasmussen IA Jr, Lindseth F, Rygh OM *et al.* Functional neuronavigation combined with intra-operative 3D ultrasound: initial experiences during surgical resections close to eloquent brain areas and future directions in automatic brain shift compensation of preoperative data. *Acta Neurochirurgica* 149(4), 365–378 (2007).
- 57 Reinges MH, Nguyen HH, Krings T, Hutter BO, Rohde V, Gilsbach JM. Course of brain shift during microsurgical resection of supratentorial cerebral lesions: limits of conventional neuronavigation. *Acta Neurochirurgica* 146(4), 369–377; discussion 377 (2004).
- 58 Kuhnt D, Bauer MH, Becker A *et al.* Intraoperative visualization of fiber tracking based reconstruction of language pathways in glioma surgery. *Neurosurgery* 70(4), 911–919; discussion 919–920 (2012).
- 59 Kuhnt D, Becker A, Ganslandt O, Bauer M, Buchfelder M, Nimsky C. Correlation of the extent of tumor volume resection and patient survival in surgery of glioblastoma multiforme with high-field intraoperative MRI guidance. *Neuro Oncol.* 13(12), 1339–1348 (2011).
- 60 Bertani G, Carrabba G, Raneri F *et al.* Predictive value of inferior fronto-occipital fasciculus (IFO) DTI-fiber tracking for determining the extent of resection for surgery of frontal and temporal gliomas preoperatively. *J. Neurosurg. Sci.* 56(2), 137–143 (2012).
- 61 Gaillard WD, Berl MM. Functional magnetic resonance imaging: functional mapping. *Handb. Clin. Neurol.* 107, 387–398 (2012).
- 62 Kapsalakis IZ, Kapsalaki EZ, Gotsis ED *et al.* Preoperative evaluation with fMRI of patients with intracranial gliomas. *Radiol. Res. Pract.* 2012, 727–810 (2012).
- 63 Giussani C, Roux FE, Ojemann J, Sganzerla EP, Pirillo D, Papagno C. Is preoperative functional magnetic resonance imaging reliable for language areas mapping in brain tumor surgery? Review of language functional magnetic resonance imaging and direct cortical stimulation correlation studies. *Neurosurgery* 66(1), 113–120 (2010).
- 64 Sanai N, Mirzadeh Z, Berger MS. Functional outcome after language mapping for glioma resection. *N. Engl. J. Med.* 358(1), 18–27 (2008).
- 65 Maldonado IL, Moritz-Gasser S, de Champfleury NM, Bertram L, Moulinie G, Duffau H. Surgery for gliomas involving the left inferior parietal lobule: new insights into the functional anatomy provided by stimulation mapping in awake patients. *J. Neurosurg.* 115(4), 770–779 (2011).
- 66 Bilotta F, Rosa G. 'Anesthesia' for awake neurosurgery. *Curr. Opin Anaesthesiol.* 22(5), 560–565 (2009).
- **Emphasizing the importance of the anesthetist in awake neurosurgery.**
- 67 Unsgard G, Solheim O, Lindseth F, Selbekk T. Intra-operative imaging with 3D ultrasound in neurosurgery. *Acta Neurochir. Suppl.* 109, 181–186 (2011).
- 68 Solheim O, Selbekk T, Jakola AS, Unsgard G. Ultrasound-guided operations in unselected high-grade gliomas – overall results, impact of image quality and patient selection. *Acta Neurochirurgica* 152(11), 1873–1886 (2010).
- 69 Wirtz CR, Bonsanto MM, Knauth M *et al.* Intraoperative magnetic resonance imaging to update interactive navigation in neurosurgery: method and preliminary experience. *Comput. Aided Surg.* 2(3–4), 172–179 (1997).
- 70 Nimsky C, Ganslandt O, Cerny S, Hastreiter P, Greiner G, Fahlbusch R. Quantification of, visualization of, and compensation for brain shift using intraoperative magnetic resonance imaging. *Neurosurgery* 47(5), 1070–1079; discussion 1079–1080 (2000).
- 71 Kubben PL, ter Meulen KJ, Schijns OE, ter Laak-Poort MP, van Overbeeke JJ, van Santbrink H. Intraoperative MRI-guided resection of glioblastoma multiforme: a systematic review. *Lancet Oncol.* 12(11), 1062–1070 (2011).
- 72 Senft C, Bink A, Franz K, Vatter H, Gasser T, Seifert V. Intraoperative MRI guidance and extent of resection in glioma surgery: a randomised, controlled trial. *Lancet Oncol.* 12(11), 997–1003 (2011).
- 73 Soultz CB, Canute GS, Ryken TC. Evidence-based review of the role of reoperation in the management of malignant glioma. *Neurosurg. Focus* 4(6), e11 (1998).
- 74 Barbagallo GM, Jenkinson MD, Brodbelt AR. 'Recurrent' glioblastoma multiforme, when should we reoperate? *Br. J. Neurosurg.* 22(3), 452–455 (2003).
- 75 Mandl ES, Dirven CM, Buis DR, Postma TJ, Vandertop WP. Repeated surgery for glioblastoma multiforme: only in combination with other salvage therapy. *Surg. Neurol.* 69(5), 506–509 (2008).

- 76 van den Bent MJ, Chinot O, Boogerd W *et al.* Second-line chemotherapy with temozolomide in recurrent oligodendroglioma after PCV (procarbazine, lomustine and vincristine) chemotherapy: EORTC Brain Tumor Group Phase II study 26972. *Ann. Oncol.* 14(4), 599–602 (2003).
- 77 Van den Bent MJ, Taphoorn MJ, Brandes AA *et al.* Phase II study of first-line chemotherapy with temozolomide in recurrent oligodendroglial tumors: the European Organization for Research and Treatment of Cancer Brain Tumor Group Study 26971. *J. Clin. Oncol.* 21(13), 2525–2528 (2003).
- 78 Yung WK, Prados MD, Yaya-Tur R *et al.* Multicenter Phase II trial of temozolomide in patients with anaplastic astrocytoma or anaplastic oligoastrocytoma at first relapse. Temodal Brain Tumor Group. *J. Clin. Oncol.* 17(9), 2762–2771 (1999).
- 79 Brem H, Piantadosi S, Burger PC *et al.* Placebo-controlled trial of safety and efficacy of intraoperative controlled delivery by biodegradable polymers of chemotherapy for recurrent gliomas. The Polymer-brain Tumor Treatment Group. *Lancet* 345(8956), 1008–1012 (1995).
- 80 Stewart LA. Chemotherapy in adult high-grade glioma: a systematic review and meta-analysis of individual patient data from 12 randomised trials. *Lancet* 359(9311), 1011–1018 (2002).
- 81 Nabavi A, Thurm H, Zountsas B *et al.* Five-aminolevulinic acid for fluorescence-guided resection of recurrent malignant gliomas: a Phase II study. *Neurosurgery* 65(6), 1070–1076 (2009).
- 82 Gross CP, Herrin J, Wong N, Krumholz HM. Enrolling older persons in cancer trials: the effect of sociodemographic, protocol, and recruitment center characteristics. *J. Clin. Oncol.* 23(21), 4755–4763 (2005).
- 83 Hutchins LF, Unger JM, Crowley JJ, Coltman CA Jr, Albain KS. Underrepresentation of patients 65 years of age or older in cancer-treatment trials. *N. Engl. J. Med.* 341(27), 2061–2067 (1999).
- 84 Laws ER, Parney IF, Huang W *et al.* Survival following surgery and prognostic factors for recently diagnosed malignant glioma: data from the Glioma Outcomes Project. *J. Neurosurg.* 99(3), 467–473 (2003).
- 85 Iwamoto FM, Reiner AS, Panageas KS, Elkin EB, Abrey LE. Patterns of care in elderly glioblastoma patients. *Ann. Neurol.* 64(6), 628–634 (2008).
- 86 Kita D, Ciernik IF, Vaccarella S *et al.* Age as a predictive factor in glioblastomas: population-based study. *Neuroepidemiology* 33(1), 17–22 (2009).
- 87 Stummer W, Nestler U, Stockhammer F *et al.* Favorable outcome in the elderly cohort treated by concomitant temozolomide radiochemotherapy in a multicentric Phase II safety study of 5-ALA. *J. Neurooncol.* 103(2), 361–370 (2011).
- 88 Scott JG, Suh JH, Elson P *et al.* Aggressive treatment is appropriate for glioblastoma multiforme patients 70 years old or older: a retrospective review of 206 cases. *Neuro Oncol.* 13(4), 428–436 (2011).
- 89 Hiroko O, Kleihues P. Population-based studies on incidence, survival rates, and genetic alterations in astrocytic and oligodendroglial gliomas. *J. Neuropathol. Exp. Neurol.* 64(6), 479–489 (2005).