Background: Nursing home (NH) residents with dementia exhibit challenging behaviors or resistiveness to care (RTC) that increase staff time, stress, and NH costs. RTC is linked to elderspeak communication. Communication training (CHAT) was provided to staff to reduce their use of elderspeak.

Hypothesis: We hypothesized that CHAT would improve staff communication and subsequently reduce RTC.

Methods: Thirteen NHs were randomized to intervention and control groups. Dyads ($n = 42$) including 29 staff and 27 persons with dementia (PWD) were videorecorded during care before and/or after the intervention, and at a three month follow-up. Videos were behaviorally coded for 1) staff communication (normal, elderspeak, or silence) and 2) resident behaviors (cooperative or RTC). Linear mixed modeling was used to evaluate training effects.

Results: On average elderspeak declined from 34.6% ($SD = 18.7$) at baseline by 13.6 percentage points ($SD = 20.00$) post-intervention and 12.2 percentage points ($SD = 22.0$) at 3-month follow-up. RTC declined from 35.7% ($SD = 23.2$) by 15.3 percentage points ($SD = 32.4$) post-intervention and 13.4 percentage points ($SD = 33.7$) at 3-months. Linear mixed modeling determined that change in elderspeak was predicted by the intervention ($b = -12.20$, $p = .028$) and baseline elderspeak ($b = -0.65$, $p < .001$) while RTC change was predicted by elderspeak change ($b = 0.43$, $p < .001$); baseline RTC ($b = -0.58$, $p < .001$); and covariates.

Conclusions: A brief intervention can improve communication and reduce RTC, providing an effective nonpharmacological intervention to manage behavior and improve the quality of dementia care. No adverse events occurred.

SESSION 4625 (SYMPOSIUM)

HOSPITAL AT HOME-A DELIVERY MODEL FOR 21ST CENTURY GERIATRICS CARE

Chair: J.M. Jacobs, *Hadassah-Hebrew University Hospital, Jerusalem, Israel*

Co-Chair: G. Caplan, *Prince of Wales Hospital*

The primary purpose of Hospital at Home (HaH) is to enable the delivery of complex care at home as an alternative to in-patient care. Traditional in-patient care for older people frequently has numerous negative repercussions, and the rising number of HaH models from different Healthcare systems bears witness to this developing niche in health care delivery. Compared to usual hospital care, HaH has proven high patient satisfaction, quality of care outcomes, and evidence favors reduced mortality and overall costs. Furthermore, innovative and available technological advancements are expanding the range of treatment options available at home, and facilitating the delivery of care to a wide spectrum of increasingly more complex patients.

Yet, HaH dissemination has been limited by numerous barriers, including challenging logistics, bias towards facility-based care, and lack of payment models. Recent advances in HaH models seek to ameliorate such barriers, but they still exist. Moreover, the implementation of HaH on a widespread basis has not been well described in the research literature.

In addition to a critical review and meta-analysis of current literature, this symposium will present innovative models

of HaH providing a wide platform of care for older adults, and technologies facilitating care for specific patient groups with acute heart failure, or requiring long term mechanical ventilation. Challenges facing the implementation of widespread HaH will be addressed.

Understanding the impact of new modes of Hospital at Home care delivery, and ways at promoting implementation, is of critical importance to geriatric health service delivery in the 21st century.

A META-ANALYSIS OF HOSPITAL IN THE HOME

G. Caplan¹, N. Sulaiman¹, D. Mangin⁴, N. Aimonino Ricauda², A. Wilson³, L. Barclay¹, 1. *Department of Geriatric Medicine, Prince of Wales Hospital, Sydney, Sydney, New South Wales, Australia*, 2. *Unit of Geriatrics and Metabolic Bone Diseases, Molinette Hospital, "AOU Città della Salute e della Scienza" of Torino, Torino, Italy*, 3. *University of Leicester, Leicester, United Kingdom*, 4. *University of Otago, Christchurch, New Zealand*

This meta-analysis of Hospital-in-the-Home (HITH) compared to in-hospital care identified 61 Randomised Controlled Trials from MEDLINE, Embase, Social Sciences Citation Index, CINAHL, EconLit, PsycINFO and the Cochrane Database of Systematic Reviews. HITH care led to reduced mortality (odds ratio [OR] 0.81; 95% CI, 0.69 to 0.95; $p = 0.008$; 42 RCTs; 6992 patients), readmission rates (OR 0.75; 95% CI, 0.59 to 0.95; $p = 0.02$; 41 RCTs; 5372 patients) and cost (mean-difference 1567.11; 95% CI, 2069.53 to 1064.69; $p < 0.001$; 11 RCTs; 1215 patients). The number needed to treat at home to prevent one death was 50. Mortality data was homogenous, but heterogeneity was observed for readmission rates and cost. HITH had higher patient and carer satisfaction (21/ 22; 6/8 studies respectively); carer burden was nonsignificantly lower (8/11 studies).

HITH is associated with reductions in mortality, readmission rates and cost, and increases in patient and carer satisfaction, but no change in carer burden.

THE HOSPITAL AT HOME/MACT 'PLATFORM' MODEL—A SPECTRUM OF FACILITY-LEVEL CARE PROVIDED AT HOME

B.A. Leff¹, L. De Cherrie², A. Wajnberg², A. Federman², T. Soones², A. Brody³, A. Siu², 1. *Johns Hopkins University School of Medicine, Baltimore, Maryland*, 2. *Icahn School of Medicine, Mount Sinai, New York, New York*, 3. *New York University, New York, New York*

To date, Hospital at Home (HaH) care has focused on substitutive admission avoidance or early discharge models. In the context of a U.S. Federal innovation award program for the Center for Medicare and Medicaid Services, we have evolved the HaH model into a care "platform" for older adults. In addition to providing substitutive admission avoidance care, the platform includes: 1) "observation" stay at home; 2) acute palliative care at home; 3) acute care for hospital adverse patients (people who refuse hospital admission under all circumstances) at home; and 4) subacute rehabilitation care at home as a substitute for admission to inpatient subacute rehabilitation. This platform evolved in recognition of the evolving health care delivery system, the need to create greater demand for HaH resources, provide adequate patient