Published in final edited form as:

J Am Chem Soc. 2017 October 25; 139(42): 15259–15264. doi:10.1021/jacs.7b09491.

A Boron Protecting Group Strategy for 1,2-Azaborines

Andrew W. Baggett[†] and Shih-Yuan Liu^{*}

Department of Chemistry, Boston College, Chestnut Hill, Massachusetts 02467-3860, United States

Abstract

Upon reaction with either molecular oxygen or di-*tert*-butylperoxide in the presence of a simple copper(I) salt and an alcohol, a range of 1,2-azaborines readily exchange *B*-alkyl or *B*-aryl moieties for *B*-alkoxide fragments. This transformation allows alkyl and aryl groups to serve for the first time as removable protecting groups for the boron position of 1,2-azaborines during reactions that are not compatible with the easily modifiable *B*-alkoxide moiety. This reaction can be applied to synthesize a previously inaccessible BN isostere of ethylbenzene, a compound of interest in biomedical research. A sequence of epoxide ring opening using N-deprotonated 1,2-azaborines followed by an intramolecular version of the boron deprotection reaction can be applied to access the first examples of BN isosteres of dihydrobenzofurans and benzofurans, classes of compounds that are important to medicinal chemistry and natural product synthesis.

Graphical Abstract

1. INTRODUCTION

BN/CC isosterism of classic organic arenes¹ has emerged as a viable strategy to expand structural diversity of arenes and produce molecules with applications in biomedical research² and materials science.^{1c,d,3} In order to access a broad range of monocyclic 1,2-dihydro-1,2-azaborine substrates (abbreviated 1,2-azaborines), our group primarily utilizes late-stage functionalization strategies to diversify an assembled 1,2-azaborine core that is generated from a modified ring-closing metathesis route⁴ pioneered by Ashe.⁵ Despite recent advances including late-stage functionalization of 1,2-azaborines at the C3 position

^{*}Corresponding Author: shihyuan.liu@bc.edu.

[†]Present Address: Department of Chemistry, Linfield College, McMinnville, Oregon 97128-8626, United States

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.7b09491.

Experimental procedures, spectroscopic data (PDF)

The authors declare no competing financial interest.

via an EAS (electrophilic aromatic substitution)/Negishi coupling sequence⁶ and at the C6 position via a C-H borylation/Suzuki coupling sequence (Figure 1),⁷ the development of applications of these heterocycles is still hampered by the lack of further viable functionalization strategies.

Manipulation of the boron position of azaborines forms the core of our synthetic efforts as the nature of the boron substituent greatly influences properties of azaborines such as solubility, volatility, stability toward air and moisture,⁸ reactivity toward nucleophiles, and stability during the course of relatively harsh transformations such as Suzuki coupling. For example, *C6*-B(pin) azaborines bearing alkyl or aryl groups at the ring boron position react exclusively at the B(pin) moiety during Suzuki couplings with a diverse range of aryl bromide coupling partners, but *C6*-B(pin) azaborines bearing alkoxy groups or hydrogen at the ring boron position only tolerate a greatly restricted set of cross coupling conditions and coupling partners.⁷ Conversely, azaborines bearing alkoxy groups, chloride, or hydrogen at boron readily react at boron via nucleophilic addition/elimination,^{4,9,10} rhodium-catalyzed arylation,¹¹ or rhodium-catalyzed dehydrogenative borylation,¹² whereas azaborines bearing alkyl or aryl groups at boron do not undergo these reactions.

In contrast to the abundance of available transformations of *B*-alkoxy, *B*-Cl, or *B*-H azaborines into *B*-alkyl or *B*-aryl azaborines, previous examples¹³ of the reverse transformation are restricted to polycyclic azaborines and are narrow in scope (Scheme 1, eq 1). The development of a more general transformation for monocyclic azaborines—and perhaps for boron-containing heteroarenes of any type—is highly desirable because it would enable alkyl or aryl groups to serve as removable boron protecting groups during reaction sequences requiring synthetic steps incompatible with azaborines bearing labile substituents at boron (Scheme 1, eq 2). We envisioned that a reasonable strategy to carry out this transformation would involve oxidative conversion of a *B*-alkyl or *B*-aryl azaborine to a versatile *B*-alkoxy azaborine.¹⁴ Conversion of alkyl boranes to alkyl boronates through the use of hydrogen peroxide reagents¹⁵ is a cornerstone transformation in the synthesis of alcohols from alkenes,¹⁶ but application of similar conditions to azaborine substrates results in decomposition of the aromatic heterocycles.

A report from Jiao and co-workers describing the relatively mild copper-catalyzed oxidation of aryl ketones to aryl esters using molecular oxygen as the sole oxidant caught our attention as providing a potentially viable alternative (Scheme 2, eq 1).¹⁷ We postulated that perceived similarities between the reactivity of the electrophilic carbonyl carbon involved in this transformation (Scheme 2, eq 2) and the known reactivity of the electrophilic boron atom of azaborines (Scheme 2, eq 3) may allow an analogous transformation to occur for azaborine substrates under similar conditions using a copper(I) source and either molecular oxygen or a dialkylperoxide¹⁸ as the stoichiometric oxidant. Herein we describe the successful application of a copper-catalyzed protocol to carry out the azaborine *B*-R to *B*-alkoxy transformation while leaving the azaborine ring intact. We describe our initial investigations of the mechanistic features of this transformation, how our experiments reveal a mechanistic picture considerably different from what we expected based on the work of Jiao, and the applications of this method to access azaborines that require removal of a *B*-alkyl group as a key synthetic step.

2. RESULTS AND DISCUSSION

At the outset of this project, we tested the activity of the readily available *N*-H, *B*-*n*-Bu azaborine **1**¹⁴ under conditions¹⁷ adapted from Jiao and co-workers (Scheme 3). We were delighted to find that a ¹¹B NMR spectrum of the reaction revealed the complete consumption of *B*-butyl azaborine **1** (¹¹B shift ~35 ppm) and appearance of a peak corresponding with the desired *B*-butoxy material **2**¹⁴ (¹¹B shift ~30 ppm) and a minor peak assigned to one or more B(OR)₃ species produced as undesired side products (¹¹B shift ~20 ppm).

Encouraged by this promising result, we explored the consequences of modifying various reaction parameters. We replaced *n*-butanol with *n*-dodecanol in the reaction for two reasons: the longer alkyl chain grants increased stability toward silica gel chromatography to the *B*-alkoxide product **3**, and we were able to distinguish products arising from incorporation of the alcohol versus those arising from potential net oxygen insertion. After brief preliminary reaction optimizations including a wide survey of alternate metal salts in place of copper(I) bromide (Tables S1 and S2) resulting in the standard conditions shown in entry 1 of Table 1, we confirmed that a supply of oxygen was necessary for reactivity (entries 1–2) and that a sealed system with pure oxygen led to a cleaner reaction than those run sealed with air or performed open to air at reflux (entries 3–4). Removing the *n*-dodecanol additive from the system led to an inefficient production of **2**, indicating that a net oxygen insertion is operative to some extent in production of this species (entry 5). Adding *n*-butanol to the reaction granted **2** in a yield comparable to the total yield of *B*-OR material obtained from the standard reaction conditions, demonstrating that an alcohol additive is essential to achieve a relatively clean reaction (entry 6). Removing the copper (entry 7) or everything besides **1**, the solvent, and oxygen (entry 8) resulted in high conversion of **1** but poor yield of *B*-OR compounds **2** or **3**, showing a largely unfavorable direct reaction between molecular oxygen and the azaborine¹⁹ that was either suppressed or outpaced in the presence of the copper and alcohol additives. The *B*-OR products demonstrated a high degree of oxygen stability compared to that of *B*-alkyl **1**, since the yield of **2** and **3** remained high after extended reaction time (entry 9). Finally, adding butylated hydroxytoluene (BHT) to the reaction strongly suppressed the conversion of **1** to any product or byproduct, suggesting that radical pathways are essential to the conversion of **1** in the presence of molecular oxygen (entry 10).

Although we demonstrated that the desired *B*-R to *B*-OR transformation was possible under the aerobic conditions inspired by Jiao and co-workers, the best yields were only modest due to competitive decomposition of the azaborine material to B(OR)₃ species during the course of the reaction. Furthermore, we failed to detect any aldehyde byproducts that would be expected if the mechanism proposed by Jiao was operative in our case and suspected the carbon leaving group on the azaborine may instead be expelled as a radical during the progress of the reaction. The lessons we learned from these aerobic reactions guided us to explore an alternate system. First, we used di-*tert*-butylperoxide (DTBP)²⁰ under a nitrogen atmosphere as the oxidant to potentially reduce the prevalence of unwanted side reactions between the azaborine substrate and molecular oxygen. Second, we examined the benzyl group as the boron leaving group, since it should impart the same stability toward cross

coupling or other nucleophilic reaction conditions that *n*-butyl would, and it would become a relatively stable radical species in case this expulsion contributed to limiting the rate of the overall reaction.

Our synthesis of *B*-benzyl azaborines followed straightforward protocols involving addition of a benzyl Grignard reagent to *N*-TBS, *B*-Cl azaborine **4**⁴ to generate the *N*-TBS, *B*-benzyl azaborines **5** and **6** on gram scale in excellent yield. Both could undergo silyl deprotection with TBAF in good yield on gram scale to yield *N*-H, *B*-benzyl azaborines **7** and **8** (Scheme 4).

With *N*-H, *B*-benzyl substrate **7** in hand, we investigated the new set of conditions as illustrated in Table 2. Switching oxidants and substrates resulted in a cleaner transformation to the desired product **3** with only a small amount of the *B*-O-*t*-Bu product **9** observed and a nearly stoichiometric amount of bibenzyl **10** formed as a byproduct (entry 1). In contrast to the reaction using molecular oxygen, removal of the copper nearly completely suppressed the conversion of **7** (entry 2), and some conversion was only seen when heating the system to 130 °C (entry 3). The generation of bibenzyl was also suppressed upon removal of CuBr. Removal of pyridine or the substitution of pyridine with triethylamine or bipyridine resulted in similarly low conversions and yields for the reaction (entries 4–6). In the absence of *n*-dodecanol the *B*-O-*tert*-butyl product **9** was formed via incorporation of one-half of DTBP into the azaborine (entry 7). Without the oxidant present, the desired reaction did not occur, and only minimal conversion took place (entry 8). The reaction required elevated temperatures to occur (entry 9), but could be run 40 °C lower than the reactions described in Table 1. When we replaced DTBP with *tert*-butyl hydrogen peroxide (TBHP) we observed a less efficient reaction with a worse gap between conversion and yield of **3** (entry 10). We ran a small number of experiments on *B*-benzyl substrate **7** with other copper sources and with molecular O₂ as oxidant, but observed less efficient reactions in all cases (Table S3).

The experiments in Table 2 and Table S3 and the detailed mechanistic studies performed by others^{18,20} prompt us to propose a tentative mechanism for the *B*-R to *B*-OR transformation using the system based on copper and DTBP (Figure 2). We propose an initial generation of a Cu(II) alkoxide species from the reaction of a Cu(I) species with the DTBP oxidant with concomitant formation of a *tert*-butoxide radical.²¹ Next, the alkoxide radical attacks the azaborine boron atom via a one-electron pathway to generate azaborine radical **11**. We believe that the Cu metal may serve as a reservoir for the *tert*-butoxide radical. A Cu(I) species may then mediate the removal of the boron benzyl group to generate the azaborine product **9**, which exchanges alkoxide groups with the *n*-dodecanol in solution to generate the ultimate azaborine product **3**. The copper(II) benzyl species could finally mediate removal of a second benzyl group to generate another equivalent of **9** and a Cu(III) bisalkyl species²² that could produce bibenzyl by reductive elimination and regeneration of a Cu(I) species.

With a set of optimized reaction conditions in hand, we explored the substrate scope with respect to the azaborine reagent in this transformation (Table 3). Substrates with *N*-TBS groups gave good isolated yields in comparison to those with *N*-H groups (entries 1 vs 2) likely due to the greater tendency of *N*-H, *B*-OR azaborines to bind to silica gel compared to the low silica gel affinity of the silylated substrates. The only bibenzyl species observed

following the deprotection of **8** was the symmetrical product bearing two OMe groups, demonstrating that there is no crossover between the benzyl leaving group and the toluene solvent during the course of this reaction (entry 3). Azaborines bearing alkyl and aryl leaving groups also underwent the deprotection, albeit with reduced yields compared to substrates bearing benzyl groups (entries 4–10). The ability of the leaving group to stabilize a radical seems to correlate with the degree of success of the reaction. Azaborines bearing either a bromine or an *n*-propyl group at the C3-position and a benzyl leaving group underwent the reaction in modest yield (entries 11–12). Finally, an azaborine bearing a mesityl group at boron failed to undergo the reaction, likely due to the steric bulk of the mesityl group preventing nucleophilic attack at boron.

With our new protocol in place for the boron deprotection of *B*-benzyl azaborines, we used this strategy as a key step in the synthesis of *C6*-ethyl BN-ethylbenzene **25** (Figure 3, eq 1). Our group has investigated the activity of BN-ethylbenzenes **23** and **24** toward ethylbenzene dehydrogenase^{2b} as well as ligands for engineered T4 lysozymes,^{2a,e} and we desire access to more regioisomers of BN-ethylbenzene to develop a systematic structure-activity relationship analysis of the effects of BN/CC isosterism in biological settings. A straightforward strategy⁷ involving the cross coupling between vinyl bromide and *C6*-borylated azaborines with labile groups already installed at the boron atom failed due to the sensitivity of these azaborines toward the reaction conditions (Figure 3, eq 2).

Initial borylation of *N*-H, *B*-benzyl **7** under standard iridiumcatalyzed borylation conditions²³ granted *C6*-B(pin) azaborine **26** in excellent yield, and this substrate underwent cross coupling without difficulty to generate *C6*-vinyl azaborine **27** (Scheme 5). Compound **27** was hydrogenated under mild conditions to produce *C6*-ethyl azaborine **28**, which cleanly underwent the boron deprotection protocol to grant access to *B*-OR azaborine **29**. As the final step in the synthesis, we reduced **29** to the target compound **25** using LiAlH₄, followed by a mild acid, and isolated over 100 mg of the volatile *C6*-ethyl, *N*-H, *B*-H azaborine.

We next employed an intramolecular variant of this method to generate BN-dihydrobenzofurans, which serve as BN/CC isosteres of the dihydrobenzofuran moiety found in a variety of natural products and drugs²⁴ and as potential precursors to BN-benzofurans. To the best of our knowledge, there are no examples of BN-isosteres of either class of compounds. The synthesis of BN-dihydrobenzofurans involved a two-step process: a deprotonation/epoxide ring opening sequence starting from azaborine **7**, followed by the copper-catalyzed boron oxidation reaction which resulted in cyclization (Table 4). Six BN-dihydrobenzofuran isosteres were produced in good yield over two steps with this method including the parent compound **36**, demonstrating the potential for this method to be used to generate a library of BN-dihydrobenzofurans by varying the substitution on the epoxide reaction partner.

Next, we explored the oxidation of several BN-dihydrobenzofurans to the corresponding BN-benzofurans (Scheme 6). Phenyl-substituted **39** was converted to the substituted BN-benzofuran **42** in good yield after refluxing in toluene in the presence of Pd/C for 24 h; however, only limited success was observed when the same conditions were applied to the

other substrates. Vinyl-substituted **37** underwent isomerization rather than oxidation, leading to the isolation of **43** in modest yield. Neither **36** nor **40** was converted to the corresponding BN-benzofurans under these conditions, suggesting that the successful observed reactions may be due to the driving force resulting from extension of π conjugation in **42** and **43**.

3. CONCLUSION

We developed two copper-catalyzed reaction systems using either molecular oxygen or DTBP as the stoichiometric oxidant that convert *B*-alkyl or *B*-aryl azaborine substrates to *B*-OR materials that can undergo further functionalization at the boron atom following the transformation. Both reactions are assisted by the presence of a copper salt as an additive, but both reactions still take place without the copper, albeit with a significant loss of efficiency. No metal or Lewis acid additive besides copper was able to efficiently promote the reaction, and we determined that a key species involved in the azaborine oxidation was most likely an alkoxy radical. We investigated the mechanistic features of the reactions by designing straightforward experiments and discovering that bibenzyl is formed as a stoichiometric byproduct during the deprotection of *B*-benzyl substrates. The optimized deprotection protocol is highly effective for the removal of benzyl groups from azaborines, presumably because the benzyl radical that is expelled is relatively stabilized compared to other alkyl or aryl radicals. We applied the new method as the key step in the synthesis of *C6*-ethyl BN-ethylbenzene and in the synthesis of BN-dihydrobenzofurans and BN-benzofurans which represent unexplored classes of heterocycles. This method will be highly useful to chemists carrying out multistep transformations of azaborines in particular and perhaps boron heteroarenes in general that require removal of carbon-based boron protecting groups as part of the synthetic plans.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This research was supported by the National Institutes of Health NIGMS (R01-GM094541) and Boston College start-up funds. A.W.B. thanks the LaMattina Family Graduate Fellowship in Chemical Synthesis for support.

REFERENCES

- (1). An overview: Bosdet MJD; Piers WE *Can. J. Chem* 2009, 87, 8–29. Campbell PG; Marwitz AJV; Liu S-Y *Angew. Chem., Int. Ed* 2012, 51, 6074–6092. Wang X-Y; Wang J-Y; Pei, J. *Chem. - Eur. J* 2015, 21, 3528–3539. Morgan MM; Piers WE *Dalton Trans* 2016, 45, 5920–5924. Liu Z; Marder TB *Angew. Chem., Int. Ed* 2008, 47, 242–244.
- (2). Liu L; Marwitz AJV; Matthews BW; Liu S-Y *Angew. Chem., Int. Ed* 2009, 48, 6817–6819. Knack DH; Marshall JL; Harlow GP; Dudzik A; Szaleniec M; Liu S-Y; Heider J *Angew. Chem., Int. Ed* 2013, 52, 2599–2601. Vlasceanu A; Jessing M; Kilburn JP *Bioorg. Med. Chem* 2015, 23, 4453–4461. Rombouts FJR; Tovar F; Austin N; Tresadern G; Trabanco AA *J. Med. Chem* 2015, 58, 9287–9295. Lee H; Fischer M; Shoichet BK; Liu SY *J. Am. Chem. Soc* 2016, 138, 12021–12024. Zhao P; Nettleton DO; Karki RG; Zecri FJ; Liu SY *ChemMedChem* 2017, 12, 358–361.
- (3). Taniguchi T; Yamaguchi S *Organometallics* 2010, 29, 5732–5735. Marwitz AJV; Jenkins JT; Zakharov LN; Liu S-Y *Angew. Chem., Int. Ed* 2010, 49, 7444–7447. Marwitz AJV; Lamm AN; Zakharov LN; Vasiliu M; Dixon DA; Liu S-Y *Chem. Sci* 2012, 3, 825–829. Braunschweig H;

- Horl C; Mailander L; Radacki K; Wahler, J. Chem. - Eur. J 2014, 20, 9858–9861. Baggett AW; Guo F; Liu S-Y; Jakle F Angew. Chem., Int. Ed 2015, 54, 11191–11195. Wan WM; Baggett AW; Cheng F; Lin H; Liu SY; Jakle F Chem. Commun 2016, 52, 13616–13619.
- (4). Marwitz AJV; Abbey ER; Jenkins JT; Zakharov LN; Liu S-Y Org. Lett 2007, 9, 4905–4908. [PubMed: 17944483]
- (5). Ashe AJ, III; Fang X Org. Lett 2000, 2, 2089–2091. [PubMed: 10891237]
- (6). Pan J; Kampf JW; Ashe AJ, III Org. Lett 2007, 9, 679–681. Brown AN; Li B; Liu S-Y J. Am. Chem. Soc 2015, 137, 8932–8935. [PubMed: 17249680]
- (7). Baggett AW; Vasiliu M; Li B; Dixon DA; Liu S-YJ Am. Chem. Soc 2015, 137, 5536–5541.
- (8). Lamm AN; Liu S-Y Mol. BioSyst 2009, 5, 1303–1305. [PubMed: 19823745]
- (9). Marwitz AJV; Matus MH; Zakharov LN; Dixon DA; Liu S-Y Angew. Chem., Int. Ed 2009, 48, 973–977.
- (10). Lamm AN; Garner EB; Dixon DA; Liu S-Y Angew. Chem., Int. Ed 2011, 50, 8157–8160.
- (11). Rudebusch GE; Zakharov LN; Liu S-Y Angew. Chem., Int. Ed 2013, 52, 9316–9319.
- (12). Brown AN; Zakharov LN; Mikulas T; Dixon DA; Liu S-Y Org. Lett 2014, 16, 3340–3343. [PubMed: 24916200]
- (13). Paetzold PI; Stohr G; Maisch H; Lenz H Chem. Ber 1968, 101, 2881–2888. Molander GA; Wisniewski SR J. Org. Chem 2014, 79, 8339–8347. Li G; Xiong W-W; Gu P-Y; Cao J; Zhu J; Ganguly R; Li Y; Grimsdale AC; Zhang Q Org. Lett 2015, 17, 560–563.
- (14). Abbey ER; Lamm AN; Baggett AW; Zakharov LN; Liu S-YJ Am. Chem. Soc 2013, 135, 12908–12913.
- (15). Johnson JR; Van Campen MG, Jr. J. Am. Chem. Soc 1938, 60, 121–124.
- (16). Brown HC; Rao BC S. J. Am. Chem. Soc 1956, 78, 5694–5695.
- (17). Huang X; Li X; Zou M; Song S; Tang C; Yuan Y; Jiao NJ Am. Chem. Soc 2014, 136, 14858–14865.
- (18). Jang ES; McMullin CL; Kas M; Meyer K; Cundari TR; Warren TH J. Am. Chem. Soc 2014, 136, 10930–10940. [PubMed: 24940616]
- (19). Nozaki K; Oshima K; Utimoto K J. Am. Chem. Soc 1987, 109, 2547–2549. Ollivier C; Renaud P Chem. Rev 2001, 101, 3415–3434.
- (20). For recent work using the Cu/DTBP reagent combination, see: Tran BL; Driess M; Hartwig JF J. Am. Chem. Soc 2014, 136, 17292–17301. Chikkade PK; Kuninobu Y; Kanai M Chem. Sci 2015, 6, 3195–3200. Ha TM; Chatalova-Sazepin C; Wang Q; Zhu J Angew. Chem., Int. Ed 2016, 55, 9249–9252. [PubMed: 25389772]
- (21). Gephart RT, III; McMullin CL; Sapiezynski NG; Jang ES; Aguila MJB; Cundari TR; Warren TH J. Am. Chem. Soc 2012, 134, 17350–17353. [PubMed: 23009158]
- (22). Yoshikai N; Nakamura E Chem. Rev 2012, 112, 2339–2372. Shen R; Iwasaki T; Terao J; Kambe N Chem. Commun 2012, 48, 9313–9315. Iwasaki T; Imanishi R; Shimizu R; Kuniyasu H; Terao J; Kambe N J. Org. Chem 2014, 79, 8522–8532. [PubMed: 22111574]
- (23). Ishiyama T; Takagi J; Hartwig JF; Miyauchi N Angew. Chem., Int. Ed 2002, 41, 3056–3058.
- (24). Sneader W Drug Discovery: A History; Wiley: Hoboken, NJ, 2005; p 119 Adams H; Gilmore NJ; Jones S; Muldowney MP; von Reuss SH; Vemula R Org. Lett 2008, 10, 1457–1460. Koizumi Y; Kobayashi H; Wakimoto T; Furuta T; Fukuyama T; Kan T J. Am. Chem. Soc 2008, 130, 16854–16855.

Figure 1.
Ring position labeling of 1,2-azaborines.

Figure 3.
BN-ethylbenzene analogs.

Scheme 1. Removal of *B*-Alkyl or *B*-Aryl Groups from Azaborines

Mechanism proposed by Jiao

Conceptual similarity of BN system

Scheme 2.
 Mechanistic Inspiration

Scheme 3.
Initial *B*-Alkyl to *B*-Alkoxy Transformation

Scheme 4.
Generation of *B*-Benzyl Azaborines

Scheme 5.
Synthesis of *C*6-Ethyl BN-Ethylbenzene

Scheme 6.
Oxidation to BN-Benzofurans

Table 1.

Reaction Using Molecular Oxygen As Oxidant

Entry	Change from Standard Conditions	Conversion of 1 (%) ^a	Yield of 2 (%) ^a	Yield of 3 (%) ^a
1	none	94	15	64
2	nitrogen atmosphere	2	0	0
3	air	44	4	32
4	air, reflux	53	5	42
5	<i>n</i> -dodecanol removed	77	25	0
6	<i>n</i> -butanol instead of <i>n</i> -dodecanol	89	73	0
7	CuBr removed	79	5	39
8	CuBr, alcohol, and pyridine removed	74	14	0
9	reaction run for 2 h	100	14	64
10	BHT (2.0 equiv) added	5	0	4

^aConversion and yield obtained by GC using dodecane as a calibrated internal standard.

Table 2.

CuBr/DTBP-Catalyzed Boron Deprotection of 7

Entry	Change from Standard Conditions	Conversion of 7 (%) ^a	Yield of 9 (%) ^a	Yield of 3 (%) ^a	Yield of 10 (%) ^a
1	none	100	1	75	81
2	CuBr removed	1	0	0	1
3	CuBr removed, run at 130 °C	14	0	5	7
4	pyridine removed	24	0	15	23
5	NEt ₃ instead of pyridine	16	0	9	16
6	bipy (1.0 equiv), no pyridine	72	1	48	60
7	<i>n</i> -dodecanol removed	81	59	0	59
8	DTBP removed	2	0	0	2
9	reaction run for 2 h at RT	0	0	0	0
10	TBHP instead of DTBP	76	0	42	37

^aConversion and yield obtained by GC using dodecane as a calibrated internal standard.

Table 3.

Scope of Deprotection Method

Entry	Starting Material	Product	% Yield Isolated ^[b] (NMR) ^[c]
1	 5	 12	90 (99)
2	 7	 3	66 (97)
3	 8	3	61 (94)
4	 13	12	69 (70)
5	 1	3	59 (78)
6	 14	12	54 (66)
7	 15	12	49 (57)

Entry	Starting Material	Product	% Yield Isolated ^b (NMR) ^c
8	16	3	35 (57)
9	17	12	37 (53)
10	18	12	54 (59)
11	19	20	40 (57)
12	21	22	53 (65)

^a Azaborines with R² = mesityl failed to undergo the reaction.

^b Isolated yields are an average of two runs.

^c NMR yields obtained using 1,5-cyclooctadiene (cod) as a calibrated internal standard. All NMR yields are an average of two runs.

Table 4.

Synthesis of BN-Dihydrobenzofurans

entry	epoxide	ring-opened product (% Yield) ^[a]	BN-dihydrobenzofuran (% Yield) ^[a]
1		 30 97% Yield	 36 81% Yield
2		 31 71% Yield	 37 77% Yield
3		 32 81% Yield	 38 88% Yield
4		 33 80% Yield	 39 79% Yield

entry	epoxide	ring-opened product (% Yield) ^[a]	BN-dihydrobenzofuran (% Yield) ^[a]
5		34 95% Yield	40 84% Yield
6		35 40% Yield	41 26% Yield

^aYields shown are isolated yields and are the average of two runs.