Published in final edited form as:

J Org Chem. 1998 September 18; 63(19): 6735–6738. doi:10.1021/jo9808066.

Stereoselective Synthesis of Protected Thymine Polyoxin C via [2,3]-Wittig-Still Rearrangement of Ribose-Derived Allylic Stannyl Ethers

Arun K. Ghosh and Yong Wang

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607

The [2,3]-Wittig rearrangement leading to stereocontrolled carbon-carbon bond formation has been increasingly utilized in organic synthesis.¹ The synthetic potential of this reaction is extended even further in Still's variant of the [2,3]-Wittig rearrangement (Wittig-Still rearrangement) where unstable oxymethyl lithium is generated at low temperature by a tin-lithium exchange reaction.² Recently, we reported a stereoselective synthesis of an antifungal nucleoside, sinefungin.³ In our continuing interest in the development of synthetic methodologies for bioactive amino acid nucleosides, we have investigated the [2,3]-Wittig rearrangement of *E* and *Z*-allylic stannyl ethers derived from an isopropylidene D-ribose derivative. Herein we report that the [2,3]-rearrangement of *Z*-allylic stannyl ether proceeded with nearly complete syn selectivity and excellent isolated yield. The resulting [2,3]-rearranged product was converted to the protected thymine polyoxin C, which is a basic component of many bioactive polyoxins, including polyoxin J (**1**). Polyoxins are important antifungal agents with chitin synthetase inhibitory properties.⁴ Synthesis and biological evaluation of polyoxins and their variants have been the subject of immense interest over the years.^{5,6} Early syntheses of polyoxin nucleosides via cyanohydrin formation at the C-5' aldehyde of uridine have been reported.^{5g,h} A very useful strategy to polyoxin C and other glycosyl R-amino acids has been developed from D-serinal.^{5c} Protected ribose-derived nitro olefin and nitron-based strategies have also been utilized in the synthesis of polyoxin C.^{5b,d} Thymine polyoxin has been synthesized from myoinositol and other noncarbohydrate precursors as well.^{5a,e,6c} Our approach to polyoxin C synthesis is based upon the chain lengthening of sugars by a highly stereoselective [2,3]-Wittig rearrangement.

The known⁷ methyl glycoside **2** was readily converted to *trans*- α,β -unsaturated ester **3** by Swern oxidation followed by Horner-Emmons olefination with sodium hydride and triethyl phosphonoacetate as described previously.³ The *cis*- α,β -unsaturated ester **4** was prepared selectively by Still's variant of Horner-Emmons olefination with bis(2,2,2-trifluoroethyl)methoxycarbonylmethyl phosphonate.⁸ Reduction of these esters with DIBAL in CH₂Cl₂ at -78 °C for 2 h provided the respective *trans*- and *cis*-allylic alcohols **5** and **6** in excellent yields (Scheme 1). Etherification of alcohols **5** and **6** with potassium hydride and

Supporting Information Available: ¹H NMR spectra for compounds **6-8**, **12**, and **14-15** and ¹³C NMR spectra for **10** and **15** (8 pages). This material is contained in libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.
JO9808066

$\text{Bu}_3\text{SnCH}_2\text{I}$ in the presence of catalytic amounts of ${}^n\text{Bu}_4\text{N}^+\text{I}^-$ afforded the respective *E*- and *Z*-stannylated ethers **7** and **8**. Tin lithium exchange of *E*-stannyl ether **7** with ${}^n\text{BuLi}$ at $-78\text{ }^\circ\text{C}$ in THF resulted in a syn/anti mixture (4.4:1 by 400MHz ${}^1\text{H}$ NMR) of rearranged product **9** and its epimer in 84% yield after silica gel chromatography. On the other hand, the reaction of *Z*-stannyl ether **8** with ${}^n\text{BuLi}$ provided exclusively the syn product **9** in 74% isolated yield. ${}^1\text{H}$ NMR (400 MHz) revealed the presence of a single isomer.

The initial stereochemical assignment of the [2,3]rearranged syn product **9** has been made on the basis of previous stereochemical studies of [2,3]-Wittig-Still rearrangement by Brückner and co-workers.⁹ Further evidence of this stereochemical assignment was provided after the syn-rearranged product **9** was converted to thymine polyoxin C. The stereochemical assignment is also consistent with “Houk-like” transition-state models as shown in Figure 1.¹⁰ In these models, the allylic C-O bond is orthogonal to the plane of the allylic CdC and is antiperiplanar with respect to the impending carbanion. Consistent with these models, both compounds **7** (*E*-isomer) and **8** (*Z*-isomer) provided the same major rearranged product **9** through the more favorable transition states **A** and **C**, respectively. Between the corresponding diastereomeric transition states **B** (*E*-isomer) and **D** (*Z*-isomer), transition state **D** is highly disfavored because of severe developing allylic [1,3] strain. This may explain why compound **8** (*Z*-isomer) provided exclusively syn-rearranged product **9** compared to compound **7** (*E*-isomer), which has exhibited a syn/anti product ratio of 4.4:1.

As shown in Scheme 2, alcohol **9** was converted to Cbzprotected amine derivative **11** by oxidation followed by Curtius rearrangement of the resulting carboxylic acid. Thus, Swern oxidation of **9** provided the corresponding aldehyde which, upon exposure to sodium chlorite in *tert*-butyl alcohol, afforded the acid **10**. It is important to note that the use of triethylamine in the standard Swern oxidation protocol resulted in a substantial amount of epimerization of the resulting β,γ -unsaturated carboxaldehyde. This epimerization was circumvented by the use of diisopropylethylamine instead of triethylamine. The above acid **10** without further purification was exposed to diisopropylethylamine, methyl chloroformate, and sodium azide at $0\text{ }^\circ\text{C}$. The resulting acyl azide was dissolved in toluene and heated at $100\text{ }^\circ\text{C}$ for 1 h. After this period, benzyl alcohol was added and the mixture was heated at reflux for 12 h to furnish the Cbz derivative **11** in 68% yield (from **9**) after silica gel chromatography. Since the Curtius rearrangement proceeds with retention of configuration of the migrating carbon atom, the stereochemistry of the urethane-bearing chiral center in **11** was assigned accordingly.¹¹ Of particular note, the attempted Curtius rearrangement of **10** with diphenylphosphoryl azide and triethylamine followed by treatment with benzyl alcohol and subsequent heating at reflux for 10 h resulted in α - β -unsaturated ester **12** as the major product, along with a small amount of Cbz derivative **11**.¹² For conversion of the vinyl group in **11** to methyl ester **13**, ozonolytic cleavage was carried out in methanolic sodium hydroxide solution to afford **13** in 94% yield after chromatography.¹³

To install thymine at the anomeric center, the protecting group interconversion of **13** to triacetate **14** was carried out in a three-step sequence: (1) removal of the isopropylidene group with Dowex 50W H^+ in methanol, (2) acetylation of the resulting diol with acetic anhydride in pyridine, and (3) acetal exchange by treatment with acetic anhydride and a

catalytic amount of concentrated sulfuric acid in a mixture of glacial acetic acid and CH_2Cl_2 at 23 °C for 2 h (70%, 2:1 mixture of anomers). Triacetate **14** was exposed to Vorbrüggen reaction conditions¹⁴ with 5-methyl-2,4-bis(trimethylsilyloxy)pyrimidine in the presence of TMSOTf in dichloroethane at 45 °C for 2 h to afford the protected β -nucleoside **15** (mp 81–83 °C) in 80% yield after silica gel chromatography. The spectral properties (^1H and ^{13}C NMR) of **15** ($[\alpha]^{23}_{\text{D}} +17.1$, c 0.45, CHCl_3 ; lit.^{5a} $[\alpha]^{23}_{\text{D}} +19.8$, c 0.56, CHCl_3) are in full agreement with the reported values.

In conclusion, a stereoselective synthesis of protected thymine polyoxin C has been accomplished starting from D-ribose. The key steps in the synthesis are a highly diastereoselective [2,3]-sigmatropic reaction of *Z*-allylic stannyl ethers, followed by conversion of the resulting hydroxymethyl group to the Cbz-protected amine derivative by a Curtius rearrangement. The stereoelectronic effects of the [2,3]-Wittig rearrangement of the ribosederived *E*- and *Z*-stannyl ethers are noteworthy. Further application of this stereoselective [2,3]-Wittig-Still rearrangement in synthesis is in progress.

Experimental Section

All melting points were recorded and uncorrected. Anhydrous solvents were obtained as follows: 1,2-dichloroethane was first refluxed for 2 h over P_2O_5 and then distilled, tetrahydrofuran was distilled from sodium and benzophenone, methylene chloride was distilled from CaH_2 and trimethylchlorosilane and pyridine were distilled from CaH_2 . All other solvents were HPLC grade. Column chromatography was performed with Whatman 240400 mesh silica gel under a low pressure of 5–10 psi. Thinlayer chromatography (TLC) was carried out with Merck silica gel 60 F-254 plates.

Methyl (*E*)-5,6-Dideoxy-2,3-*O*-isopropylidene- β -D-ribohept-5-enofuranoside (**5**).

To a stirred solution of ester **3**³ (1.7 g, 6.62 mmol) in CH_2Cl_2 (10 mL) at –78 °C was added DIBAL (1 M in hexane, 19 mL, 19 mmol) solution, and the resulting mixture was stirred at –78 °C for 2 h. After this period, the reaction was carefully quenched with water (3 mL) and the mixture was allowed to warm to 23 °C. The mixture was filtered through a glass wool plug, and the solid was rinsed several times with CH_2Cl_2 and water. The combined filtrate was transferred to a separatory funnel, and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (2×20 mL), and the combined organic layers were dried over anhydrous Na_2SO_4 . Evaporation of the solvent gave a residue which was purified on a silica gel column (50% ethyl acetate/hexane) to afford **5** as a colorless oil (1.335 g, 92%): $[\text{R}]^{23}_{\text{D}} -42.2$ (c 0.83, CHCl_3); IR (neat) 3423, 2950, 2930, 1630, 1374, 1088 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 5.82 (m, 2H), 4.98 (s, 1H), 4.66 (d, 1H, J 8.2 Hz), 4.61 (s, 2H), 4.15 (t, 1H, J 5.4 Hz), 3.34 (s, 3H), 1.49 (s, 3H), 1.31 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 24.9, 26.4, 54.7, 62.6, 84.5, 85.4, 87.4, 109.3, 112.3, 130.4, 132.4; MS (EI) m/z 230 (M^+), 215 ($\text{M}^+ - \text{Me}$), 198, 155. Anal. Calcd for $\text{C}_{11}\text{H}_{18}\text{O}_5$: C, 57.38; H, 7.88. Found: C, 57.22; H, 7.94.

Methyl (*Z*)-5,6-Dideoxy-2,3-*O*-isopropylidene- β -D-ribohept-5-enofuranoside (**6**).

To a stirred solution of ester **4**⁸ (1.07 g, 4.15 mmol) in CH_2Cl_2 (15 mL) at –78 °C was added DIBAL (1.0 M in hexane, 12.4 mL, 12.4 mmol) solution, and the resulting mixture was

stirred for 3 h at -78°C . After this period, the reaction was worked up as described for **5** to furnish the allyl alcohol **6** as a colorless oil (930 mg, 95%): ^1H NMR (400 MHz, CDCl_3) δ 5.74 (m, 1H), 5.61 (m, 1H), 4.97 (s, 1H), 4.95 (d, 1H, J 9.5 Hz), 4.62 (d, 1H, J 5.9 Hz), 4.58 (d, 1H, J 5.9 Hz), 4.33 (m, 1H), 4.24 (m, 1H), 3.31 (s, 3H), 1.49 (s, 3H), 1.30 (s, 3H); MS (EI) m/z 230 (M^+), 215 ($\text{M}^+ - \text{Me}$), 198, 155.

Methyl 5-(S)-Deoxy-5-vinyl-2,3-O-isopropylidene- β -Dribo-hexanofuranoside (**9**).

To a suspension of KH (418 mg, 10.4 mmol, prewashed with hexane) and tetrabutylammonium iodide (25 mg) in dry THF (10 mL) at 0°C was added dropwise a solution of alcohol **5** (1.2 g, 5.23 mmol) in THF (2 mL). The resulting red suspension was allowed to warm to 23°C , and the mixture was stirred for 1 h. The reaction was cooled to 0°C , and $\text{Bu}_3\text{SnCH}_2\text{I}$ (2.1 g, 6.8 mmol) in THF (2 mL) was added dropwise over a period of 2 min. The mixture was stirred at 0 – 23°C for 4 h. After this period, the reaction was quenched with a saturated NH_4Cl solution (10 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were dried over anhydrous Na_2SO_4 . Evaporation of the solvents provided the allyl stannylmethyl ether **7** as a colorless oil. This material was used for the next reaction without further purification. An analytical sample of **7** was obtained by flash chromatography over silica gel (5% ethyl acetate/hexane): ^1H NMR (400 MHz, CDCl_3) δ 5.71–5.74 (m, 2H), 4.98 (s, 1H), 4.65 (d, 1H, J 10.9 Hz), 4.61 (s, 2H), 3.85 (d, 2H, J 6.8 Hz), 3.69 (s, 2H), 3.32 (s, 3H), 1.49 (s, 3H), 1.35 (s, 3H), 1.20–1.62 (m, 12H), 0.85–0.94 (m, 15H).

To a stirred solution of above crude stannyl ether **7** in THF (10 mL) at -78°C , was added $^t\text{BuLi}$ (2.5 M in hexane, 3.2 mL, 8 mmol) dropwise by a syringe pump over a period of 1 h. The resulting mixture was stirred for 2 h at -78°C . After this period, the reaction was quenched with saturated NH_4Cl solution and the resulting mixture was warmed to 23°C . The layers were separated, and the aqueous layer was extracted with ethyl acetate (2×20 mL). The combined organic layers were dried over anhydrous Na_2SO_4 . Evaporation of the solvents gave a residue which was chromatographed over silica gel (25% ethyl acetate/hexane) to furnish an inseparable mixture (4.4:1 by ^1H NMR) of **9** and its epimer (1.07 g, 84% from **5**) as a colorless oil. Major isomer **9**: IR (neat) 3454, 2988, 2936, 1643, 1374, 1090 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 5.81 (m, 1H), 5.27 (dd, 1H, J 1.1, 10 Hz), 5.21 (dd, 1H, J 1.1, 18 Hz), 4.95 (s, 1H), 4.71 (dd, 1H, J 1.5, 6.2 Hz), 4.53 (d, 1H, J 6.2 Hz), 4.25 (dd, 1H, J 1.5, 8 Hz), 3.72 (m, 1H), 3.59 (m, 1H), 3.35 (s, 3H), 2.46 (m, 1H), 1.49 (s, 3H), 1.35 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ

Methyl 5-(S)-Deoxy-5-vinyl-2,3-O-isopropylidene- β -Dribo-hexanofuranoside (**9**).

(prepared from *Z*-alcohol **6**). The *Z*-alcohol **6** (672.8 mg, 2.92 mmol) was converted to *Z*-stannyl ether **8** as a colorless oil by following the procedure described for *E*-stannyl ether **7**. This material was used for the next reaction without further purification. An analytical sample of **8** was obtained by flash chromatography over silica gel (5% ethyl acetate/hexane). ^1H NMR for **8**: (400 MHz, CDCl_3) δ 5.61 (dd, 2H, J 4.2, 5 Hz), 4.96 (s, 1H), 4.92 (d, 1H, J 6.7 Hz), 4.62 (d, 1H, J 5.8 Hz), 4.56 (d, 1H, J 5.6 Hz), 4.0 (d, 1H, J 4.1 Hz), 3.72 (s, 2H), 3.31 (s, 3H), 1.43 (s, 3H), 1.29 (s, 3H), 1.24–1.63 (m, 12H), 0.86–0.98 (m, 15H). Wittig-Still

rearrangement of *Z*-stannyl ether **8** as described above (for stannyl ether **7**) provided the alcohol **9** (528 mg, 74% from **6**) exclusively as a colorless oil: $[R]^{23}_D -29$ (c 0.13, CHCl_3).

Methyl 5-(*R*)-*N*-[(Phenylmethoxy)carbonyl]-5,6-dideoxy-2,3-*O*-isopropylidene- β -D-ribo-hept-6-enfuranoside (11**).**

To a stirred solution of DMSO (0.18 mL, 2.54 mmol) in CH_2Cl_2 (5 mL) at -78°C was added dropwise oxalyl chloride (0.13 mL, 1.52 mmol). The resulting mixture was stirred for 2 min, and then a solution of alcohol **9** (248 mg, 1.02 mmol) in CH_2Cl_2 (2 mL) was added dropwise over a period of 1 min. The mixture was stirred at -78°C for an additional hour. The reaction was quenched with diisopropylethylamine (0.9 mL, 5.08 mmol), and the mixture was allowed to warm to 0°C for 30 min. Ethyl acetate (20 mL) was added to the reaction mixture, and the mixture was successively washed with a 1 M aqueous NaHSO_4 solution (20 mL) and pH 7 buffered solution (20 mL). The organic layer was dried over anhydrous Na_2SO_4 . Evaporation of the solvent gave the crude aldehyde, which was used for the next reaction immediately without further purification.

To a stirred solution of the above aldehyde in a mixture of *tert*-butyl alcohol (10 mL) and 2-methyl-2-butene (3 mL) at 23°C were added sodium chlorite (900 mg, 10 mmol) in water (1 mL) and NaH_2PO_4 (1.38 g, 10 mmol) in water (1 mL) sequentially. The resulting mixture was stirred at 23°C for 12 h. After this period, the reaction mixture was diluted with saturated NH_4Cl (10 mL) and the mixture was extracted with ethyl acetate (3×20 mL). The combined organic layers were dried over anhydrous Na_2SO_4 . Evaporation of the solvent provided the acid, which was directly used for the next reaction.

To a stirred solution of the above acid in acetone (4 mL) was added diisopropylethylamine (0.21 mL, 1.22 mmol) at 0°C . The resulting mixture was stirred for 10 min, and ClCO_2Me (0.1 mL, 1.32 mmol) was added dropwise over a period of 1 min. The reaction mixture was stirred at 0°C for 30 min, and then sodium azide (132 mg, 2.03 mmol) in water (2 mL) was added. The mixture was stirred for an additional 30 min at 0°C . After this period, CH_2Cl_2 (20 mL) and water (20 mL) were added and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 (2×20 mL). The combined organic layers were successively washed with 1 M NaHSO_4 (20 mL) and pH 7 buffer (20 mL) and dried over anhydrous Na_2SO_4 . Evaporation of the solvent under reduced pressure gave a colorless oil, which was dissolved in toluene (5 mL). The resulting mixture was heated at 100°C for 1 h. After this period, benzyl alcohol (0.21 mL, 2 mmol) was added and the resulting mixture was refluxed for 12 h. The reaction was cooled to 23°C , and the solvent was evaporated to give a yellow residue, which was chromatographed over silica gel (5% ethyl acetate/benzene) to furnish the Cbz derivative **11** (249 mg, 68% from **9**) as a white solid: mp $80\text{--}82^\circ\text{C}$; $[R]^{23}_D -11$ (c 0.18, CHCl_3); IR (neat) 3447, 2925, 1643, 1239, 1092 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.26–7.37 (m, 5H), 5.95 (m, 1H), 5.28 (d, 1H, J 11.4 Hz), 5.25 (dd, 1H, J 1, 4.7 Hz), 5.12 (dd, 1H, J 4.7, 17 Hz), 4.98 (s, 1H), 4.97 (br s, 1H), 4.74 (d, 1H, J 4.9 Hz), 4.59 (d, 1H, J 4.9 Hz), 4.29 (dd, 1H, J 5.6, 9.9 Hz), 3.98 (dd, 1H, J 1.1, 9.4 Hz), 3.34 (s, 3H), 1.46 (s, 3H), 1.29 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 25.0, 26.5, 55.0, 55.5, 66.9, 88.9, 109.7, 112.5, 116.8, 126.9, 127.5, 128.0, 128.1, 128.4, 135.3, 136.2, 155.9; MS (EI) m/z 363 (M^+),

331, 173, 146, 115, 91. Anal. Calcd for $C_{19}H_{25}NO_6$: C, 62.80; H, 6.93; N, 3.86. Found: C, 62.19; H, 7.14; N, 3.58.

Methyl (Methyl-5-*N*-[(phenylmethoxy)carbonyl]-5-deoxy-2,3-*O*-isopropylidene- β -D-allofuranoside)uronate (13).

To a stirred solution of **11** (36.5 mg, 0.1 mmol) in a mixture of CH_2Cl_2 (3 mL) and MeOH (1 mL) was added solid NaOH (20 mg), and a stream of ozonized oxygen was bubbled through this stirred mixture at $-78^\circ C$ until the color of the solution changed from yellow to blue. The reaction mixture was diluted with water and allowed to warm to $23^\circ C$. The layers were separated, and the aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were dried over anhydrous Na_2SO_4 . Evaporation of the solvent gave a residue, which was chromatographed over silica gel (15% ethyl acetate/hexane) to provide the methyl ester **13** (37.1 mg, 94%) as a white solid: mp $66-68^\circ C$; $[R]^{23}_D -14$ (c 2.81, $CHCl_3$); IR (neat) 3428, 1720, 1643, 1109 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 7.32–7.41 (m, 5H), 5.52 (d, 1H, J 8.3 Hz), 5.11 (brs, 2H), 4.96 (s, 1H), 4.93 (d, 1H, J 5.8 Hz), 4.57 (d, 1H, J 5.8 Hz), 4.49 (t, 1H, J 8 Hz), 4.33 (d, 1H, J 7.6 Hz), 3.75 (s, 3H), 3.32 (s, 3H), 1.51 (s, 3H), 1.30 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 25.0, 26.4, 52.5, 55.8, 56.4, 67.3, 81.3, 85.0, 87.6, 110.1, 112.6, 128.2, 128.5, 135.9, 155.7, 170.5; MS (EI) m/z 396 ($M^+ + H$), 395 (M^+), 173, 91. Anal. Calcd for $C_{19}H_{25}NO_8$: C, 57.71; H, 6.31; N, 3.54. Found: C, 57.53; H, 6.36; N, 3.75.

Methyl 5-Deoxy-5-[(phenylmethoxy)carbonyl]amino]- β -D-allofuranuronate-1,2,3-triacetate (14).

To a stirred solution of **13** (446.5 mg, 1.14 mmol) in MeOH (20 mL) was added Dowex 50W H^+ resin (600 mg), and the mixture was heated at $65^\circ C$ for 12 h. After this period, the resin was filtered off and the solvent was removed under reduced pressure. The residue was dissolved in pyridine (4 mL), Ac_2O (1 mL) was added, and the resulting mixture was stirred at $23^\circ C$ for 4 h. The reaction mixture was poured into ice, stirred for 30 min, and extracted with ethyl acetate (2×20 mL). The combined organic layers were successively washed with saturated aqueous $CuSO_4$ solution and brine and then dried over anhydrous Na_2SO_4 . Evaporation of the solvent gave a residue, which was dissolved in a mixture of CH_2Cl_2 (2 mL) and AcOH (2 mL). To this mixture at $0^\circ C$ was added Ac_2O (0.5 mL), followed by a drop of concentrated H_2SO_4 . The resulting mixture was stirred at $0^\circ C$ for 1 h and then at $23^\circ C$ for 2 h. The reaction mixture was poured onto ice and stirred for 30 min. The layers were separated, and the aqueous layer was extracted with CH_2Cl_2 (2×10 mL). The combined organic layer was successively washed with saturated $NaHCO_3$ and brine and dried over anhydrous Na_2SO_4 . Evaporation of the solvent provided a residue, which was chromatographed on silica gel (33% ethyl acetate/hexane) to furnish the triacetate **14** (373 mg, 70% from **13**) as a colorless oil: IR (neat) 3420, 1750, 1720 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 7.35 (s, 5H), 6.36 (d, 0.33H, J 8.9 Hz, R-anomer), 6.09 (s, 0.67H, β -anomer), 5.54 (m, 1H), 5.29 (d, 1H, J 4.7 Hz), 5.11 (s, 2H), 4.67 (dd, 1H, J 4.4, 8.7 Hz), 4.45 (dd, 1H, J 4.4, 7.6 Hz), 3.76 (s, 3H), 2.10 (s, 3H), 2.01 (s, 6H).

**Methyl 1,5-Dideoxy-1-(3,4-dihydro-5-methyl-2,4-dioxo1(2H)-pyrimidinyl)-5-
[[[(phenylmethoxy)carbonyl]amino] β -D-allofuranuronate-2,3-diacetate (15).**

To a stirred suspension of thymine (17 mg, 0.13 mmol) in hexamethyl disilazane (2 mL) was added trimethylchlorosilane (0.1 mL), and the resulting mixture was heated at 120 °C for 5 h. The mixture was cooled to 23 °C, and the solvent was removed under reduced pressure to give the crude bis-silylated thymine. The residue was dissolved in dichloroethane (2 mL), and a solution of triacetate **14** (20 mg, 0.043 mmol) in dichloroethane (1 mL) followed by TMSOTf (46 μ L, 0.126 mmol) was added at 23 °C. The resulting mixture was heated at 45 °C for 2 h, cooled to 23 °C, and quenched with saturated sodium bicarbonate solution. The layers were separated, and the aqueous layer was extracted with CH₂Cl₂ (3 $\times$ 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, and the solvents were evaporated under reduced pressure. The resulting residue was chromatographed on silica gel (40% ethyl acetate/benzene) to give the title nucleoside **15** (18.4 mg, 80%) as a pale yellow solid: mp 81–83 °C; [R]²³_D +17.1 (*c* 0.45, CHCl₃); IR (neat), 3380, 1750, 1715 cm⁻¹; ¹H NMR (200 MHz, CDCl₃) δ 8.15 (s, 1H), 7.35 (s, 5H), 7.05 (d, 1H, *J* 0.7 Hz), 5.93 (d, 1H, *J* 5.6 Hz), 5.79 (d, 1H, *J* 8.1 Hz), 5.51 (t, 1H, *J* 5.9 Hz), 5.26 (t, 1H, *J* 5.9 Hz), 5.14 (s, 2H), 4.81 (dd, 1H, *J* 3.7, 8.5 Hz), 4.39 (dd, 1H, *J* 3.7, 4.7 Hz), 3.81 (s, 3H), 2.09 (s, 6H), 1.87 (d, 3H, *J* 0.7 Hz); ¹³C NMR (100 MHz, CD₃COCD₃) δ 12.3, 20.3, 52.8, 56.1, 67.1, 70.6, 72.9, 81.8, 89.3, 111.6, 128.5, 128.6, 129.1, 137.3, 137.7, 151.4, 157.4, 164.0, 169.9, 170.0, 170.1; MS (EI) *m/z* 534 (M⁺+ H), 408, 91.

Acknowledgment.

Financial support of our work by the National Institutes of Health (GM 55600) is gratefully acknowledged.

References

- (1). (a) Nakai T; Mikami K *Org. React* 1994, 46, 105.
(b) Marshall JA *Comprehensive Organic Synthesis*; Trost BM, Fleming I, Eds.; Pergamon Press: Oxford, 1991; Vol. 3, p 975 and references therein.
- (2). (a) Still WC; Mitra A J. *Am. Chem. Soc* 1978, 100, 1927.
(b) Bruckner R; Priepeke H *Angew. Chem., Int. Ed. Engl.* 1988, 27, 278.
- (3). Ghosh AK; Liu WJ *Org. Chem.* 1996, 61, 6175.
- (4). (a) Isono K; Suzuki S *Heterocycles* 1979, 13, 333.
(b) Gooday GW *Abh. Dtsch. Akad. Wiss. DDR, Abt. Math. Naturwiss.; Tech, Issue 2N*; 1979; p 159.
(c) Mori M; Kakiki K; Misato T *Agric. Biol. Chem.* 1974, 38, 699.
- (5). For synthesis of polyoxin nucleotides, see: (a) Dehoux C; Fontaine E; Escudier J-M; Baltas M; Gorrichon L J. *Org. Chem* 1998, 63, 2601. [PubMed: 11672125]
(b) Dondoni A; Junquera F; Merchan FL; Merino P; Tejero T *Tetrahedron Lett.* 1994, 35, 9439.
(c) Garner P; Park JM J. *Org. Chem* 1990, 55, 3772.
(d) Barrett AGM; Lebold SA J. *Org. Chem* 1990, 55, 3853.
(e) Auberson Y; Vogel P *Tetrahedron* 1990, 46, 7019.
(f) Tabusa F; Yamada T; Suzuki K; Mukaiyama T *Chem. Lett* 1984, 405.
(g) Damodaran NP; Jones GH; Moffatt JG J. *Am. Chem. Soc* 1971, 93, 3812.
(h) Naka T; Hashizume T; Nishimura M *Tetrahedron Lett.* 1971, 95. [PubMed: 5545643]
- (6). For the total synthesis of polyoxins, see: (a) Akita H; Uchida K; Kato K *Heterocycles* 1998, 47, 157.
(b) Dondoni A; Junquera F; Merchan FL; Merino P; Tejero T J. *Org. Chem* 1997, 62, 5497.

- (c) Chida N; Koizumi K; Kitada Y; Yokoyama C; Ogawa S *J. Chem. Soc. Chem. Commun* 1994, 111.
 (d) Mukaiyama T; Suzuki K; Yamada T; Tabusa F *Tetrahedron* 1990, 46, 265.
 (e) Kazuhara H; Ohru H; Emoto S *Tetrahedron Lett.* 1973, 5055 and references therein.
 (7). (a) Ghosh AK; McKee SP; Sanders WM; Darke PL; Zugay JA; Emini EA; Schleif WA; Quintero JC; Huff JR; Anderson PS *Drug Des. Discovery* 1993, 10, 77.
 (b) Levene PA; Stiller ET *J. Biol. Chem.* 1934, 299.
 (8). (a) Still WC; Gennari C *Tetrahedron Lett.* 1983, 24, 4405.
 (b) Spada MR; Ubukata M; Isono K *Heterocycles* 1992, 34, 1147–6736 *J. Org. Chem*, Vol. 63, No. 19, 1998
 (9). (a) Priepke H; Brückner R *Chem. Ber* 1990, 123, 153.
 (b) Priepke H; Brückner R; Harms K *Chem. Ber* 1990, 123, 555.
 (c) Scheuplein SW; Kusche A; Brückner R; Harms K *Chem. Ber.* 1990, 123, 917 and references therein.
 (10). Houk KN; Paddon-Row MN; Rondan NG; Wu Y-D; Brown FK; Spellmeyer DC; Metz JT; Li Y; Loncharich RJ *Science* 1986, 231, 1108. [PubMed: 3945819]
 (11). (a) Ghosh AK; Hussain KA; Fidanze S *J. Org. Chem* 1997, 62, 6080.
 (b) Grunewald GL; Ye Q *J. Org. Chem* 1988, 53, 4021.
 (c) Ninomiya K; Iorio T; Yamada S *Tetrahedron* 1974, 30, 2151 and references therein. *J. Org. Chem*, Vol. 63, No. 19, 1998
 (12). The tentative α - β -unsaturated ester **12** was formed as a major product (50% yield).

Compound **11** was isolated in a small amount (<10% yield): ^1H NMR (200 MHz, CDCl_3) δ 7.35 (m, 5H), 6.16 (q, 1H, J 7.4 Hz), 5.23 (brs, 1H), 5.12 (d, 2H, J 4.7 Hz), 5.03 (s, 1H), 4.65 (dd, 1H, J 1.0, 6.0 Hz), 4.59 (d, 1H, J 6.0 Hz), 3.43 (s, 3H), 1.77 (d, 3H, J 7.4 Hz), 1.52 (s, 3H), 1.30 (s, 3H). 136, 135.4, 118.9, 112.6, 109.4, 87.3, 85.3, 82.2, 64.1, 63.2, 55.6; MS (EI) m/z 213 (M^+ - OMe), 197, 173, 59. Anal. Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_5$: C, 59.0; H, 8.25. Found: C, 58.90; H, 8.40. Minor isomer: ^1H NMR (400 MHz, CDCl_3) δ 5.15 (m, 1H), 5.29 (m, 2H), 4.99 (s, 1H), 4.68 (dd, 1H, J 1, 5.9 Hz), 4.57 (d, 1H, J 5.9 Hz), 4.14 (d, 1H, J 11.2 Hz), 3.81 (m, 1H), 3.66 (m, 1H), 3.40 (s, 3H), 2.46 (m, 1H), 1.48 (s, 3H), 1.31 (s, 3H).

- (13). Marshall JA; Garofalo AW; Sedrani R *Synlett* 1992, 643.
 (14). Vorbrüggen H; Krolkiewicz K; Bennua B *Chem. Ber.* 1981, 114, 1234.

Figure 1.

Scheme 1.

a^a Key: (a) DIBAL, CH₂Cl₂, -78 °C, 2–3 h (92–95%); (b) KH, Bu₄N⁺I⁻, Bu₃SnCH₂I, THF, 23 °C, 4 h; (c) ⁿBuLi, THF, -78 °C, 2 h (74–84%).

Scheme 2.

a ^a Key: (a) Swern oxidation; (b) NaClO₂, Me₂CdCHMe, ^tBuOH, 23 °C, 12 h; (c) MeOCOC₂H₅, ^tPr₂NEt, NaN₃ then PhCH₂OH, PhMe, 100 °C, 12 h (68% from 9); (d) O₃, NaOH, CH₂Cl₂, MeOH, -78 to 23 °C, 2 h (94%); (e) Dowex 50W, MeOH, 65 °C, 12 h; (f) Ac₂O, Py, 23 °C, 4 h; (g) Ac₂O, AcOH, CH₂Cl₂, cat. H₂SO₄, 23 °C, 2 h (70% from 13); (h) Thymine-bis-TMS, Cl (CH₂)₂Cl, TMSOTf, 45 °C, 2 h (80%).