RESEARCH ARTICLE

Radiographic features of lingual mandibular bone depression using dental cone beam computed tomography

Liu Liu, Byung Cheol Kang, Suk Ja Yoon, Jae Seo Lee and Sel Ae Hwang

Department of Oral and Maxillofacial Radiology, School of Dentistry, Dental Science Research Institute, Chonnam National University, Gwangju, South Korea

Objectives: To describe accurately the radiographic characteristics of lingual mandibular bone depression (LMBD) using CBCT images.

Methods: This study included 30 cases of LMBD with CBCT images from 29 patients. Age, sex, location, shape, size, and the relationships of the LMBDs to adjacent anatomical structures such as the mandibular inferior cortex and mandibular canal were recorded. The correlation between age and size was evaluated using Pearson's correlation coefficient.

Results: The average age of the patients was 57 years. LMBDs occurred primarily in males (93%), occurrence in the right ($n = 14$) and left ($n = 16$) mandibles was approximately equal. The shapes were ovoid, round, peanut, triangular, or rectangular on panoramic radiographs. The occurrences in the molar, distal to the molar, and ramus regions were 63.3, 33.3, and 3.3% respectively. The mean size on CBCT was 15.4 mm in length, 11.2 mm in height, and 6.4 mm in depth. The relationship to the mandibular canal was categorized as separated (48.2%), contacted (31%), or go-through (20.7%). The relationship to the mandibular inferior cortex was classified as separated (20.7%), contacted (27.6%), or resorbed (51.7%). The Pearson's correlation coefficient showed a positive correlation between age and depth ($r = 0.38$; $p = 0.038$).

Conclusions: The typical features of a LMBD were ovoid, round or multilobated radiolucent shape, well-defined sclerotic border, and position in the posterior area of the mandible between the mandibular canal and inferior mandibular cortex. The buccolingual depth of LMBD was deeper with increasing age.

Dentomaxillofacial Radiology (2018) 47, 20170383. doi: [10.1259/dmfr.20170383](https://doi.org/10.1259/dmfr.20170383)

Cite this article as: Liu L, Kang BC, Yoon SJ, Lee JS, Hwang SA. Radiographic features of lingual mandibular bone depression using dental cone beam computed tomography. *Dentomaxillofac Radiol* 2018; 47: 20170383.

Keywords: Cone beam computed tomography; salivary gland; mandible; ultrasonography

Introduction

The lingual mandibular bone depression (LMBD) was first reported by Stafne in 1942 to describe 35 asymptomatic bone cavities in the angle of the mandible.¹ Since then, there have been many articles about the condition. There are several synonyms for the condition, including Stafne bone cyst or cavity,² Stafne's idiopathic bone cavity or cyst,^{3,4} static bone cavity,⁵ lingual mandibular salivary gland depression,⁶ lingual cortical mandibular bone depression,⁷ and combinations of the above. The

most common location is below the mandibular canal, within the submandibular salivary gland fossa and often close to the inferior border of the mandible, but it can also be found in the anterior part of the mandible. Due to their asymptomatic nature, most depressions were found incidentally during radiographic examination. The typical radiographic feature of an LMBD is a well-demarcated round or ovoid radiolucency with a well-defined sclerotic border. Most cases are unilateral and unilocular depressions. Bilateral, bilobate, multilobate, simultaneous unilateral anterior and posterior defects have also been reported.⁸⁻¹¹ Evidence of a

Correspondence to: Dr Byung Cheol Kang, E-mail: bckang@jnu.ac.kr

Received 16 October 2017; revised 19 March 2018; accepted 23 March 2018

concavity with a lingual opening on CBCT images could exclude most of the differential diagnosis of other types of bone lesions for which surgical treatment is needed.¹² CT studies showed that the contents of LMBD were not always aberrations of the submandibular gland, but were fatty tissues, blood vessels and soft tissue.^{2,5}

Previous studies have measured the size of the LMBD on panoramic radiograph,¹³ which are inherently unreliable for horizontal measurement.¹⁴ The typical feature of LMBD is the mandible lingual surface opening which could not be interpreted on two-dimensional image.¹⁵ CBCT-enabled two-dimensional assessment of craniofacial bone structures with low radiation dose and high resolution image. So, it can be used for differential diagnosis of LMBD and investigating the relationship between depression and adjacent anatomic structures.

The following survey is based on 30 unpublished cases from South Korea. The aim of this study was to use CBCT to measure the exact size of the LMBD and determine the relationship between LMBDs and adjacent anatomic structures.

Methods and materials

We retrieved 120 LMBD cases (including six bilateral cases) from Picture Archiving and Communication System (PACS) stored from December 2008 to November 2016 at Chonnam University Dental Hospital (Gwangju, South Korea). The cases included 98 males and 16 females, aged 26–87 years. 90 cases with panoramic images only were excluded. 30 cases with CBCT and panoramic images were selected for this study. Inclusion criteria were no clinical symptoms, well-defined radiolucency of the mandible, and concavity on the lingual surface of the mandible on CBCT images.

All of the radiographs including the panoramic (OP100, Planmeca, Helsinki, Finland; Kodak 8000C, Carestream, Rochester, NY) and CBCT images (Pax-i3D, VATECH, Seoul, Korea; CS 9300, Carestream, Atlanta, GA; CB MercuRay, Hitachi, Tokyo, Japan) were evaluated by PiViewSTAR (Infinite Healthcare, Seoul, Korea). Ultrasonographic examination (LOGIQ 7, GE, Milwaukee, WI) was performed to evaluate content of a LMBD with buccal cortical perforation during the patient's enlarged submandibular salivary gland examination.

The age, sex, location, size and relationship to adjacent anatomical structures were recorded. The locations of the centers of the depressions were classified as incisor–premolar, molar, distal to molar, and ramus (Figure 1).

The size of the depressions was measured on CBCT images. All of the measurements were made by two experienced oral and maxillofacial radiologists blinded to each other on the monitor (LG IPS FULLHD 34 inch monitor with 1920 × 1080 resolution). The depth was measured as the maximum distance from the lingual surface to the bottom of the depression (facial cortical bone) in axial section image; the length was the maximum

Figure 1 The four possible locations of LMBD. The line AB is the bisecting line formed by the intersection of the mandibular and ramal planes. LMBD, lingual mandibular bone depression.

anteroposterior distance, and the height was the maximum superoinferior distance in the cross section image.

The relationship between the LMBD and mandibular canal was examined on CBCT and the relation was divided into three types excluding one ramus case.¹⁶

Separated: mandibular canal is separated from the LMBD (Figure 2a).

Contacted: mandibular canal is in contact with the LMBD (Figure 2b).

Go through: mandibular canal goes through the LMBD (Figure 2c).

The relationship between the LMBD and mandibular inferior cortex was divided into three types excluding one ramus case.

Separated: the inferior border of depression separated from the mandibular inferior cortex (Figure 3a).

Contacted: the inferior border of depression contacted with the mandibular cortex (Figure 3b).

Involved: the inferior border of depression involved the mandibular inferior cortex (Figure 3c).

The relationship between depression and buccal cortical plate, was classified as suggested by Arijj *et al.*²

Separated: the bottom of the depression did not reach the buccal cortical plate (Figure 3a).

Contacted: the bottom of the depression reached the buccal cortical plate, but there was no expansion or distortion of the plate (Figure 3c).

Figure 2 The relationships between the depression and mandibular canal. (a) Separated: the mandibular canal is separated from the depression. (b) Contacted: the mandibular canal is in contact with the depression. (c) Resorbed: the mandibular canal goes through the depression.

Figure 3 The relationships between the depression and mandibular inferior cortex. (a) The bottom of the depression is separated from the mandibular inferior cortex. (b) The bottom of the depression is in contact with the mandibular inferior cortex, which is intact. (c) The bottom of the depression is involved the mandibular inferior cortex which is resorbed.

Expanded: the depression was characterized by a buccal expansion of the cortical plate (Figure 2a).

Statistical analysis was performed with SPSS Statistics for Windows, v. 17.0 (SPSS Inc, Chicago, IL). A non-parametric Kruskal–Wallis test was used to compare inter-reliability of size measurements between examiners. Cronbach’s α test was used to test intraexaminers agreement. The correlation between age and size was evaluated using Pearson’s correlation coefficient.

Results

A total of 30 cases were diagnosed in 29 patients (one patient presented two depressions, Figure 4a,b). All patients were asymptomatic. The average age was 57.4 years (range, 33–77). The male to female (M/F) ratio was 14:1, and the right to left mandibular occurrence ratio was 1:1.1. 29 cases were found in the posterior region of the mandible, and 1 was found at the ramus. There was no depression in incisor-premolar area. The occurrences at the “molar”, “distal to molar” and “ramus” were 63.3, 33.3, and 3.3% (19, 10, and 1 case/s) respectively. The mean size of the depression was 15.4 mm in length (range, 6.4–33.3 mm), 11.2 mm in height (4.1–19.8 mm), and 6.4 mm in depth (2.3–13.8 mm).

Figure 4 (a) Two adjacent LMBDs were detected on a panoramic radiograph. (b) CBCT showed the first depression (short arrow) in contact with the posterior second one (long arrow). LMBD, lingual mandibular bone depression.

Table 1 Relationships with adjacent anatomical structures

	Type	Type	Type
Relationship with the mandibular canal ^{a,16}	Separated	Contacted	Go through
Number	14	9	6
Incidence	48.30%	31%	20.70%
Relationship with the mandibular inferior cortex ^a	Separated	Contacted	Involved
Number	6	8	15
Incidence	20.70%	27.60%	51.70%
Relationship with the buccal cortical plate ²	Separated	Contacted	Expanded
Number	7	15	8
Incidence	23.30%	50%	26.70%

^aExcluding one ramus case.

Most of the LMBDs were separated from the mandibular canal (48.2%, $n = 14$), followed by contacted with the canal (31%, $n = 9$), and go through the canal (20.7%, $n = 6$), excluding the ramus case.

Most of the LMBDs involved the mandibular inferior cortex (51.7%, $n = 15$), followed by contacted with the cortex (27.6%, $n = 8$), and separated from the cortex (20.7%, $n = 6$), excluding the ramus case.

Most bottoms were contacted with the buccal cortical plate (50%, $n = 15$), followed by expanded the buccal cortical plate (26.7%, $n = 8$), and separated from the buccal cortical plate (23.3%, $n = 7$) (Table 1).

One case presented mandibular inferior border destruction and mandibular buccal plate expansion and perforation on CBCT (Figure 5a,b). We got a non-interrupted hyperechoic outer convex images on sonography (arrow, Figure 5c), meaning the expanded mandibular facial cortical bone was not perforated.

There was no significant difference between examiners for size measurement ($p > 0.05$). Cronbach’s α values showed excellent intraobserver agreement ($\alpha > 0.9$) The Pearson’s correlation coefficient showed a weak positive correlation between LMBD depth and age ($r = 0.38$, $p = 0.038$) (Figure 6). There was no correlation among height, length and age.

Discussion

The incidence of LMBD is 0.009–0.7% on radiographs.^{17,18} Bone study revealed a higher incidence rate of 1.7–2.4% (4.5% in male, 0.4% in female).^{19,20} The depressions could be detected on panoramic radiographs only when the depth is greater than 1.5 mm, therefore, the actual incidence of LMBD for the living populations might be underestimated.²⁰ In the present study, all depressions’ depths were more than 2 mm.

The M/F ratio in this study was 14:1, indicating a strong male preponderance. The ratio was slightly higher than those reported in previous studies (M/F ranges from 6:1–1:1).^{7,19} Expanding our cases to those

Figure 5 (a) LMBD on a panoramic radiograph. (b) Thinning and perforation of the expansive buccal cortical bone on cross-sectional CBCT image. (c) The arrow represent hyperechoic convex of buccal cortical expansion of the LMBD representing advanced thinning on a sonographic image. LMBD, lingual mandibular bone depression.

with and without CBCT images, M/F ratio of the first retrieved 120 cases was 6:1.

In the present study, all of the LMBDs were on the lingual aspect of the mandible. Except 1 case on the ramus, 29 LMBDs were located in the submandibular fossa, between the mandibular canal and the mandibular inferior cortex.

The average size of the depressions in our study was 15.4 mm in length, 11.2 mm in height, and 6.4 mm in depth. The present study showed that the buccolingual depth of LMBD was increased with aging (Figure 6). The length and width had no correlation with age. In contrast to our result, Mourao et al reported that the sizes of LMBD over age 55 were smaller than that of people under age 55. They measured only the antero-posterior and vertical dimensions of the 20 cases of LMBD utilizing panoramic radiographs.⁷

The pathogenesis of LMBD is still controversial and not fully understood. Several hypotheses have been proposed. Stafne suggested that the depressions were congenital and caused by a failure of fusion in the areas formerly occupied by Meckel's cartilage.¹ Campos et al reported that the LMBD should not be seen as a

salivary gland-related bone defect, because there was no contact between the major salivary glands and the bone surface.²¹ They preferred a congenital origin, and theorized that the depressions were the result of a focal failure during mandibular intramembranous ossification. However, the theory could not explain that no occurrence has been documented in children younger than 11 years old.²²

The general consensus is the “glandular pressure theory”. The well defined depression was thought to be associated with the pressure from major salivary glands, adjacent to the lingual surface of the mandible. Posterior depressions are associated with the submandibular glands, anterior depressions are associated with the sublingual gland, and depressions on the ascending ramus thought to be associated with the parotid gland.¹⁴ CT examinations, sialography, and surgical explorations had showed salivary gland tissue in the LMBD.^{23,24}

A LMBD is usually encountered incidentally on routine plain radiographs. The appearance and location of LMBD are characteristic and easily identified. LMBD can be readily differentiated from odontogenic lesions such as cysts because the epicenter of odontogenic lesions is located above the inferior alveolar canal.^{3,14,25}

The depression may be difficult to detect on panoramic radiographs because the degree of resorption is not sufficient to produce radiographic changes. However, the depression can be detected easily on CBCT images (Figure 4a,b). The CBCT examination is beneficial for confirming lingual opening and depth measurement of LMBD. CBCT has more advantages compared with Multi-Slice Computer Tomograph due to low exposure dose of radiation, effectively used in dental clinics and the high resolution of images.²⁶ However, dental CBCT provides limited quantitative CT values for each pixel on sliced images for the differentiation of air, water and soft tissues, so CBCT could not determinate the contents of depressions.²⁷ Therefore, sometimes additional diagnostic procedures such as medical CT, MRI and sialography techniques are needed to confirm.^{5,28,29}

In conclusion, the present study showed the radiological features of LMBDs, and revealed a relationship

Figure 6 Correlation between LMBD depth and age. ($r = 0.38$, $p = 0.038$). LMBD, lingual mandibular bone depression.

between LMBDs and adjacent anatomy based on 30 cases from South Korea. The deeper LMBD occurred in older patients. The buccolingual depth of LMBD was deeper with increasing age. CBCT is an effective and suitable approach for investigating LMBDs.

References

1. Stafne EC. Bone cavities situated near the angle of the mandible. *J Am Dent Assoc* 1942; **29**: 1969–72. doi: <https://doi.org/10.14219/jada.archive.1942.0315>
2. Arijji E, Fujiwara N, Tabata O, Nakayama E, Kanda S, Shiratsuchi Y, et al. Stafne's bone cavity. Classification based on outline and content determined by computed tomography. *Oral Surg Oral Med Oral Pathol* 1993; **76**: 375–80. doi: <https://doi.org/10.1038/sj.dmf.4600294>
3. Parvizi F, Rout PG. An ossifying fibroma presenting as stafne's idiopathic bone cavity. *Dentomaxillofac Radiol* 1997; **26**: 361–3. doi: <https://doi.org/10.1038/sj.dmf.4600294>
4. Ogunsalu C, Pillai K, Barclay S. Radiological assessment of type II Stafne idiopathic bone cyst in a patient undergoing implant therapy: a case report. *West Indian Med J* 2006; **55**: 447–50. doi: <https://doi.org/10.1590/S0043-31442006000600016>
5. Minowa K, Inoue N, Sawamura T, Matsuda A, Totsuka Y, Nakamura M. Evaluation of static bone cavities with CT and MRI. *Dentomaxillofac Radiol* 2003; **32**: 2–7. doi: <https://doi.org/10.1259/dmfr/92201340>
6. Cataldo E, Kabani S. A clinico-pathological presentation. Lingual mandibular salivary gland depression. *J Mass Dent Soc* 1985; **34**: 8.
7. Mourão CF, Miranda AM, Santos EJ, Pires FR. Lingual cortical mandibular bone depression: frequency and clinical-radiological features in a Brazilian population. *Braz Dent J* 2013; **24**: 157–62. doi: <https://doi.org/10.1590/0103-6440201302091>
8. Kim H, Seok JY, Lee S, An J, Kim NR, Chung DH, et al. Bilateral stafne bone cavity in the anterior mandible with heterotopic salivary gland tissue: a case report. *Korean J Pathol* 2014; **48**: 248–9. doi: <https://doi.org/10.4132/KoreanJPathol.2014.48.3.248>
9. Ozaki H, Ishikawa S, Kitabatake K, Yusa K, Tachibana H, Iino M. A case of simultaneous unilateral anterior and posterior stafne bone defects. *Case Rep Dent* 2015; **2015**: 1–5. doi: <https://doi.org/10.1155/2015/983956>
10. Boffano P, Gallesio C, Daniele D, Rocchia F. An unusual trilobate stafne bone cavity. *Surg Radiol Anat* 2013; **35**: 351–3. doi: <https://doi.org/10.1007/s00276-012-1043-7>
11. Aguiar LB, Neves FS, Bastos LC, Crusoe-Rebello I, Ambrosano GM, Campos PS. Multiple stafne bone defects: a rare entity. *ISRN Dent* 2011; **2011**: 1–3. doi: <https://doi.org/10.5402/2011/792145>
12. Monteiro LS, Câmara MID, Tadeu F, Salazar F, Pacheco JJ. Posterior lingual bone depression diagnosis using 3D-computed tomography. *Revista Portuguesa de Estomatologia, Medicina Dentária e Cirurgia Maxilofacial* 2012; **53**: 170–4. doi: <https://doi.org/10.1016/j.rpemd.2012.05.007>
13. Tsui SH, Chan FF. Lingual mandibular bone defect. Case report and review of the literature. *Aust Dent J* 1994; **39**: 368–71. doi: <https://doi.org/10.1111/j.1834-7819.1994.tb03108.x>
14. White SC, Pharoah MJ. *Oral radiology: principles and interpretation*: Elsevier Health Sciences; 2014. pp. 168.
15. Mann RW, Tuamsuk P. Stafne's defects in a sample of adult Thai Mandibles in the Khon Kaen University Osteology Collection, Thailand. *Int J Osteoarchaeol* 2016; **26**: 906–11. doi: <https://doi.org/10.1002/oa.2508>
16. Adisen MZ, Yilmaz S, Misirlioglu M, Atil F. Evaluation of volumetric measurements on CBCT images using stafne bone cavities as an example. *Med Oral Patol Oral Cir Bucal* 2015; **20**: e580–e586. doi: <https://doi.org/10.4317/medoral.20633>
17. Assaf AT, Solaty M, Zrnc TA, Fuhrmann AW, Scheuer H, Heiland M, et al. Prevalence of stafne's bone cavity-retrospective analysis of 14,005 panoramic views. *In Vivo* 2014; **28**: 1159–64.
18. Schneider T, Filo K, Locher MC, Gander T, Metzler P, Grätz KW, et al. Stafne bone cavities: systematic algorithm for diagnosis derived from retrospective data over a 5-year period. *Br J Oral Maxillofac Surg* 2014; **52**: 369–74. doi: <https://doi.org/10.1016/j.bjoms.2014.01.017>
19. Shields ED, Mann RW. Salivary glands and human selection: a hypothesis. *J Craniofac Genet Dev Biol* 1996; **16**: 126–36.
20. Wasterlain SN, Silva AM. Study of stafne's defects in Late Neolithic, Late Roman, Medieval and Modern skeletal samples from Portugal. *Int J Osteoarchaeol* 2012; **22**: 423–34. doi: <https://doi.org/10.1002/oa.1216>
21. Campos PS, Panella J, Crusoe-Rebello IM, Azevedo RA, Pena N, Cunha T. Mandibular ramus-related stafne's bone cavity. *Dentomaxillofac Radiol* 2004; **33**: 63–6. doi: <https://doi.org/10.1259/dmfr/39682286>
22. Friedrich RE, Scheuer HA, Gröbe A. Anterior lingual mandibular bone depression in an 11-year-old child. *In Vivo* 2012; **26**: 1103–7.
23. Quesada-Gómez C, Valmaseda-Castellón E, Berini-Aytés L, Gay-Escoda C. Stafne bone cavity: a retrospective study of 11 cases. *Med Oral Patol Oral Cir Bucal* 2006; **11**: E277–80.
24. Sisman Y, Miloglu O, Sekerci AE, Yilmaz AB, Demirtas O, Tokmak TT. Radiographic evaluation on prevalence of Stafne bone defect: a study from two centres in Turkey. *Dentomaxillofac Radiol* 2012; **41**: 152–8. doi: <https://doi.org/10.1259/dmfr/10586700>
25. Prapanpoch S, Langlais RP. Lingual cortical defect of the mandible: an unusual presentation and tomographic diagnosis. *Dentomaxillofac Radiol* 1994; **23**: 234–7. doi: <https://doi.org/10.1259/dmfr.23.4.7835531>
26. Sisman Y, Etöz OA, Mavili E, Sahman H, Tarim Ertas E. Anterior stafne bone defect mimicking a residual cyst: a case report. *Dentomaxillofac Radiol* 2010; **39**: 124–6. doi: <https://doi.org/10.1259/dmfr/49320253>
27. Yamashina A, Tanimoto K, Sutthiprapaporn P, Hayakawa Y. The reliability of computed tomography (CT) values and dimensional measurements of the oropharyngeal region using cone beam CT: comparison with multidetector CT. *Dentomaxillofac Radiol* 2008; **37**: 245–51. doi: <https://doi.org/10.1259/dmfr/45926904>
28. Shimizu M, Osa N, Okamura K, Yoshiura K. CT analysis of the stafne's bone defects of the mandible. *Dentomaxillofac Radiol* 2006; **35**: 95–102. doi: <https://doi.org/10.1259/dmfr/71115878>
29. Li B, Long X, Cheng Y, Wang S. Cone beam CT sialography of stafne bone cavity. *Dentomaxillofac Radiol* 2011; **40**: 519–23. doi: <https://doi.org/10.1259/dmfr/84472023>

Funding

This study was approved by the Institutional Review Board of the Chonnam National University Dental Hospital (CNUHD-2017-002).