Published in final edited form as:

Chemosphere. 2014 November ; 114: 219–225. doi:10.1016/j.chemosphere.2014.04.066.

Adjusting serum concentrations of organochlorine compounds by lipids and symptoms: A causal framework for the association with K-ras mutations in pancreatic cancer

Tomás López^{a,b}, José A. Pumarega^{a,c}, Anna Z. Pollack^d, Duk-Hee Lee^e, Lorenzo Richiardi^f, David R. Jacobs Jr.^g, Enrique F. Schisterman^d, and Miquel Porta^{a,b,c,*}

^aHospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

^bFacultat de Medicina, Universitat Autònoma de Barcelona, Catalonia, Spain

^cCIBER en Epidemiología y Salud Pública (CIBERESP), Spain

^dEpidemiology Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, Rockville, MD, USA

^eDepartment of Preventive Medicine and Health Promotion Research Center, School of Medicine, Kyungpook National University, Daegu, Republic of Korea

^fCancer Epidemiology Unit, Department of Medical Sciences, University of Turin and CPO-Piemonte, Torino, Italy

^gDivision of Epidemiology, School of Public Health, University of Minnesota, Minneapolis, MN, USA

Abstract

In clinically aggressive diseases, patients experience pathophysiological changes that often alter concentrations of lipids and environmental lipophilic factors; such changes are related to disease signs and symptoms. The aim of the study was to compare the effects of correcting for total serum lipids (TSL) and other clinical factors on the odds of mutations in the K-ras oncogene by organochlorine compounds (OCs), in logistic models, in 103 patients with exocrine pancreatic cancer (EPC) using a causal directed acyclic graph (DAG) framework. Results and likelihood of bias were discussed in the light of possible causal scenarios. The odds of K-ras mutated EPC was associated with some TSL-corrected OCs, including *p,p'*-DDT (*p*-value: 0.008) and polychlorinated biphenyl 138 (*p*-trend: 0.024). When OCs were not corrected by TSL, the OR of a K-ras mutation was significant for *p,p'*-DDT (*p*-trend: 0.035). Additionally adjusting for cholestatic syndrome increased the ORs of TSL-corrected OCs. When models were adjusted by the interval from first symptom to blood extraction (ISE), the ORs increased for both TSL-corrected and uncorrected OCs. Models with TSL-corrected OCs and adjusted for cholestatic syndrome or ISE yielded the highest ORs. We show that DAGs clarify the covariates necessary to

*Corresponding author. Address: IMIM PRBB UAB, Carrer del Dr. Aiguader 88, E-08003 Barcelona, Catalonia, Spain. Tel.: +34 93 316 0700/0781; fax: +34 93 316 0410., mporta@imim.es (M. Porta).

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.chemosphere.2014.04.066>.

minimize bias, and demonstrate scenarios under which adjustment for TSL-corrected OCs and failure to adjust for symptoms or ISE may induce bias. Models with TSL-uncorrected OCs may be biased too, and adjusting by symptoms or ISE may not control such biases. Our findings may have implications as well for studying environmental causes of other clinically aggressive diseases.

Keywords

Causal directed acyclic graph (DAG); Pancreatic neoplasm; Organochlorine compounds; Lipids; K-*ras* gene; Disease progression

1. Introduction

Organochlorine compounds (OCs) are highly lipophilic and resistant to degradation. Human contamination is common and exposure occurs primarily through consumption of fatty foods (Porta et al., 2008a; Reed et al., 2007). A substantial number of studies have shown associations between internal contamination by OCs and intermediate disorders (e.g., immunological) and severe diseases, including cancer (Centers for Disease Control and Prevention, 2009; Leijds et al., 2009; Porta, 2012), but it has often been difficult to rule out “disease progression bias” or “reverse causality” (Porta, 2008).

Indeed, there are still many unresolved methodological questions related to the toxicokinetics of OCs in humans, particularly when they are sick. Correction of blood concentration of OCs by total serum lipids (TSL) is frequently used in studies, on the premise that lipophilic substances are in a state of equilibrium across body compartments (blood, fat, organs) (Porta et al., 2009a). Thus, the lipid correction of OC blood levels improved precision, and it became common to express OCs on a lipid-basis. However, such conditions of equilibrium may not hold in overt disease, or even in the subclinical phases of disease, long before diagnosis (Porta, 2001; Schisterman et al., 2005; Wolff et al., 2007). It is also plausible that some OCs affect lipid concentrations (e.g., by inducing lipid synthesis), thus, even regression adjustment by TSL might entail mis-adjustment by a variable in the causal pathway (Goncharov et al., 2008; Lee et al., 2011; Porta et al., 2009c).

Clinical, metabolic and toxicological factors may influence the accumulation of OCs and other lipophilic compounds in fatty tissues, their mobilization, the blood/fat ratio, and the blood/organ tissue ratio. Both subtle and abrupt changes in these processes are common in clinically aggressive diseases as cancer (Kritchevsky et al., 1991; Porta, 2001; Wolff et al., 2007). Little is known about the kinetics of lipophilic environmental pollutants during the prediagnostic phases of cancer and other disease periods (Wolff et al., 2000). Statistical approaches to control lipid mobilization and the other pathophysiologic changes that commonly take place during cancer progression are thus necessary to prevent biased estimates of associations between OCs and risk of cancer (Baris et al., 2000; Lee et al., 2011; Spinelli et al., 2007).

Disease signs and symptoms reflect underlying pathophysiological processes (Porta et al., 2003, 2009c, 2012). In this paper we will consider where such symptoms fit on the causal spectrum in determining whether characteristics of the disease influence associations

between the occurrence of disease-related genetic alterations and environmental exposures, keeping in mind that the associations may also be reflective of disease-related lipid alterations.

We wish to estimate the possible causal effect of OC levels on *Kras* mutations in exocrine pancreatic cancer (EPC) without bias. To this end, the use of causal directed acyclic graphs (DAGs) can be helpful. DAGs can clarify issues relating to adjustment and bias (Greenland et al., 1999; Hernán et al., 2002). Using DAGs, Schisterman et al. (2005) showed that lipid adjustment may lead to bias, depending on the underlying causal relationship in question. As many other types of cancer, EPC itself affects lipid levels and, therefore, cannot be considered in the context of this previous work. In particular, patients with a highly aggressive disease as EPC experience pathophysiological changes (e.g., part of the constitutional or the cholestatic syndrome) that profoundly alter TSL levels. TSL thus vary in different stages of the disease (Porta et al., 2008b). It is important to note that in EPC *K-ras* mutations do not affect lipid metabolism, BMI or the prognosis of the disease. Therefore, utilizing a causal understanding of the relationships between *K-ras* mutations, OC exposure levels, lipid correction, age, sex, and cancer symptoms, we sought to clarify understanding of appropriate adjustment for confounding factors to minimize bias in a clinical series of EPC patients.

The aim of the present study was to compare alterations to the odds of having a *K-ras* mutated EPC vs. a *K-ras* wild-type EPC, given a certain concentration of an OC, from additional adjustment for TSL, disease symptoms and other clinical factors in patients with EPC.

2. Material and methods

Methods and strategies of analysis of the PANKRAS II study have been previously described in detail (Crous-Bou, 2009; Porta et al., 2008b). Briefly, subject recruitment took place between 1992 and 1995 at five general hospitals in the Mediterranean part of Spain, where 185 incident cases of EPC were prospectively identified. The present report is based on 103 EPC patients with known *K-ras* status and with analysis of OCs.

2.1. Clinicopathological information and personal interviews

Patients were interviewed face-to-face by trained monitors during hospital stay, close to the time of diagnosis. Over 40% of the blood draws took place in the same day of the interview, and over 94% of the interviews took place in the 2 weeks before or after the blood extraction. Interviews included questions about occupation, lifestyle, and past clinical history (Crous-Bou, 2009; Porta et al., 2009b). On the basis of previous research (Porta et al., 2003), detailed information on signs and symptoms was obtained from two sources: medical records and patient interviews. The cholestatic syndrome involved jaundice, hypocholia and choluria, while the constitutional syndrome comprised asthenia, anorexia and weight loss (Porta et al., 2008b).

Blood was drawn in the early morning and with patients fasting (Porta et al., 2007). Each subject had one measure of serum concentrations of organochlorines. The median time between the first symptom of cancer and blood extraction (ISE) was 73 days.

2.2. Laboratory analysis

K-ras mutations were analyzed by nested polymerase chain reaction (PCR) and direct DNA sequencing as described previously (Crous-Bou, 2009; Gasull et al., 2010). It should be noted that the occurrence of *K-ras* mutations is not affected by lipid mobilization, EPC symptoms, or any clinical markers of disease emergence or progression.

Details of laboratory protocols of OCs have also been described elsewhere (Gasull et al., 2013; Porta et al., 2009b). Briefly, gas chromatography analyses were performed and selected samples were analyzed by negative-ion chemical ionization gas chromatography-mass spectrometry. Statistical analyses were limited to compounds that were detected above the detection limit >85% of subjects. When a sample had an OC concentration below the detection threshold, it was assigned the mid-value of this limit, when an OC was detected but under the quantification threshold, the midvalue between detection and quantification limits was assigned.

Total cholesterol and triglycerides were determined enzymatically, using serum obtained at the same time as the serum used for the OC analyses. TSL were calculated by the traditional or standard formula 2 proposed by Phillips et al. (1989) which uses total cholesterol and triglycerides. Among cases, mean (standard deviation) serum concentrations of total cholesterol, triglycerides and TSL were, respectively, 209 (102), 178 (94) and 714 (293) mg dL⁻¹. The main variables in this study were TSL-corrected OC, organochlorine concentrations individually corrected by TSL and expressed in ng g⁻¹ lipid, the lipid correction is performed by simply dividing the serum OC value by the serum TL value and TSL-uncorrected OC, organochlorine concentrations uncorrected by TSL and expressed in ng mL⁻¹ (i.e. TSL-uncorrected *p,p'*-DDE).

2.3. Statistical analyses

We use multivariate-adjusted odds ratios (OR) and their corresponding 95% confidence intervals (CI), calculated by unconditional logistic regression. We categorized the OCs (corrected and uncorrected by TSL) in tertiles, the cutoff points of the tertiles are shown in Table A.1. Categorical ordinal variables were analyzed for a linear dose-response relation through the multivariate analogue of Mantel's extension test, when a linear trend was not apparent, Wald's test was used. Age and sex were assessed in all models as potential confounders. Final models were chosen coherently with the study objectives and the nature of the variables, i.e. variable selection was primarily based on coherence with the explicit DAGs. In final models age, sex, ISE and cholestatic and constitutional syndrome were not associated with *K-ras* mutational status (both in models with OCs corrected and uncorrected by TSL). The level of statistical significance was set at 0.05, and all tests are two-tailed. Analyses were performed using SPSS, version 12.0.

We used the DAG framework (Greenland et al., 1999; Hernán et al., 2002) to express our assumptions regarding the underlying causal structure of the variables. DAGs serve as a

graphical method to facilitate causal inference in epidemiology and other fields of research. We provide greater detail regarding DAG methodology in the Appendix A.

We used heuristic diagrams to assess different scenarios based on results from previous analyses (Porta et al., 2009c). The heuristic diagrams in Fig. A.1 are useful to construct DAGs. The DAG in the Fig. 1 includes time-dependent variables (namely OCs1, OCs2, OCs3, lipids1, lipids2, lipids3, EPC1 and EPC2), they are actually variables related to disease progression. OCs1 and lipids1 represent concentrations of OCs and lipids before development of EPC, OCs2 and lipids2 represent concentrations when EPC was in the initial subclinical phase (EPC1), OCs3 and lipids3 represent concentrations when EPC was symptomatic (EPC2). The PANKRAS II study obtained measures only of OCs3. The DAG in Fig. 1 also shows that: OCs1 influence the occurrence of *K-ras* mutations, OCs1, *K-ras* mutations, and other effects (U) are associated with development of EPC1, EPC1 causes signs and symptoms (including cholestatic syndrome or the constitutional syndrome), and affects lipids2 and OCs2. In the next disease stage, EPC2 affects OCs3 and lipids3, while the pathophysiological processes underlying some signs and symptoms (like the cholestatic syndrome or the constitutional syndrome) affect concentrations of lipids3. Therefore, some EPC signs and symptoms are associated with blood concentrations of some OCs (certainly, through lipids, perhaps directly as well) measured after the symptomatic diagnosis of EPC (Porta et al., 2009c). It follows that adjustment for certain signs and symptoms may be necessary to estimate the causal association between OC and *K-ras* mutations. Variables we adjusted for in our analyses are depicted with boxes around them, consistent with DAG notation.

3. Results

In models adjusted for age and sex, we compared the OR of a *K-ras* mutation in relation to OC levels both correcting for TSL and not correcting for TSL. Overall, we found that several OCs in tertiles were statistically significantly associated with the odds of *K-ras* mutation, but that additional adjustment for TSL strengthened the point estimates for p,p' -DDE, PCB 138 and PCB 153. TSL-corrected concentrations of p,p' -DDT, PCB 138 and PCB 153 were significantly associated with an increased odds of a *K-ras* mutated EPC (Table 1). When OCs in tertiles were uncorrected by TSL, only p,p' -DDT and HCB were statistically significantly associated with the odds of a *K-ras* mutation, with increasing levels of HCB associated with a decreased odds of a *K-ras* mutation. The ORs of *K-ras* mutation for TSL-corrected p,p' -DDE were slightly higher than those of TSL-uncorrected p,p' -DDE for the middle and upper tertiles compared with the lowest, respectively (1.61 and 2.35 vs. 1.16 and 1.74, respectively). For TSL-corrected PCBs 138 and 153, the ORs in the mid-tertile were lower than the ORs for TSL-uncorrected PCBs 138 and 153, however, this was not so for the ORs in the upper tertile. Patients in the upper tertile of TSL-corrected PCBs 138 and 153 were over 4 and 5 times more likely to have a *K-ras* mutated tumor than patients in the lower tertile, whereas the ORs were only 2.7 and 2.2 in the upper tertile of TSL-uncorrected PCBs 138 and 153, respectively. There was a statistically significant linear trend for these PCBs in the models with the TSL-corrected OCs in tertiles (p for trend 0.024 and 0.017, respectively), while the p -values for these TSL-uncorrected PCBs were not statistically significant. Only for HCB the p -value was smaller in the TSL-uncorrected

model (p -value 0.044) than in the model for TSL-corrected HCB (p -value 0.175) (Table 1). When we considered the OCs as continuous variables (log-transformed), the relations of *K-ras* with OCs corrected and uncorrected by lipids were very similar (similar ORs and p -values).

Additional adjustment for cholestatic syndrome, while lipid correcting, strengthened the associations between the following OCs in tertiles and the odds of a *K-ras* mutation: p,p' -DDT, PCB 138, and PCB 153. The pattern shown in models adjusted by age and sex was similar when the models were further adjusted by the cholestatic syndrome (Table 1). Most ORs from the models for TSL-corrected OCs in tertiles increased slightly when adjusting by cholestatic syndrome (as compared to models adjusted by age and sex), by contrast, the ORs for the TSL-uncorrected OCs in tertiles were not affected. Adjustment for constitutional syndrome resulted in slight changes in the ORs of TSL-corrected p,p' -DDT and β -HCH (Table A.2). ORs for the rest of the OCs, TSL-corrected and TSL-uncorrected, were not affected when the models were adjusted by the constitutional syndrome (i.e., parameters they were similar to those shown in Table 1, models adjusted by age and sex). The ORs in models adjusted by the interval from first symptom to blood extraction (ISE) increased slightly (both in the models corrected and uncorrected by TSL) compared with the models adjusted only by age and sex (Table A.3).

The direction of the change in the ORs of the OCs (TSL-corrected and TSL-uncorrected) when adjusted by signs, symptoms, syndromes and ISE is summarized in Table A.4. Adjustment for cholestatic syndrome and ISE together altered all 7 of the OCs we evaluated when additionally TSL corrected. Adjustment for ISE and pruritus altered levels of 6 of the 7 OCs when additionally TSL corrected. TSL correction was associated with greater percent change in estimates compared with not correcting for TSL. When the models were adjusted by constitutional syndrome, only the ORs of p,p' -DDT (TSL-corrected) and β -HCH (TSL-corrected and TSL-uncorrected) increased (changes between 10 and 20%) compared with the models without constitutional syndrome. There were no relevant changes when the models were adjusted by asthenia. When weight loss was included in the models, the odds of a *K-ras* mutation increased for TSL-corrected p,p' -DDT and decreased for TSL-corrected β -HCH. However, when the models were adjusted by cachexia, the most important change was observed in the lowering of the ORs of TSL-uncorrected HCB.

4. Discussion

Serum concentrations of TSL-corrected p,p' -DDT, PCB 138 and PCB 153 were associated with *K-ras* mutations in EPC patients (Porta et al., 2009b). Two OCs were associated with *K-ras* mutations when they were not TSL-corrected: p,p' -DDT and HCB. With the exception of HCB, the associations were stronger after TSL correction. In healthy populations, under the assumption that body concentrations of OCs are in a state of equilibrium (between blood and fat tissue), models corrected by TSL are deemed preferable than models uncorrected by TSL (Phillips et al., 1989). This assumption may not hold in populations with significant changes in lipid concentrations, such as EPC patients. The change in the OR after adjustment suggests that TSL confounded the association between p,p' -DDT, PCB 138 and PCB 153 and *K-ras*.

Therefore, TSL and EPC symptoms could be considered as confounders among EPC patients, and we provide DAGs to demonstrate these associations (Fig. 2a). Adjustment and modeling strategies alone cannot prove that TSL-corrected models are better than TSL-uncorrected models, because modeling strategies alone are inadequate to determine the impact of unmeasured confounding, time varying confounding, and additional bias, and demand an explicit causal framework to support these strategies. Residual confounding may persist due to the large variation in lipid concentrations and EPC related symptoms in patients with EPC. Diminished bias following adjustment for EPC symptoms would be expected if an unmeasured variable was associated with *K-ras* mutations and symptoms, but was not affected by OC levels. In Fig. 2a we summarize this causal structure, and based upon our structured analysis including and excluding likely confounding factors, the DAG that best exemplifies our findings is demonstrated in Fig. 2a.

OCs3 was the only measured OCs in our study, and it is the only OCs that can be obtained in case-control and case-case studies. However, OCs3 is a proxy for OCs1 and all our comments here on the association between OCs3 and *K-ras* mutations refer to the possible causal effect of OCs1 on *K-ras* mutations.

The results show that signs and symptoms affected the TSL-corrected OCs more than they affected TSL-uncorrected OCs, changes in most ORs were higher in TSL-corrected models than in TSL-uncorrected models (compared with models adjusted for age and sex). Cholestatic syndrome, constitutional syndrome, and vomiting reflect a range of severity that may not be fully captured by adjustment for either having such symptoms or not. Therefore, some degree of residual confounding may be present, even in cases where adjustment for EPC symptoms was appropriate. Inclusion of signs and symptoms in TSL-uncorrected models increased the ORs only slightly, indicating that signs and symptoms did not negatively bias the associations between *K-ras* mutation status and OCs.

Another important factor to consider is the disease stage (e.g., tumoral stage) when biological samples are obtained. In the case of pancreatic cancer, tumor stage may not be a powerful discriminatory variable, because most patients are diagnosed at stage IV (Porta, 2001; Porta et al., 2012). By contrast, the interval between first symptom and blood extraction (ISE) may be one of the possible accurate measures of duration of disease until blood draw. In principle, ISE is not a better proxy of disease progression than tumor stage because diseases do not progress uniformly as time increases. Thus, adjusting for ISE may be appropriate both when analyzing TSL-corrected and TSL-uncorrected OCs, but might not be sufficient. An aggressive disease process further implies that patients begin to develop different signs and symptoms (e.g., weight loss, jaundice, hypocholia).

Fig. 2a illustrates that adjustment for cholestatic syndrome and TSL at time 3 (when EPC was symptomatic), together with adjustment for age and sex are sufficient to control for confounding in the association between OCs at time 3 and *K-ras* mutation. Cholestatic syndrome and TSL together are a consequence of an unmeasured confounding factor, represented by a U on the DAG (Fig. 2a). The best candidates for U are the pathophysiological processes inherent to EPC. As control for an unmeasured confounding factor is impossible, measuring a variable that is affected by such a factor is one way to

control for it. Fig. 2b illustrates the causal structure under which adjustment for EPC-related symptoms and TSL is not appropriate. In this case, neither variable may be considered a confounding factor in the association between *K-ras* mutation and OCs at time 3. Therefore, adjustment for TSL or EPC symptoms would not diminish bias. However, as it is not possible to detect the underlying causal structure by comparing the magnitude of ORs adjusted and not adjusted for particular factors, the proposed causal structure must be determined *a priori*. Addressing such issues through a causal DAG approach is a good solution, as relying on change in estimate criteria –though commonly done –cannot reliably differentiate between true confounding factors and factors that introduce bias if they are conditioned on.

Additional criteria support the inclusion of TSL when evaluating the association between OCs and *K-ras* mutations. Den Hond et al. (2009), conclude that in terms of adjusted R^2 , the regression model with TSL-corrected OCs was superior to models with TSL-uncorrected OCs. In previous articles we concluded that correcting OCs by TSL may be inappropriate in patients with severe diseases as pancreatic cancer (Porta et al., 2009a). Schisterman et al. (2005) and Gaskins and Schisterman (2009) reported that lipid standardization, or the division of serum concentrations of PCBs by serum lipids (in a simulation study of patients with breast cancer) was highly prone to bias, but that simple lipid adjustment did not induce significant bias in similar settings. The reason for bias with TSL-standardized OCs may be that the formula to calculate TSL (Phillips et al., 1989) stemmed from a study in healthy patients, in whom an equilibrium exists between serum lipids and lipids in adipose tissue; as mentioned, such an equilibrium is rarely present among individuals with cancer (Porta et al., 2009c). Alternatively, lipids may not be true confounders of the relationship of interest, as suggested by Fig. 2b. Other formulas to calculate TSL are based on the same principle (Bergonzi et al., 2009).

Our work has some limitations. The study size was small. Furthermore, this is the largest study on OCs, genetic alterations and clinical information in pancreatic cancer, and one of the largest with analyses on OCs and tumor DNA in any type of cancer (Fryzek et al., 2006; Slebos et al., 2000). Also, a high response rate was achieved and a high proportion of cases were interviewed face-to-face and had blood drawn around the time of diagnosis. Our objective was to compare the alterations to the odds of a *Kras* mutation in relation to OC exposure from additional adjustment for lipid, disease symptoms and other clinical factors, in patients with EPC. As adjustment for TSL strengthened associations of interest, and based on *a priori* knowledge, in our view Fig. 2a best exemplifies that TSL and EPC symptoms are descendents of a common cause between the exposure and outcome of interest. Therefore, adjustment for these factors diminished confounding bias. If the study measures OCs prior to the onset of symptoms and lipid mobilization, correction for lipids and symptoms may not be necessary based upon our proposed DAGs. If OCs are measured after disease onset, then correcting by lipids may bias estimates, and one may have no alternative: crude OC concentrations will also be biased (Porta et al., 2009c).

It is important to note how different these two situations are: (1) OCs affect lipids, and (2) the disease process affects OCs and lipids. The present paper deals with (2) and in this situation, adjusting by symptoms of the disease (which reflect pathophysiologic processes

that causally influence lipids) while simultaneously adjusting for TSL would minimize bias, with the additional advantage that cancer symptoms are seldom or never instantaneous, i.e., they have a temporal dimension. There is also another situation (3): lipid correction or adjustment is a valid option (it is probably required) to deal with lack of information on fasting status, and when we are not in situation (1) (i.e., OCs do not affect lipids), nor are we in situation (2) (the disease has not affected OCs). In other words, correction of OCs by TSL is a valid option when the bias that may cause ignorance of fasting status is potentially much higher than the bias that could cause the influence of OCs on lipids.

If OCs alter lipid concentrations, profiles or patterns over a long time and subtly (which is likely) (Goncharov et al., 2008; Lee et al., 2011): (a) it will be scientifically relevant to focus on such changes as an intermediate outcome (e.g., influence of OCs on lipids as a main study aim); (b) it will be important to have longitudinal data with a long time between measurement(s) of OCs and measurement(s) of lipids; and (c) measurements of metabolic biomarkers and other intermediate biomarkers will also be of interest (Leijds et al., 2009; Porta, 2012).

In aggressive diseases that may alter lipid profiles, correcting OCs by lipids and adjusting the models for variables such as the cholestatic syndrome or the ISE may generate less bias or palliate some bias. Our findings might have implications as well for studying environmental causes of other clinically aggressive diseases. Nonetheless, further research is needed to refute or to confirm these results before methodological implications are considered. The rationale for the current practice of correcting OCs and other lipophilic biomarkers of exposure in research on aggressive diseases is yet unclear.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors gratefully acknowledge scientific and technical assistance provided by Magda Gasull, Joan Alguacil, and Yolanda Rovira.

Abbreviations:

CI	confidence interval
BMI	body mass index
DAG	causal directed acyclic graph
EPC	exocrine pancreatic cancer
HCBz	hexachlorobenzene
HCH	hexachlorocyclohexane
ISE	interval from first symptom of EPC to blood extraction

OCs	organochlorine compounds
OR	odds ratio
PCBs	polychlorinated biphenyls
<i>p,p'</i>-DDE	dichlorodiphenyldichloroethene
<i>p,p'</i>-DDT	dichlorodiphenyltrichloroethane
TSL	total serum lipids.

References

- Baris D, Kwak LW, Rothman N, Wilson W, Hartge P, 2000 Blood levels of organochlorines before and after chemotherapy among non-Hodgkin's lymphoma patients. *Cancer Epidem. Biomarkers* 9,193–197.
- Bergonzi R, De Palma G, Tomasi C, Ricossa MC, Apostoli P, 2009 Evaluation of different methods to determine total serum lipids for normalization of circulating organochlorine compounds. *Int. Arch. Occ. Env. Hea* 82, 1241–1247.
- Centers for Disease Control and Prevention, 2009 Fourth National Report on Human Exposure to Environmental Chemicals Centers for Disease Control and Prevention, Atlanta.
- Crous-Bou M, 2009 Clinical and Environmental Influences on the Prevalence of Mutations in the K-ras Oncogene in Patients with Pancreatic Ductal Adenocarcinoma [doctoral dissertation, dir.: Porta M]. Barcelona: Universitat Autònoma de Barcelona <<http://www.imim.es/programesrecerca/epidemiologia/documentsgreem.html>>.
- Den Hond E, Govarts E, Bruckers L, Schoeters G, 2009 Determinants of polychlorinated aromatic hydrocarbons in serum in three age classes. Methodological implications for human biomonitoring. *Environ. Res.* 109, 495–502. [PubMed: 19278675]
- Fryzek JP, Garabrant DH, Schenk M, Sarkar FH, 2006 The association between selected risk factors for pancreatic cancer and the expression of p53 and K-ras codon 12 mutations. *Int. J. Gastro. Cancer* 37,139–145.
- Gaskins AJ, Schisterman EF, 2009 The effect of lipid adjustment on the analysis of environmental contaminants and the outcome of human health risks. *Methods Mol. Biol* 580, 371–381. [PubMed: 19784610]
- Gasull M, Porta M, Pumarega J, Vioque J, Morales E, Grimalt JO, Malats N, 2010 The relative influence of diet and serum concentrations of organochlorine compounds on K-ras mutations in exocrine pancreatic cancer. *Chemosphere* 79, 686–697. [PubMed: 20350743]
- Gasull M, Pumarega J, Rovira G, López T, Alguacil J, Porta M, 2013 Relative effects of educational level and occupational social class on body concentrations of persistent organic pollutants in a representative sample of the general population of Catalonia, Spain. *Environ. Int* 60, 190–201. [PubMed: 24064380]
- Goncharov A, Haase RF, Santiago-Rivera A, Morse G, Carpenter DO, 2008 High serum PCBs are associated with elevation of serum lipids and cardiovascular disease in a Native American population. *Environ. Res* 106, 226–239. [PubMed: 18054906]
- Greenland S, Pearl J, Robins JM, 1999 Causal diagrams for epidemiologic research. *Epidemiology* 10, 37–48. [PubMed: 9888278]
- Hernán MA, Hernández-Díaz S, Werler MM, Mitchell AA, 2002 Causal knowledge as a prerequisite for confounding evaluation: an application to birth defects epidemiology. *Am. J. Epidemiol* 155, 176–184. [PubMed: 11790682]
- Kritchevsky SB, Wilcosky TC, Morris DL, Truong KN, Tyroler HA, 1991 Changes in plasma lipid and lipoprotein cholesterol and weight prior to the diagnosis of cancer. *Cancer Res* 51, 3198–3203. [PubMed: 2039996]

- Lee DH, Steffes MW, Sjödin A, Needham LL, Jacobs DR, Jr., 2011 Low dose organochlorine pesticides and polychlorinated biphenyls predict obesity, dyslipidemia, and insulin resistance among people free of diabetes. *PLoS ONE* 6, e15977. [PubMed: 21298090]
- Leijds MM, Koppe JG, Olie K, van Aalderen WM, de Voogt P, ten Tusscher GW, 2009 Effects of dioxins, PCBs, and PBDEs on immunology and hematology in adolescents. *Environ. Sci. Technol* 43, 7946–7951. [PubMed: 19921918]
- Phillips DL, Pirkle JL, Burse VW, Bernert JT, Jr., Henderson LO, Needham LL, 1989 Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. *Arch. Environ. Con. Tox* 18, 495–500.
- Porta M, 2001 Role of organochlorine compounds in the etiology of pancreatic cancer: a proposal to develop methodological standards. *Epidemiology* 12, 272–276. [PubMed: 11246593]
- Porta M, Fernandez E, Alguacil J, 2003 Semiology, proteomics, and the early detection of symptomatic cancer. *J. Clin. Epidemiol* 56, 815–819. [PubMed: 14505764]
- Porta M, Pumarega J, Ferrer-Armengou O, Lóez T, Alguacil J, Malats N, Fernández E, 2007 Timing of blood extraction in epidemiologic and proteomic studies: Results and proposals from the PANKRAS II Study. *Eur. J. Epidemiol* 22, 577–588. [PubMed: 17636417]
- Porta M, Puigdomènech E, Ballester F, Selva J, Ribas-Fitó N, Llop S, López T, 2008a Monitoring concentrations of persistent organic pollutants in the general population: the international experience. *Environ. Int* 34, 546–561. [PubMed: 18054079]
- Porta M, Ferrer-Armengou O, Pumarega J, López T, Crous-Bou M, Alguacil J, Fitó M, Morales E, Puigdomènech E, Gupta N, 2008b Exocrine pancreatic cancer clinical factors were related to timing of blood extraction and influenced serum concentrations of lipids. *J. Clin. Epidemiol* 61, 695–704. [PubMed: 18538264]
- Porta M, Jarrod M, López T, Pumarega J, Puigdomènech E, Malats N, Grimalt JO, Real FX, 2009a Correcting serum concentrations of organochlorine compounds by lipids: alternatives to the organochlorine/total lipids ratio. *Environ. Int* 35, 1080–1085. [PubMed: 19581003]
- Porta M, López T, Pumarega J, Jarrod M, Crous-Bou M, Grimalt JO, Malats N, Real FX, 2009b In pancreatic ductal adenocarcinoma blood concentrations of some organochlorine compounds and coffee intake are independently associated with KRAS mutations. *Mutagenesis* 24, 513–521. [PubMed: 19797353]
- Porta M, Pumarega J, López T, Jarrod M, Marco E, Grimalt JO, 2009c Influence of tumor stage, symptoms and time of blood draw on serum concentrations of organochlorine compounds in exocrine pancreatic cancer. *Cancer Causes Control* 20, 1893–1906. [PubMed: 19562493]
- Porta M, Pumarega J, Guarner L, Malats N, Solà R, Real FX, 2012 Relationships of hepatic and pancreatic biomarkers with the cholestatic syndrome and tumor stage in pancreatic cancer. *Biomarkers* 17, 557–565. [PubMed: 22793268]
- Porta M, 2012 Human contamination by environmental chemical pollutants: can we assess it more properly? *Prev. Med.* 55, 560–562. [PubMed: 23036521]
- Porta M, ed. *A Dictionary of Epidemiology*, 2008 5th edition. New York, Oxford University Press.
- Reed L, Buchner V, Tchounwou PB, 2007 Environmental toxicology and health effects associated with hexachlorobenzene exposure. *Rev. Environ. Health* 22, 213–243. [PubMed: 18078005]
- Schisterman EF, Whitcomb BW, Louis GM, Louis TA, 2005 Lipid adjustment in the analysis of environmental contaminants and human health risks. *Environ. Health Persp* 113, 853–857.
- Slebos RJ, Hoppin JA, Tolbert PE, Holly EA, Brock JW, Flake GP, Taylor JA, 2000 K-ras and p53 in pancreatic cancer: association with medical history, histopathology, and environmental exposures in a population-based study. *Cancer Epidemi. Biomark. Prev* 9, 1223–1232.
- Spinelli JJ, Ng CH, Weber JP, Connors JM, Gascoyne RD, Berry BR, Gallagher RP, 2007 Organochlorines and risk of non-Hodgkin lymphoma. *Int. J. Cancer* 121, 2767–2775. [PubMed: 17722095]
- Wolff MS, Zeleniuch-Jacquotte A, Dubin N, Toniolo P, 2000 Risk of breast cancer and organochlorine exposure. *Cancer Epidemi. Biomark. Prev* 9, 271–277.
- Wolff MS, Anderson HA, Britton JA, Rothman N, 2007 Pharmacokinetic variability and modern epidemiology—the example of dichlorodiphenyltrichloroethane, body mass index, and birth cohort. *Cancer Epidemi. Biomark. Prev* 16, 1925–1930.

HIGHLIGHTS

- The appropriateness of correcting blood levels of OCs for lipids is unclear.
- Causal directed acyclic graphs (DAGs) clarified covariates necessary to minimize bias.
- Lipid-correction and adjustment for cholestasis yielded the highest odds ratios.
- Adjustment for lipids and failure to adjust for symptoms may induce bias.
- Models with lipid-uncorrected organochlorine compounds (OCs) may be biased too.

Fig. 1. DAG to express effects of EPC progression on lipids, OCs and symptoms. ‘Serum lipids’ refers to Total serum lipids (TSL). Dashed lines represent the causal relationship between unmeasured factors (in most instances, the upstream variable was unmeasured), solid lines denote that such factors were measured. A square around a variable denotes that the variable was measured in the study.

Fig. 2.

a. DAG 2a: Adjustment for age, sex, serum lipids (TSL) at time 3 and cholestatic syndrome (which may serve as a proxy for other EPC-related symptoms, i.e., nausea, ISE, pruritus) is appropriate in investigating the association between OC levels and odds of K-ras mutation.

b. DAG 2b: Adjustment for TSL at time 3 and cholestatic syndrome (which may serve as a proxy for other EPC-related symptoms i.e., nausea, ISE, pruritus) is unnecessary to evaluate the association between OC levels and odds of K-ras mutation. 'Serum lipids' refers to Total

serum lipids (TSL). Dashed lines represent the causal relationship between unmeasured factors (in most instances, the upstream variable was unmeasured), solid lines denote that such factors were measured. A square around a variable denotes that the variable was measured in the study.

Table 1

Odds of mutation in the *K-ras* oncogene by concentrations of organochlorine compounds in cases of EPC.

Teriles of OC	OCs total lipids corrected ^a (N=103)			OCs total lipids uncorrected ^a (N=103)			OCs total lipids corrected ^b (N=103)			OCs total lipids uncorrected ^b (N=103)		
	OR	(95% CI)	P	OR	(95% CI)	P	OR	(95% CI)	P	OR	(95% CI)	P
<i>p,p'</i> -DDT												
Low tertile	1.00		0.008 ^c	1.00		0.035 ^d	1.00		0.009 ^c	1.00		0.046 ^d
Mid tertile	26.8	(3.06–234.5)		3.50	(1.08–11.30)		31.1	(3.33–290.6)		3.66	(1.06–12.62)	
Upper tertile	2.61	(0.80–8.52)		3.47	(0.94–12.85)		2.17	(0.64–7.37)		3.36	(0.89–12.69)	
<i>p,p'</i> -DDE												
Low tertile	1.00		0.176 ^d	1.00		0.393 ^d	1.00		0.245 ^d	1.00		0.566 ^d
Mid tertile	1.61	(0.53–4.85)		1.16	(0.38–3.48)		1.47	(0.47–4.54)		1.09	(0.35–3.42)	
Upper tertile	2.35	(0.67–8.25)		1.74	(0.50–6.10)		2.13	(0.58–7.73)		1.47	(0.41–5.28)	
<i>PCB 138</i>												
Low tertile	1.00		0.024 ^d	1.00		0.083 ^d	1.00		0.014 ^d	1.00		0.075 ^d
Mid tertile	2.08	(0.62–6.96)		2.40	(0.74–7.83)		2.00	(0.58–6.87)		2.37	(0.72–7.84)	
Upper tertile	4.02	(1.16–13.85)		2.68	(0.83–8.67)		4.84	(1.35–17.36)		2.81	(0.85–9.26)	
<i>PCB 153</i>												
Low tertile	1.00		0.017 ^d	1.00		0.157 ^d	1.00		0.012 ^d	1.00		0.158 ^d
Mid tertile	1.66	(0.55–5.00)		2.37	(0.69–8.13)		1.56	(0.50–4.83)		2.31	(0.67–7.90)	
Upper tertile	5.17	(1.30–20.47)		2.24	(0.72–7.00)		6.03	(1.47–24.77)		2.25	(0.71–7.12)	
<i>PCB 180</i>												
Low tertile	1.00		0.161 ^c	1.00		0.203 ^d	1.00		0.171 ^c	1.00		0.219 ^d
Mid tertile	0.41	(0.13–1.33)		1.51	(0.49–4.70)		0.43	(0.13–1.38)		1.38	(0.44–4.37)	
Upper tertile	1.15	(0.30–4.33)		2.20	(0.65–7.50)		1.21	(0.31–4.72)		2.17	(0.63–7.44)	
<i>HCB</i>												
Low tertile	1.00		0.175 ^d	1.00		0.044 ^c	1.00		0.388 ^c	1.00		0.048 ^c
Mid tertile	0.94	(0.27–3.22)		1.81	(0.48–6.77)		1.14	(0.32–4.01)		1.82	(0.47–7.03)	
Upper tertile	0.34	(0.08–1.44)		0.28	(0.07–1.15)		0.44	(0.10–1.93)		0.28	(0.06–1.23)	
<i>β-HCH</i>												

Teriles of OC	OCs total lipids corrected ^a (N =103)				OCs total lipids uncorrected ^a (N =103)				OCs total lipids corrected ^b (N =103)				OCs total lipids uncorrected ^b (N =103)			
	OR	(95% CI)	P		OR	(95% CI)	P		OR	(95% CI)	P		OR	(95% CI)	P	
Low tertile	1.00		0.133 ^c		1.00		0.143 ^c		1.00		0.193 ^c		1.00		0.139 ^c	
Mid tertile	3.63	(1.03–12.77)			3.06	(0.95–9.82)			3.28	(0.91–11.81)			3.04	(0.93–9.90)		
Upper tertile	1.44	(0.44–4.70)			2.19	(0.59–8.06)			1.55	(0.46–5.28)			2.43	(0.64–9.22)		

^aOR adjusted by age and sex.

^bOR adjusted by age, sex and cholestatic syndrome.

^cp- Value derived from Wald test.

^dTest for linear trend (multivariate analogue of Mantel’s extension test).