Addendum: Abdominal Wall Reconstruction: An Integrated Approach

Stefanos Boukovalas, MD¹ Geoffrey Sisk, MD¹ Jesse C. Selber, MD, MPH, FACS¹

¹Department of Plastic Surgery, University of Texas MD Anderson Cancer Center, Houston, Texas

Address for correspondence Jesse C. Selber, MD, MPH, FACS, Department of Plastic Surgery, The University of Texas, MD Anderson Cancer Center, Houston, TX (e-mail: jcselber@mdanderson.org).

Semin Plast Surg 2018;32:199–202.

The authors have informed the publisher that the Suggested Readings list was mistakenly omitted in the above article in *Seminars in Plastic Surgery*, Volume 32, Number 3, 2018 (DOI: 10.1055/s-0038-1667062). The list has now been provided here.

Suggested Readings

- Adelman DM, Selber JC, Butler CE. Bovine versus Porcine Acellular Dermal Matrix: A Comparison of Mechanical Properties. *Plast Reconstr Surg Glob Open* 2014;2(05):e155
- Albino FP, Patel KM, Nahabedian MY, Sosin M, Attinger CE, Bhanot P. Does mesh location matter in abdominal wall reconstruction? A systematic review of the literature and a summary of recommendations. *Plast Reconstr Surg* 2013;132(05):1295–1304
- Awad ZT, Puri V, LeBlanc K, et al. Mechanisms of ventral hernia recurrence after mesh repair and a new proposed classification. *J Am Coll Surg* 2005;201(01):132–140
- Badylak SF, Coffey AC, Lantz GC, Tacker WA, Geddes LA. Comparison of the resistance to infection of intestinal submucosa arterial autografts versus polytetrafluoroethylene arterial prostheses in a dog model. *J Vasc Surg* 1994;19(03):465–472
- Bailey SH, Saint-Cyr M, Wong C, et al. The single dominant medial row perforator DIEP flap in breast reconstruction: three-dimensional perforasome and clinical results. *Plast Reconstr Surg* 2010;126(03):739–751
- Bauer JJ, Harris MT, Gorfine SR, Kreel I. Rives-Stoppa procedure for repair of large incisional hernias: experience with 57 patients. *Hernia* 2002;6(03):120–123
- Bauer JJ, Harris MT, Kreel I, Gelernt IM. Twelve-year experience with expanded polytetrafluoroethylene in the repair of abdominal wall defects. *Mt Sinai J Med* 1999;66(01):20–25
- Baumann DP, Butler CE. Soft tissue coverage in abdominal wall reconstruction. *Surg Clin North Am* 2013;93(05):1199–1209
- Bellon JM, Rodriguez M, Garcia-Hondurilla N, Gomez-Gil V, Pascual G, Bujan J. Postimplant behavior of lightweight polypropylene meshes in an experimental model of abdominal hernia. *J Invest Surg* 2008;21(05):280–287
- Bingener J, Buck L, Richards M, Michalek J, Schwesinger W, Sirinek K. Long-term outcomes in laparoscopic vs open ventral hernia repair. *Arch Surg* 2007;142(06):562–567
- Blatnik J, Jin J, Rosen M. Abdominal hernia repair with bridging acellular dermal matrix—an expensive hernia sac. *Am J Surg* 2008;196(01):47–50
- Booth JH, Garvey PB, Baumann DP, et al. Primary fascial closure with mesh reinforcement is superior to bridged mesh repair for abdominal wall reconstruction. *J Am Coll Surg* 2013;217(06):999–1009
- Burger JW, Halm JA, Wijsmuller AR, ten Raa S, Jeekel J. Evaluation of new prosthetic meshes for ventral hernia repair. *Surg Endosc* 2006;20(08):1320–1325
- Burger JW, Luijendijk RW, Hop WC, Halm JA, Verdaasdonk EG, Jeekel J. Long-term follow-up of a randomized controlled trial of suture versus mesh repair of incisional hernia. *Ann Surg* 2004;240(04):578–583, discussion 583–585
- Butler CE, Langstein HN, Kronowitz SJ. Pelvic, abdominal, and chest wall reconstruction with AlloDerm in patients at increased risk for mesh-related complications. *Plast Reconstr Surg* 2005;116(05):1263–1275, discussion 1276–1277
- Butler CE, Prieto VG. Reduction of adhesions with composite AlloDerm/polypropylene mesh implants for abdominal wall reconstruction. *Plast Reconstr Surg* 2004;114(02):464–473
- Carbonell AM, Criss CN, Cobb WS, Novitsky YW, Rosen MJ. Outcomes of synthetic mesh in contaminated ventral hernia repairs. *J Am Coll Surg* 2013;217(06):991–998
- Clagett GP, Bowers BL, Lopez-Viego MA, et al. Creation of a neo-aortoiliac system from lower extremity deep and superficial veins. *Ann Surg* 1993;218(03):239–248, discussion 248–249
- Clemens MW, Selber JC, Liu J, et al. Bovine versus porcine acellular dermal matrix for complex abdominal wall reconstruction. *Plast Reconstr Surg* 2013;131(01):71–79
- Cobb WS, Harris JB, Lokey JS, McGill ES, Klove KL. Incisional herniorrhaphy with intraperitoneal composite mesh: a report of 95 cases. *Am Surg* 2003;69(09):784–787
- Cobb WS, Kercher KW, Heniford BT. The argument for lightweight polypropylene mesh in hernia repair. *Surg Innov* 2005;12(01):63–69
- de Vries Reilingh TS, van Geldere D, Langenhorst B, et al. Repair of large midline incisional hernias with polypropylene mesh: comparison of three operative techniques. *Hernia* 2004;8(01):56–59
- de Vries Reilingh TS, van Goor H, Charbon JA, et al. Repair of giant midline abdominal wall hernias: “components separation technique” versus prosthetic repair: interim analysis of a randomized controlled trial. *World J Surg* 2007;31(04):756–763

Issue Theme Business and Technology;
Guest Editor: Edward M. Reece, MD, MBA

Copyright © 2018 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

DOI <https://doi.org/10.1055/s-0038-1673696>.
ISSN 1535-2188.

- 24 de Vries Reilingh TS, van Goor H, Koppe MJ, Bodegom ME, Hendriks T, Bleichrodt RP. Interposition of polyglactin mesh does not prevent adhesion formation between viscera and polypropylene mesh. *J Surg Res* 2007;140(01):27–30
- 25 de Vries Reilingh TS, van Goor H, Rosman C, et al. “Components separation technique” for the repair of large abdominal wall hernias. *J Am Coll Surg* 2003;196(01):32–37
- 26 Deeken CR, Abdo MS, Frisella MM, Matthews BD. Physicomechanical evaluation of polypropylene, polyester, and polytetrafluoroethylene meshes for inguinal hernia repair. *J Am Coll Surg* 2011;212(01):68–79
- 27 Deeken CR, Abdo MS, Frisella MM, Matthews BD. Physicomechanical evaluation of absorbable and nonabsorbable barrier composite meshes for laparoscopic ventral hernia repair. *Surg Endosc* 2011;25(05):1541–1552
- 28 den Hartog D, Dur AH, Tuinebreijer WE, Kreis RW. Open surgical procedures for incisional hernias. *Cochrane Database Syst Rev* 2008;(03):CD006438
- 29 Diaz JJ Jr, Guy J, Berkes MB, Guillaumondegui O, Miller RS. Acellular dermal allograft for ventral hernia repair in the compromised surgical field. *Am Surg* 2006;72(12):1181–1187, discussion 1187–1188
- 30 DiBello JN Jr, Moore JH Jr. Sliding myofascial flap of the rectus abdominus muscles for the closure of recurrent ventral hernias. *Plast Reconstr Surg* 1996;98(03):464–469
- 31 Dunne JR, Malone DL, Tracy JK, Napolitano LM. Abdominal wall hernias: risk factors for infection and resource utilization. *J Surg Res* 2003;111(01):78–84
- 32 Emans PJ, Schreinemacher MH, Gijbels MJ, et al. Polypropylene meshes to prevent abdominal herniation. Can stable coatings prevent adhesions in the long term? *Ann Biomed Eng* 2009;37(02):410–418
- 33 Ennis LS, Young JS, Gampper TJ, Drake DB. The “open-book” variation of component separation for repair of massive midline abdominal wall hernia. *Am Surg* 2003;69(09):733–742, discussion 742–743
- 34 Espinosa-de-los-Monteros A, de la Torre JI, Marrero I, Andrades P, Davis MR, Vásconez LO. Utilization of human cadaveric acellular dermis for abdominal hernia reconstruction. *Ann Plast Surg* 2007;58(03):264–267
- 35 Fabian TC, Croce MA, Pritchard FE, et al. Planned ventral hernia. Staged management for acute abdominal wall defects. *Ann Surg* 1994;219(06):643–650, discussion 651–653
- 36 Finan KR, Vick CC, Kiefe CI, Neumayer L, Hawn MT. Predictors of wound infection in ventral hernia repair. *Am J Surg* 2005;190(05):676–681
- 37 Flum DR, Horvath K, Koepsell T. Have outcomes of incisional hernia repair improved with time? A population-based analysis. *Ann Surg* 2003;237(01):129–135
- 38 Franz MG. The biology of hernias and the abdominal wall. *Hernia* 2006;10(06):462–471
- 39 Garvey PB, Giordano SA, Baumann DP, Liu J, Butler CE. Long-Term Outcomes after Abdominal Wall Reconstruction with Acellular Dermal Matrix. *J Am Coll Surg* 2017;224(03):341–350
- 40 Garvey PB, Martinez RA, Baumann DP, Liu J, Butler CE. Outcomes of abdominal wall reconstruction with acellular dermal matrix are not affected by wound contamination. *J Am Coll Surg* 2014;219(05):853–864
- 41 Ghali S, Turza KC, Baumann DP, Butler CE. Minimally invasive component separation results in fewer wound-healing complications than open component separation for large ventral hernia repairs. *J Am Coll Surg* 2012;214(06):981–989
- 42 Giordano S, Garvey PB, Baumann DP, Liu J, Butler CE. Primary fascial closure with biologic mesh reinforcement results in lesser complication and recurrence rates than bridged biologic mesh repair for abdominal wall reconstruction: A propensity score analysis. *Surgery* 2017;161(02):499–508
- 43 Giordano S, Schaverien M, Garvey PB, Baumann DP, Liu J, Butler CE. Advanced age does not affect abdominal wall reconstruction outcomes using acellular dermal matrix: A comparative study using propensity score analysis. *Am J Surg* 2017;213(06):1046–1052
- 44 Glasberg SB, D’Amico RA. Use of regenerative human acellular tissue (AlloDerm) to reconstruct the abdominal wall following pedicle TRAM flap breast reconstruction surgery. *Plast Reconstr Surg* 2006;118(01):8–15
- 45 Goodney PP, Birkmeyer CM, Birkmeyer JD. Short-term outcomes of laparoscopic and open ventral hernia repair: a meta-analysis. *Arch Surg* 2002;137(10):1161–1165
- 46 Gravvanis A, Dionysiou DD, Chandrasekharan L, Francis I, Smith RW. Paramuscular and paraneural perforators in DIEAP flaps: radiographic findings and clinical application. *Ann Plast Surg* 2009;63(06):610–615
- 47 Gupta A, Zahriya K, Mullens PL, Salmassi S, Keshishian A. Ventral herniorrhaphy: experience with two different biosynthetic mesh materials, Surgisis and Alloderm. *Hernia* 2006;10(05):419–425
- 48 Guyatt G, Gutterman D, Baumann MH, et al. Grading strength of recommendations and quality of evidence in clinical guidelines: report from an american college of chest physicians task force. *Chest* 2006;129(01):174–181
- 49 Harrell AG, Novitsky YW, Kercher KW, et al. In vitro infectability of prosthetic mesh by methicillin-resistant *Staphylococcus aureus*. *Hernia* 2006;10(02):120–124
- 50 Harrell AG, Novitsky YW, Peindl RD, et al. Prospective evaluation of adhesion formation and shrinkage of intra-abdominal prosthetics in a rabbit model. *Am Surg* 2006;72(09):808–813, discussion 813–814
- 51 Heartsill L, Richards ML, Arfai N, et al. Open Rives-Stopppa ventral hernia repair made simple and successful but not for everyone. *Hernia* 2005;9(02):162–166
- 52 Helton WS, Fischella PM, Berger R, Horgan S, Espat NJ, Abcarian H. Short-term outcomes with small intestinal submucosa for ventral abdominal hernia. *Arch Surg* 2005;140(06):549–560, discussion 560–562
- 53 Hijjawi JB, Blondeel PN. Advancing deep inferior epigastric artery perforator flap breast reconstruction through multidetector row computed tomography: an evolution in preoperative imaging. *J Reconstr Microsurg* 2010;26(01):11–20
- 54 Holton LH III, Kim D, Silverman RP, Rodriguez ED, Singh N, Goldberg NH. Human acellular dermal matrix for repair of abdominal wall defects: review of clinical experience and experimental data. *J Long Term Eff Med Implants* 2005;15(05):547–558
- 55 Houck JP, Rypins EB, Sarfeh IJ, Juler GL, Shimoda KJ. Repair of incisional hernia. *Surg Gynecol Obstet* 1989;169(05):397–399
- 56 Iqbal CW, Pham TH, Joseph A, Mai J, Thompson GB, Sarr MG. Long-term outcome of 254 complex incisional hernia repairs using the modified Rives-Stopppa technique. *World J Surg* 2007;31(12):2398–2404
- 57 Jarman-Smith ML, Bodamyali T, Stevens C, Howell JA, Horrocks M, Chaudhuri JB. Porcine collagen crosslinking, degradation and its capability for fibroblast adhesion and proliferation. *J Mater Sci Mater Med* 2004;15(08):925–932
- 58 Jin J, Rosen MJ, Blatnik J, et al. Use of acellular dermal matrix for complicated ventral hernia repair: does technique affect outcomes? *J Am Coll Surg* 2007;205(05):654–660
- 59 Jones JW, Jurkovich GJ. Polypropylene mesh closure of infected abdominal wounds. *Am Surg* 1989;55(01):73–76
- 60 Kanters AE, Krpata DM, Blatnik JA, Novitsky YM, Rosen MJ. Modified hernia grading scale to stratify surgical site occurrence after open ventral hernia repairs. *J Am Coll Surg* 2012;215(06):787–793
- 61 Karakousis CP, Volpe C, Tanski J, Colby ED, Winston J, Driscoll DL. Use of a mesh for musculoaponeurotic defects of the abdominal

- wall in cancer surgery and the risk of bowel fistulas. *J Am Coll Surg* 1995;181(01):11–16
- 62 Kim H, Bruen K, Vargo D. Acellular dermal matrix in the management of high-risk abdominal wall defects. *Am J Surg* 2006;192(06):705–709
 - 63 Kingsnorth A. The management of incisional hernia. *Ann R Coll Surg Engl* 2006;88(03):252–260
 - 64 Kingsnorth A, LeBlanc K. Hernias: inguinal and incisional. *Lancet* 2003;362(9395):1561–1571
 - 65 Klinge U, Conze J, Kronen CJ, Schumpelick V. Incisional hernia: open techniques. *World J Surg* 2005;29(08):1066–1072
 - 66 Kolker AR, Brown DJ, Redstone JS, Scarpinato VM, Wallack MK. Multilayer reconstruction of abdominal wall defects with acellular dermal allograft (AlloDerm) and component separation. *Ann Plast Surg* 2005;55(01):36–41, discussion 41–42
 - 67 Korenkov M, Beckers A, Koebke J, Lefering R, Tiling T, Troidl H. Biomechanical and morphological types of the linea alba and its possible role in the pathogenesis of midline incisional hernia. *Eur J Surg* 2001;167(12):909–914
 - 68 Korenkov M, Neugebauer E. Comments on the letter from S. Petersen and K. Ludwig concerning our paper “Classification and surgical treatment of the incisional hernia. Results of expert meeting.” *Langenbeck's Arch Surg* 386:65–73. *Langenbecks Arch Surg* 2001;386(04):310–311
 - 69 Korenkov M, Paul A, Sauerland S, et al. Classification and surgical treatment of incisional hernia. Results of an experts' meeting. *Langenbecks Arch Surg* 2001;386(01):65–73
 - 70 Leber GE, Garb JL, Alexander AI, Reed WP. Long-term complications associated with prosthetic repair of incisional hernias. *Arch Surg* 1998;133(04):378–382
 - 71 LeBlanc KA. Laparoscopic incisional and ventral hernia repair: complications-how to avoid and handle. *Hernia* 2004;8(04):323–331
 - 72 Levine JP, Karp NS. Restoration of abdominal wall integrity as a salvage procedure in difficult recurrent abdominal wall hernias using a method of wide myofascial release. *Plast Reconstr Surg* 2001;107(03):707–716, discussion 717–718
 - 73 Lindström D, Sadr Azodi O, Wladis A, et al. Effects of a perioperative smoking cessation intervention on postoperative complications: a randomized trial. *Ann Surg* 2008;248(05):739–745
 - 74 Lowe JB, Garza JR, Bowman JL, Rohrich RJ, Strodel WE. Endoscopically assisted “components separation” for closure of abdominal wall defects. *Plast Reconstr Surg* 2000;105(02):720–729, quiz 730
 - 75 Luijendijk RW, Hop WC, van den Tol MP, et al. A comparison of suture repair with mesh repair for incisional hernia. *N Engl J Med* 2000;343(06):392–398
 - 76 Maas SM, de Vries RS, van Goor H, de Jong D, Bleichrodt RP. Endoscopically assisted “components separation technique” for the repair of complicated ventral hernias. *J Am Coll Surg* 2002;194(03):388–390
 - 77 Malone DL, Genuit T, Tracy JK, Gannon C, Napolitano LM. Surgical site infections: reanalysis of risk factors. *J Surg Res* 2002;103(01):89–95
 - 78 Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR; Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. *Infect Control Hosp Epidemiol* 1999;20(04):250–278, quiz 279–280
 - 79 Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR; Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. Guideline for Prevention of Surgical Site Infection, 1999. *Am J Infect Control* 1999;27(02):97–132, quiz 133–134, discussion 96
 - 80 Martín-Duce A, Noguezales F, Villeta R, et al. Modifications to Rives technique for midline incisional hernia repair. *Hernia* 2001;5(02):70–72
 - 81 Maurice SM, Skeete DA. Use of human acellular dermal matrix for abdominal wall reconstructions. *Am J Surg* 2009;197(01):35–42
 - 82 Milburn ML, Holton LH, Chung TL, et al. Acellular dermal matrix compared with synthetic implant material for repair of ventral hernia in the setting of peri-operative *Staphylococcus aureus* implant contamination: a rabbit model. *Surg Infect (Larchmt)* 2008;9(04):433–442
 - 83 Milburn ML, Shah PK, Friedman EB, et al. Laparoscopically assisted components separation technique for ventral incisional hernia repair. *Hernia* 2007;11(02):157–161
 - 84 Nemeth NL, Butler CE. Complex torso reconstruction with human acellular dermal matrix: long-term clinical follow-up. *Plast Reconstr Surg* 2009;123(01):192–196
 - 85 Novitsky YW, Elliott HL, Orenstein SB, Rosen MJ. Transversus abdominis muscle release: a novel approach to posterior component separation during complex abdominal wall reconstruction. *Am J Surg* 2012;204(05):709–716
 - 86 Novitsky YW, Fayeizadeh M, Majumder A, Neupane R, Elliott HL, Orenstein SB. Outcomes of Posterior Component Separation With Transversus Abdominis Muscle Release and Synthetic Mesh Sublay Reinforcement. *Ann Surg* 2016;264(02):226–232
 - 87 Novitsky YW, Harrell AG, Cristiano JA, et al. Comparative evaluation of adhesion formation, strength of ingrowth, and textile properties of prosthetic meshes after long-term intra-abdominal implantation in a rabbit. *J Surg Res* 2007;140(01):6–11
 - 88 Paton BL, Novitsky YW, Zerey M, Sing RF, Kercher KW, Heniford BT. Management of infections of polytetrafluoroethylene-based mesh. *Surg Infect (Larchmt)* 2007;8(03):337–341
 - 89 Patton JH Jr, Berry S, Kralovich KA. Use of human acellular dermal matrix in complex and contaminated abdominal wall reconstructions. *Am J Surg* 2007;193(03):360–363, discussion 363
 - 90 Pauli EM, Wang J, Petro CC, Juza RM, Novitsky YW, Rosen MJ. Posterior component separation with transversus abdominis release successfully addresses recurrent ventral hernias following anterior component separation. *Hernia* 2015;19(02):285–291
 - 91 Pessaux P, Lermite E, Blezel E, et al; French Associations for Surgical Research. Predictive risk score for infection after inguinal hernia repair. *Am J Surg* 2006;192(02):165–171
 - 92 Petersen S, Henke G, Freitag M, Faulhaber A, Ludwig K. Deep prosthesis infection in incisional hernia repair: predictive factors and clinical outcome. *Eur J Surg* 2001;167(06):453–457
 - 93 Pierce RA, Perrone JM, Nimeri A, et al. 120-day comparative analysis of adhesion grade and quantity, mesh contraction, and tissue response to a novel omega-3 fatty acid bioabsorbable barrier macroporous mesh after intraperitoneal placement. *Surg Innov* 2009;16(01):46–54
 - 94 Plymale MA, Davenport DL, Dugan A, Zachem A, Roth JS. Ventral hernia repair with poly-4-hydroxybutyrate mesh. *Surg Endosc* 2018;32(04):1689–1694
 - 95 Ramirez OM, Ruas E, Dellon AL. “Components separation” method for closure of abdominal-wall defects: an anatomic and clinical study. *Plast Reconstr Surg* 1990;86(03):519–526
 - 96 Rosen MJ, Jin J, McGee MF, Williams C, Marks J, Ponsky JL. Laparoscopic component separation in the single-stage treatment of infected abdominal wall prosthetic removal. *Hernia* 2007;11(05):435–440
 - 97 Rosen MJ, Williams C, Jin J, et al. Laparoscopic versus open-component separation: a comparative analysis in a porcine model. *Am J Surg* 2007;194(03):385–389
 - 98 Roth JS, Anthone GJ, Selzer DJ, et al. Prospective evaluation of poly-4-hydroxybutyrate mesh in CDC class I/high-risk ventral and incisional hernia repair: 18-month follow-up. *Surg Endosc* 2018;32(04):1929–1936
 - 99 Sajid MS, Bokhari SA, Mallick AS, Cheek E, Baig MK. Laparoscopic versus open repair of incisional/ventral hernia: a meta-analysis. *Am J Surg* 2009;197(01):64–72
 - 100 Sandor M, Xu H, Connor J, et al. Host response to implanted porcine-derived biologic materials in a primate model of abdominal wall repair. *Tissue Eng Part A* 2008;14(12):2021–2031

- 101 Sarikaya A, Record R, Wu CC, Tullius B, Badylak S, Ladisch M. Antimicrobial activity associated with extracellular matrices. *Tissue Eng* 2002;8(01):63–71
- 102 Saulis AS, Dumanian GA. Periumbilical rectus abdominis perforator preservation significantly reduces superficial wound complications in “separation of parts” hernia repairs. *Plast Reconstr Surg* 2002;109(07):2275–2280, discussion 2281–2282
- 103 Schreinemacher MH, Emans PJ, Gijbels MJ, Greve JW, Beets GL, Bouvy ND. Degradation of mesh coatings and intraperitoneal adhesion formation in an experimental model. *Br J Surg* 2009;96(03):305–313
- 104 Schug-Pass C, Sommerer F, Tannapfel A, Lippert H, Köckerling F. The use of composite meshes in laparoscopic repair of abdominal wall hernias: are there differences in biocompatibility?: experimental results obtained in a laparoscopic porcine model *Surg Endosc* 2009;23(03):487–495
- 105 Schumpelick V, Klinge U, Junge K, Stumpf M. Incisional abdominal hernia: the open mesh repair. *Langenbecks Arch Surg* 2004;389(01):1–5
- 106 Shekelle PG, Woolf SH, Eccles M, Grimshaw J. Developing clinical guidelines. *West J Med* 1999;170(06):348–351
- 107 Shekelle PG, Woolf SH, Eccles M, Grimshaw J. Clinical guidelines: developing guidelines. *BMJ* 1999;318(7183):593–596
- 108 Smart NJ, Bloor S. Durability of biologic implants for use in hernia repair: a review. *Surg Innov* 2012;19(03):221–229
- 109 Stone HH, Fabian TC, Turkleson ML, Jurkiewicz MJ. Management of acute full-thickness losses of the abdominal wall. *Ann Surg* 1981;193(05):612–618
- 110 Stoppa RE. The treatment of complicated groin and incisional hernias. *World J Surg* 1989;13(05):545–554
- 111 Switzer NJ, Dykstra MA, Gill RS, et al. Endoscopic versus open component separation: systematic review and meta-analysis. *Surg Endosc* 2015;29(04):787–795
- 112 Szczerba SR, Dumanian GA. Definitive surgical treatment of infected or exposed ventral hernia mesh. *Ann Surg* 2003;237(03):437–441
- 113 Tagaya N, Mikami H, Aoki H, Kubota K. Long-term complications of laparoscopic ventral and incisional hernia repair. *Surg Laparosc Endosc Percutan Tech* 2004;14(01):5–8
- 114 Temudom T, Siadati M, Sarr MG. Repair of complex giant or recurrent ventral hernias by using tension-free intraparietal prosthetic mesh (Stoppa technique): lessons learned from our initial experience (fifty patients). *Surgery* 1996;120(04):738–743, discussion 743–744
- 115 Tsimoyiannis EC, Tsimogiannis KE, Pappas-Gogos G, Nikas K, Karfis E, Sioziou H. Seroma and recurrence in laparoscopic ventral hernioplasty. *JSLs* 2008;12(01):51–57
- 116 Turner PL, Park AE. Laparoscopic repair of ventral incisional hernias: pros and cons. *Surg Clin North Am* 2008;88(01):85–100, viii–viii.
- 117 van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001;345(19):1359–1367
- 118 van ’t Riet M, de Vos van Steenwijk PJ, Bonthuis F, et al. Prevention of adhesion to prosthetic mesh: comparison of different barriers using an incisional hernia model. *Ann Surg* 2003;237(01):123–128
- 119 van’t Riet M, de Vos van Steenwijk PJ, Bonjer HJ, Steyerberg EW, Jeekel J. Mesh repair for postoperative wound dehiscence in the presence of infection: is absorbable mesh safer than non-absorbable mesh? *Hernia* 2007;11(05):409–413
- 120 Vargo D. Component separation in the management of the difficult abdominal wall. *Am J Surg* 2004;188(06):633–637
- 121 Breuing K, Butler CE, Ferzoco S, et al; Ventral Hernia Working Group. Incisional ventral hernias: review of the literature and recommendations regarding the grading and technique of repair. *Surgery* 2010;148(03):544–558
- 122 Voyles CR, Richardson JD, Bland KI, Tobin GR, Flint LM, Polk HC Jr. Emergency abdominal wall reconstruction with polypropylene mesh: short-term benefits versus long-term complications. *Ann Surg* 1981;194(02):219–223
- 123 Wassenaar EB, Raymakers JT, Rakic S. Impact of the mesh fixation technique on operation time in laparoscopic repair of ventral hernias. *Hernia* 2008;12(01):23–25
- 124 White TJ, Santos MC, Thompson JS. Factors affecting wound complications in repair of ventral hernias. *Am Surg* 1998;64(03):276–280
- 125 Xu H, Wan H, Sandor M, et al. Host response to human acellular dermal matrix transplantation in a primate model of abdominal wall repair. *Tissue Eng Part A* 2008;14(12):2009–2019