Published in final edited form as:

ACS Chem Biol. 2018 September 21; 13(9): 2655–2662. doi:10.1021/acscchembio.8b00577.

Structure-function analyses of the *N*-butanoyl L-homoserine lactone quorum-sensing signal define features critical to activity in RhIR

Michelle E. Boursier^{1,†}, Joseph D. Moore^{1,‡}, Katherine M. Heitman¹, Sally P. Shepardson-Fungairino^{1,§}, Joshua B. Combs^{1,**}, Lea C. Koenig^{1,††}, Daniel Shin², Eric C. Brown², Rajesh Nagarajan², and Helen E. Blackwell¹

¹Department of Chemistry, University of Wisconsin–Madison, 1101 University Ave., Madison, WI 53706, USA;

²Department of Chemistry and Biochemistry, Boise State University, 1910 University Dr., Boise, ID 83725, USA; blackwell@chem.wisc.edu

Abstract

Pseudomonas aeruginosa is an opportunistic pathogen that coordinates the production of many virulence phenotypes at high population density via quorum sensing (QS). The LuxR-type receptor RhIR plays an important role in the *P. aeruginosa* QS process, and there is considerable interest in the development of chemical approaches to modulate the activity of this protein. RhIR is activated by a simple, low molecular weight *N*-acyl L-homoserine lactone signal, *N*-butanoyl-L-homoserine lactone (BHL). Despite the emerging prominence of RhIR in QS pathways, there has been limited exploration of the chemical features of the BHL scaffold that are critical to its function. In the current study, we sought to systematically delineate the structure-activity relationships (SARs) driving BHL activity for the first time. A focused library of BHL analogs was designed, synthesized, and evaluated in cell-based reporter gene assays for RhIR agonism and antagonism. These investigations allowed us to define a series of SARs for BHL-type ligands and identify structural motifs critical for both activation and inhibition of the RhIR receptor. Notably, we identified agonists that have ~10-fold higher potencies in RhIR relative to BHL, are highly selective for RhIR agonism over LasR, and are active in the *P. aeruginosa* background. These compounds and the SARs reported herein should pave a route toward new chemical strategies to study RhIR in *P. aeruginosa*.

GRAPHICAL ABSTRACT

[†]Current address: Promega Corporation, Madison, WI 53711

[‡]Current address: Dow Chemical Company, Collegeville, PA 19426

[§]Current address: Department of Chemistry, Tufts University, Medford, MA 02155

^{**}Current address: Department of Chemistry, Yale University, New Haven, CT 06520

^{††}Current address: Covance Laboratories, Madison, WI 53704

Electronic supplementary information (ESI) available

Details of instrumentation, reagents, synthetic methods, primary screening data, dose-response curves, and characterization data for all new compounds.

INTRODUCTION

Certain bacteria are capable of coordinating population density with gene expression using an intercellular chemical signaling process known as quorum sensing (QS) (1, 2). QS allows bacteria to synchronize group-beneficial phenotypes only at high populations. In Gram-negative bacteria, QS is typically mediated by *N*-acyl L-homoserine lactone (AHL) signals, which are produced by LuxI-type synthases (3). These small molecules can freely diffuse out of the cell (though in select cases, export is facilitated by efflux pumps (4, 5)) and into the environment or other neighboring cells; as population density increases, the AHLs will reach an intracellular concentration at which they productively bind LuxR-type receptors. The ligand-bound LuxR-type receptors then act as transcription factors and alter gene expression levels to regulate a broad diversity of collective behaviors, including motility, biofilm formation, antibiotic and virulence factor production, and bioluminescence (6). As some of the most common agents of human infection use QS to control virulence, their QS systems have become attractive targets for infection control (7, 8). More fundamentally, identifying chemical strategies to modulate QS can provide novel entry into the molecular mechanisms of this important signaling pathway. Our laboratory and others have focused intently on developing such chemical approaches to attenuate QS in bacterial pathogens and symbionts over the past decade (9).

The Gram-negative bacterium *Pseudomonas aeruginosa* is one of the most common causes of hospital-acquired infections, and uses QS to control ~10% of its genome, much of which is involved in virulence (10). *P. aeruginosa* utilizes a relatively complex network of receptors and chemical signals to control QS, including multiple LuxI/LuxR pairs (Figure 1). The two LuxI-type synthases, LasI and RhlI, produce *N*-(3-oxo)-dodecanoyl HL (OdDHL; Figure 1) and *N*-butanoyl HL (BHL; Figure 1), respectively (11). These two signaling molecules are recognized by their cognate LuxR-type receptors, LasR and RhIR. Both QS circuits regulate a large number of virulence factors—for example, the LasI/R system regulates the production of elastase B, alkaline protease, and exotoxin A; and the RhlI/R system regulates rhamnolipid production (a rhamnose-based biosurfactant) and the toxic exofactors hydrogen cyanide and pyocyanin (3, 11). Interestingly, OdDHL is also recognized by the orphan LuxR-type receptor, QscR, which lacks its own cognate ligand. Once ligand bound, QscR has been found to both negatively regulate LasR and activate its own unique regulon in *P. aeruginosa* (12). In addition to the Las and Rhl circuits, *P. aeruginosa* has a third circuit, Pqs, which is regulated by the lysR-type receptor PqsR (unrelated to LuxR-type receptors), the

Pseudomonas quinolone signal (PQS) and the PQS biosynthetic precursor 4-hydroxy-2-heptylquinoline (HHQ), and plays a role in the regulation of pyocyanin production (13). Adding to the complexity of QS in *P. aeruginosa*, each of the QS systems can positively or negatively regulate the other QS systems, and this interregulation is exquisitely sensitive to environmental factors, allowing for nimble and intricate genome regulation (14).

LasR is generally considered to be at the top of the *P. aeruginosa* QS receptor hierarchy, as it regulates genes associated with both the *rhl* and *pqs* circuits (Figure 1) (15). Due to this prominent role, it's not surprising that LasR has been a primary target for the design of small molecule antagonists to block QS in this pathogen (9, 16–18). Far fewer research efforts have been directed toward the design of non-native ligands for RhlR, likely due to its perceived secondary role in QS. However, a growing number of reports have shown that both the nutrient conditions and stage of bacterial growth can reroute the QS regulatory circuitry so that RhlR instead is dominant, upending the traditional understanding of QS receptor regulation (14, 19–22). Experiments by the Bassler laboratory have further implicated the *rhl* system in the direct regulation of virulence factors and subsequent reduction of virulence factor production using multiple in vivo models (23, 24). In synergy with these reports, our laboratory has recently shown that inhibition of pyocyanin production can be achieved via agonism of RhlR using small molecule ligands (25).

Together, these prior results highlight that the RhlR receptor presents a significantly underdeveloped opportunity for the attenuation of *P. aeruginosa* virulence. To date, only a few studies have specifically focused on the development of synthetic modulators of RhlR. In 2003, Suga and co-workers showed that BHL analogs with cyclopentanone or cyclohexanone head groups and butanoyl tails exhibit potent agonistic activity towards RhlR in a *P. aeruginosa* reporter system (26, 27). A decade later, Bassler and co-workers reported that a *meta*-bromo aryl homocysteine thiolactone AHL mimic (**mBTL**) strongly modulated the *rhl* system, decreasing pyocyanin production and associated virulence (23). In 2015, we reported the screening of our in-house non-native AHL libraries for RhlR modulators and the discovery of a small collection of new RhlR ligands (28). Perhaps not surprisingly, the compounds that were capable of potent RhlR activation (e.g., **D8** and **S4**; Figure 2) were similar in structure to its native ligand BHL. However, to date, no systematic investigation of the structure-activity relationships (SARs) dictating BHL function have been reported. Delineation of these SARs would not only allow for a deeper fundamental understanding of the features critical to BHL activity, but also provide a framework from which to design new ligands for RhlR with improved potencies.

Herein, we report the design, synthesis, and biological characterization of a set of BHL analog libraries with the intent of discerning SAR trends important for agonism and antagonism of the RhlR receptor in *P. aeruginosa*. These studies revealed the first set of detailed SAR surrounding the BHL scaffold and some of the most potent and receptor-selective non-native RhlR agonists to be reported.

RESULTS AND DISCUSSION

RhlR-targeted library design.

An overview of our library design process and the structures of the BHL-derived library are shown in Figure 2. Our library constitutes largely new compounds. Of the subset of these compounds that have been reported previously (for references, see Table S1), none to our knowledge have been tested for direct RhlR modulation in a heterologous background (e.g., *E. coli*) lacking other competing LuxR-type receptors; previous screening efforts have utilized *P. aeruginosa* reporter strains that contain the organism's complete and inter-connected QS network. The BHL-derived library can be divided into two groups: (1) compounds with modified tails and native lactone head groups (i.e., the tail library), and (2) compounds with modified head groups and native butanoyl tails (i.e., the head library). The tail library compounds can be further divided into three subsets based on structural similarity to either BHL or our two previously reported potent RhlR activators **2** and **3** (28). The first subset of compounds (**4–11**) retains the short-chain alkyl characteristics of native ligand BHL, with compounds designed to probe both tail length and alkyl substitution on the tail. We note that compound **7** (isovaleryl HL) is actually a naturally occurring AHL, first found in the symbiont *Bradyrhizobium japonicum* (29). 2-methylbutanoyl HL (**8**) was examined as a diastereomeric mixture. The second subset of tail-modified compounds (**12–16**) expands on the structural characteristics of lead pentenyl HL **2**, probing alkene stereochemistry and position, as well as the presence of an alkene vs. an alkyne in the tail. Lastly, the third subset of tail modified compounds (**17–22**) was designed to explore structural features of lead cyclopentyl HL **3**, focusing on size and the position of small carbocycles in the acyl tail.

The head group library was similarly divided into four subsets of compounds (Figure 2). While previous studies of AHL-type ligands have explored alternate head groups expanded beyond 5-membered rings (27, 30), we elected in the current study to focus exclusively on close mimics of the homoserine lactone to hopefully delineate more direct SAR for the BHL ligand. The first subset of head-group-modified compounds investigates alternate replacements for the amide linkage in the native ligand (sulfonamide **23** and ester **24**). The second, one compound “sub-set” was designed to examine the importance of native stereochemistry for BHL activity (i.e., D-BHL **25**). The third subset incorporates one-atom modifications to the lactone ring, replacing the lactone oxygen with sulfur (**26**), nitrogen (**27**), or a methylene (**28**). The fourth subset deconstructs the five-membered lactone ring even further by removing the carbonyl (**29**), replacing it with an alcohol (**31**), or varying the heteroatom location in the ring (**30**, **32**, and **33**). We note that compounds **32** and **33** (both examined as racemates) also contain an additional methylene between the ring and the amide bond; this feature was included simply due to ease of synthesis. We reasoned that these last two compounds could allow us to also explore the effect of increasing the distance the head group extends into the RhlR ligand binding pocket (assuming these close BHL analogs target the same site).

Library synthesis.

AHL library compounds were synthesized using previously established solution-phase chemistry, mainly based on carbodiimide-mediated amide coupling procedures (see Methods) (31).

E. coli reporter screens reveal BHL SARs and new RhIR agonists and antagonists.

We first evaluated the RhIR-modulatory activity of the library compounds in an *E. coli* reporter strain (JLD271 harboring a b-galactosidase reporter for RhIR; see Methods) in order to isolate the RhIR receptor from other QS regulators in *P. aeruginosa* (most notably, LasR) and thus study its response to ligands directly. We performed initial RhIR agonism and competitive RhIR antagonism screens of the library at 10 μ M. Many compounds were RhIR agonists at this concentration; however, no compounds were able to inhibit RhIR activity at 10 μ M (Table S2). We thus submitted all compounds to a second competitive antagonism screen at 1 mM against 10 μ M BHL.

Strikingly, the primary agonism screen revealed multiple compounds with efficacies (reported as % activation, Table 1) greater than RhIR's native ligand BHL (**1**) at 10 μ M, which is BHL's approximate EC₅₀ value. The most active compounds were isovaleryl HL **7** and cyclobutyl HL **17** (86% and 79% activation, respectively), each capable of agonizing RhIR at levels equal to or greater than our previous lead RhIR agonist, **3** (**S4**). Scrutinizing the agonism data in terms of structural features critical for ligand activity, the most influential feature appeared to be substitution of the acyl tail α -carbon. Compounds with tertiary substituents at the α - or β -carbons appear to be well tolerated by RhIR, yet quaternary carbons abolished agonistic activity. For example, compounds **5–8** show modest to strong RhIR agonism activity (25–86%), whereas **9** and **10** were inactive as agonists. These data suggest that a specific handedness may be preferred at the α -carbon, and accordingly, that one of the diastereomers of 2-methylbutanoyl (**8**) is likely more active. Similarly, cyclic-tail compounds **17** and **19** were strong agonists (70% activation), but 1-methyl-cyclopropanoyl HL **20** was inactive likely due to its quaternary α -carbon. We also observed that AHLs with 3-carbon tails (e.g., propyl HL **4** and cyclopropyl HL **18**) display significantly reduced abilities to activate RhIR relative to AHL homologs with straight chain or cyclic tails expanded by just one methylene (e.g., BHL (**1**), **17** and **19**).

Amongst the set of compounds tested, the presence of an alkene or alkyne in the acyl tail had relatively little effect on RhIR agonistic activity (Table 1). Butenyl HL **14** and butynyl HL **15** were the most active of all compounds with unsaturated acyl tails: both displayed agonistic activities comparable BHL (**1**). Alkene isomers displayed different activity profiles; *trans* 2-butenyl derivative **13** exhibited a two-fold reduction in activity from its *cis* isomer **14**. The “kinked” *cis* alkene may enforce a tail conformation permitting **14** to bind RhIR in a manner similar to the active lead cyclopentyl HL **3**. The decrease in potency observed in the longer straight chain (6 carbon) alkenyl and alkynyl HLs **12** and **16** is consistent with the prior observation that AHLs with longer acyl chains are less active RhIR agonists relative to shorter chain AHLs (26, 30).

Of the cyclic-tail AHLs tested, cyclobutanoyl HL **17** displayed the strongest RhIR-agonistic activity (79% activation; Table 1). While compound **18**, with a cyclopropanoyl tail alone, fails to activate RhIR strongly (32% agonism), agonism was recovered in the one-carbon extended homolog **19** (70% activation). The larger cyclohexyl tail of **21** yielded reduced activity relative to cyclopentyl HL **3** (34% vs. 76%, respectively). Phenyl compound **22** had no agonist activity, suggesting that the tail's rigidity at the α -carbon may be detrimental to RhIR binding.

Turning to the head-group modified BHL analogs, AHLs **26** and **28** were the only compounds to show appreciable RhIR agonism at 10 μ M (Table 1). Both of these compounds maintain a carbonyl in the head group, and the more active homocysteine thiolactone analog **26** (71% activation) maintains the heteroatom electron donor. Most other head group alterations shut down all agonistic activity in RhIR at 10 μ M, including replacement of the ketone in **28** to give alcohol **31** (supporting the Suga lab's earlier result (26, 27)), replacing the native lactone with an amide in lactam **27**, removing the lactone carbonyl to give cycloether **30**, or extending the head group motif by an additional methylene as in compounds **32** and **33**. The D-stereoisomer of BHL (**25**) showed marginal activity even at 1 mM (Table S2), confirming previously reported data that the stereochemistry of BHL is critical for RhIR activation (32). The AHL amide proved to be necessary for agonistic activity, as compound **24**, the ester analog of BHL, was inactive. This result for BHL correlates with prior reports indicating that the AHL amide NH is essential for AHL:LuxR-type receptor binding, making a key hydrogen bond with a conserved aspartic acid in the ligand-binding site (33).

Turning next to the RhIR antagonism assay data for the tail group library (Table 1), only compounds **9**, **20**, and **22** were capable of inhibiting RhIR activity to a statistically significant extent (25–31%) at 1 mM. These compounds contain either a quaternary or sp^2 hybridized α -carbon in their tails, suggestive that steric bulk in close proximity to the HL engenders weak antagonistic activity for short-tail AHLs. More head group library members displayed RhIR antagonism, with compounds **23**, **27**, **29**, and **33** all showing antagonism greater than 35%. Sulfonamide **23**, a 55% antagonist, has increased steric bulk adjacent to the amide NH and potentially operates through a similar mode of action as bulky tail antagonists **9** and **20**. In turn, cyclolactam **27** and cyclopentyl derivative **29** lack key hydrogen bonding contacts presented by the native lactone HL for agonism (see above), and antagonize RhIR instead. Finally, the extended, tetrahydrofurfurylamine head group of **33** caused moderate RhIR antagonism (57% at 1 mM), suggesting that this positioning of the head group enforces inhibitory interactions.

BHL-type analogs can antagonize, but not agonize LasR.

We were also interested to examine the activity of the BHL library in LasR, as ligands selective for RhIR, or displaying opposite activities in each receptor (i.e., both a RhIR agonist and a LasR antagonist, and vice versa) would be useful as probes for teasing apart the closely interrelated QS circuit in *P. aeruginosa* (Figure 1). We thus tested the library in an *E. coli* LasR reporter strain using analogous methods to that for RhIR above (see Methods).

In general, our lead RhlR agonists showed negligible LasR agonism in the *E. coli* reporter, particularly at 10 μ M (Table S3). This result is not surprising, in view of the structural differences between the native ligands for these two receptors (short chain BHL in RhlR versus long chain OdDHL in LasR). However, we found that many of the RhlR agonists, including the native ligand BHL (**1**), *antagonize* the LasR receptor instead at high concentrations (1 mM, Table S3). To our knowledge, LasR antagonism by RhlR *agonists* (including BHL) has not been previously reported explicitly. Interestingly, the potent RhlR agonists **7** and **17** are also some of the most active LasR antagonists found in the library (>70% inhibition). This activity profile for **7** and **17** is unique, as such compounds could be utilized to suppress LasR-associated virulence factors along with those suppressed by RhlR agonism (i.e., PQS) (25). That said, the high compound concentrations necessary for this observed activity might reduce the likelihood of implementing such dual regulation in the wild type organism.

Dose-response studies reveal highly potent RhlR agonists.

To gauge the relative potencies of the lead RhlR agonists identified in the primary assays, we submitted these compounds to dose-response analyses using the *E. coli* RhlR reporter. All tail group compounds that showed activities comparable to or greater than that of BHL (**1**), along with two head group compounds (**26** and **28**) with promising activities, were evaluated (Table 2). Four of the tail library members tested were found to be significantly more potent than BHL. Cyclobutyl HL **17** and isovaleryl HL **7** rivaled in potency that of cyclopentyl HL **3**. Naturally occurring isovaleryl HL was the most potent activator of RhlR in this *E. coli* bioassay overall, with an EC₅₀ almost 10-fold lower than BHL (1 mM vs. ~9 μ M, respectively).

Of the head group library compounds tested (**26** and **28**), the homocysteine thiolactone BHL analog **26** was found to have comparable potency to BHL (Table 2). While thiolactone analogs of other AHLs have been reported to have comparable or lower agonistic activity than their parent AHL (34, 35), **26** represents to our knowledge the first thiolactone analog that is more potent than the native AHL ligand. We do note that LuxR-type receptor SdiA in *Salmonella enterica* serovar Typhimurium has been reported to be more strongly activated by ligands bearing thiolactone relative to lactone head groups, but this receptor is an “orphan” type LuxR receptor that lacks a native ligand for direct comparison (36). Turning to cyclopentanone derivative **28**, this analog was found to be just under 3-fold less active than BHL ($p < 0.05$), tracking with early reports by Suga and co-workers in *P. aeruginosa* (27). Despite its reduced potency, the ketone head group in **28** is not susceptible to hydrolysis (and its racemization rate should be slow (37)), marking it as a potentially useful moiety for use in the design of future, hydrolytically stable AHL analogs.

Library compounds capable of greater than 25% antagonistic activity against RhlR (**23**, **27**, **29**, and **33**) were also submitted to dose-response analysis in the *E. coli* RhlR reporter strain versus 10 μ M BHL. Overall, this set of compounds showed limited potencies (IC₅₀ > 60 μ M, dose-response curves shown in Figure S3); only the dose-response curve for cyclopentyl head group derivative **29** yielded a calculable IC₅₀ value (52.2 μ M). As this library was

designed to very closely mimic the BHL scaffold, it is not surprising that we mostly identified agonists as opposed to antagonists.

Lead RhlR agonists maintain their potencies in *P. aeruginosa* background.

We next performed dose-response analyses on the lead RhlR agonists in *P. aeruginosa* to determine if they maintained their potencies in the native organism. For these experiments, we used the *P. aeruginosa* double synthase mutant PAO-JP2 (*lasI rhII*) harboring the RhlR reporter plasmid *prhII-LVAgfp* (see Methods). Because the production of RhlR is dependent on LasR in *P. aeruginosa* (at least in LB medium (14)), all assays were performed in the presence of 100 nM OdDHL.

We were pleased to observe that all of the lead RhlR agonists retained similar potencies in the *P. aeruginosa* background relative to the *E. coli* RhlR reporter (Table 2). Notably, isovaleryl HL **7** and cyclobutyl HL **17** (alongside compound **3**) represent, to our knowledge, the most potent non-native agonists of RhlR in *P. aeruginosa*. The homocysteine thiolactone BHL analog **26** showed a slight but significant improvement in potency over BHL as compared to the *E. coli* reporter data ($p < 0.05$), making it the only headgroup to prove more potent than the native ligand. As the lead RhlR agonists only display appreciable LasR antagonism at low millimolar concentrations (Table S3), the observed activities of lead compounds on RhlR in *P. aeruginosa* are likely not a result of indirect modulation of RhlR via LasR, but rather are due to interaction with RhlR. We note that that we have made the assumption throughout this study that these very close BHL analogs target the BHL-binding site in RhlR and engender ligand:receptor interactions that alter RhlR function. However, we cannot discount the possibility that they also may target an allosteric site on RhlR (as either agonist or antagonists). Additional biochemical studies are required to more fully characterize their specific interactions with RhlR and are ongoing.

SUMMARY

The goal of this study was to delineate key SARs dictating the activity of the natural *P. aeruginosa* QS signal, BHL, in the RhlR receptor. We addressed this goal via the design, synthesis, and biological characterization of a focused library of non-native AHL analogs closely based on the BHL scaffold. These experiments revealed the following SARs critical to BHL activity (Figure 3): first, the head group should maintain L-stereochemistry as well as a carbonyl adjacent to the stereocenter, and there should be an amide linkage between the head and tail groups. These requirements corroborate past studies indicating that the AHL head group carbonyl and amide make key hydrogen-bonding contacts in LuxR-type receptors with well-conserved residues (38). Second, the head group can accommodate multiple heteroatom changes alpha to the head group carbonyl, but proton donors such as an amide are not tolerated. Third, the AHL tail must contain four atoms; longer or shorter tails result in reduced agonism. The tail also must have a secondary or tertiary carbon alpha to the amide linker. We also found changes resulting in compounds with improved agonism activity over BHL. These agonists have added steric bulk relative to BHL, specifically homocysteine thiolactone head groups and cyclobutyl, cyclopentyl, or isovaleryl moieties for tail groups. Interestingly, removing or altering many of the aforementioned critical

components of BHL for RhIR activation resulted in RhIR antagonists. Only the removal of the amide nitrogen resulted in compounds that had no RhIR activity. Future RhIR antagonist libraries may benefit from the removal of activating components determined in this study.

Finally, in the course of these experiments, we identified several compounds capable of fully agonizing RhIR with higher potencies than the native ligand BHL, both with and without modified head groups. Notably, isovaleryl HL **7** and cyclobutyl HL **17** are almost 10-fold more potent than BHL, active in *P. aeruginosa*, and selective for RhIR agonism. Agonism (or partial agonism) of the RhIR system has already been implicated as a possible anti-virulence strategy (23–25). As such, the structural insights and new RhIR ligands reported here will be useful in advancing new chemical probe development for further study of this important QS receptor.

METHODS

Chemistry.

Most compounds were synthesized using previously established solution-phase, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC)-mediated amide coupling procedures (31), with a few modifications (see Supporting Information for full details).

General bacterial growth conditions and assay methods.

Bacteria were cultured in Luria–Bertani broth (LB) at 37 °C with shaking at 200 rpm. Compound stock solutions (100 mM) were prepared in DMSO. Absorbance measurements were performed using a Biotek Synergy 2 plate reader and Gen 5 software (version 1.05). Bacterial growth was quantified according to absorbance at 600 nm (OD₆₀₀). All biological assay data were processed using Microsoft Excel and GraphPad Prism 6 software.

Bacterial strains.

The following reporter strains were used: (i) *E. coli* strain JLD271 (*sdhA*) (39) harboring the RhIR expression plasmid pJN105R2 (25) and the *rhII-lacZ* transcriptional fusion reporter pSC11-rhII* (25), (ii) *E. coli* strain JLD271 (*sdhA*) (39) harboring the LasR expression plasmid pJN105L (40) and the *lasI-lacZ* transcriptional fusion reporter pSC11 (12), and (iii) *P. aeruginosa* strain PAO-JP2 (*lasI rhII*) (41) harboring the *rhII-gfp* transcriptional fusion reporter *prhII-LVAgfp* (42). Reporter strains JLD271/pJN105R2/pSC11-rhII* and JLD271/pJN105L/pSC11 were grown in LB containing 100 µg mL⁻¹ ampicillin and 10 µg mL⁻¹ gentamicin.

E. coli RhIR and LasR reporter assay protocols.

The *E. coli* reporters JLD271/pJN105R2/pSC11-rhII* and JLD271/pJN105L/pSC11 were used as previously described (28), with the following modifications: in the RhIR agonism reporter assay performed with substrate chlorophenol red-β-D-galactopyranoside (CPRG), non-native compounds (at 10 µM or 1 mM) were compared to 1 mM BHL as the positive control; LasR reporter assays were processed using our reported protocol for the substrate

ortho-nitrophenyl- β -galactoside (ONPG) (43); and LasR antagonism assays were performed versus the EC₅₀ for OdDHL (2 nM).

***P. aeruginosa* RhlR reporter assay protocols.**

The *P. aeruginosa* PAO-JP2 reporter strain harboring *prhII-LVAgfp* was used as previously reported (25), with the following modifications: in the RhlR agonism assay, non-native compounds were compared to 1 mM BHL as a positive control; and RhlR antagonism assays were performed versus the EC₅₀ for BHL (10 μ M).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Financial support for H.E.B was provided by the NIH (R01 GM109403). Financial support for R.N. was provided by Boise State University start-up funds and the NIH (R15 GM117323, P20 RR016454, and P20 GM103408). M.E.B. was funded in part by the NSF through the UW–Madison Materials Research Science and Engineering Center (DMR-1121288). J.D.M was supported in part by the UW–Madison NIH Biotechnology Training Program (T32 GM08349). NMR facilities in the UW–Madison Department of Chemistry were supported by the NSF (CHE-0342998) and a gift from P. Bender. MS facilities in the Departments of Chemistry at UW–Madison and Boise State University were supported by the NSF (CHE-9974839 and CHE-0923535, respectively).

REFERENCES

1. Whiteley M, Diggle SP, and Greenberg EP (2017) Progress in and promise of bacterial quorum sensing research, *Nature* 551, 313–320. [PubMed: 29144467]
2. Camilli A, and Bassler BL (2006) Bacterial Small-Molecule Signaling Pathways, *Science* 311, 1113–1116. [PubMed: 16497924]
3. Schuster M, Sexton DJ, Diggle SP, and Greenberg EP (2013) Acyl-homoserine lactone quorum sensing: from evolution to application, *Annu. Rev. Microbiol* 67, 43–63. [PubMed: 23682605]
4. Pearson JP, Van Delden C, and Iglewski BH (1999) Active Efflux and Diffusion Are Involved in Transport of *Pseudomonas aeruginosa* Cell-to-Cell Signals, *J. Bacteriol* 181, 1203–1210. [PubMed: 9973347]
5. Moore JD, Gerdt JP, Eibergen NR, and Blackwell HE (2014) Active efflux influences the potency of quorum sensing inhibitors in *Pseudomonas aeruginosa*, *ChemBioChem* 15, 435–442. [PubMed: 24478193]
6. Rutherford ST, and Bassler BL (2012) Bacterial quorum sensing: its role in virulence and possibilities for its control, *Cold Spring Harb. Perspect. Med* 2, a012427. [PubMed: 23125205]
7. O'Connell KM, Hodgkinson JT, Sore HF, Welch M, Salmond GP, and Spring DR (2013) Combating Multidrug-Resistant Bacteria: Current Strategies for the Discovery of Novel Antibacterials, *Angew. Chem. Int. Ed* 52, 10706–10733.
8. Clatworthy AE, Pierson E, and Hung DT (2007) Targeting virulence: a new paradigm for antimicrobial therapy, *Nat. Chem. Biol* 3, 541–548. [PubMed: 17710100]
9. Welsh MA, and Blackwell HE (2016) Chemical probes of quorum sensing: from compound development to biological discovery, *FEMS Microbiol. Rev* 40, 774–794. [PubMed: 27268906]
10. Strateva T, and Mitov I (2011) Contribution of an arsenal of virulence factors to pathogenesis of *Pseudomonas aeruginosa* infections, *Ann. Microbiol* 61, 717–732.
11. Schuster M, and Greenberg EP (2008) LuxR-Type Proteins in *Pseudomonas aeruginosa* Quorum Sensing: Distinct Mechanisms with Global Implications, In *Chemical Communication among Bacteria* (Winans SC, and Bassler BL, Eds.), pp 133–144, ASM Press, Washington, DC.

12. Chugani S. a., Whiteley M, Lee KM, D'Argenio D, Manoil C, and Greenberg EP (2001) QscR, a modulator of quorum-sensing signal synthesis and virulence in *Pseudomonas aeruginosa*, *Proc. Nat. Acad. Sci. USA* 98, 2752–2757. [PubMed: 11226312]
13. Deziel E, Lepine F, Milot S, He JX, Mindrinos MN, Tompkins RG, and Rahme LG (2004) Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication, *Proc. Nat. Acad. Sci. USA* 101, 1339–1344. [PubMed: 14739337]
14. Welsh MA, and Blackwell HE (2016) Chemical Genetics Reveals Environment-Specific Roles for Quorum Sensing Circuits in *Pseudomonas aeruginosa*, *Cell Chem. Biol* 23, 361–369. [PubMed: 26905657]
15. Schuster M, Lostroh CP, Ogi T, and Greenberg EP (2003) Identification, Timing, and Signal Specificity of *Pseudomonas aeruginosa* Quorum-Controlled Genes: a Transcriptome Analysis, *J. Bacteriol* 185, 2066–2079. [PubMed: 12644476]
16. Geske GD, O'Neill JC, and Blackwell HE (2008) Expanding dialogues: from natural autoinducers to non-natural analogues that modulate quorum sensing in Gram-negative bacteria, *Chem. Soc. Rev* 37, 1432–1447. [PubMed: 18568169]
17. Galloway W, Hodgkinson JT, Bowden SD, Welch M, and Spring DR (2011) Quorum Sensing in Gram-Negative Bacteria: Small-Molecule Modulation of AHL and AI-2 Quorum Sensing Pathways, *Chem. Rev* 111, 28–67. [PubMed: 21182299]
18. Moore JD, Rossi FM, Welsh MA, Nyffeler KE, and Blackwell HE (2015) A Comparative Analysis of Synthetic Quorum Sensing Modulators in *Pseudomonas aeruginosa*: New Insights into Mechanism, Active Efflux Susceptibility, Phenotypic Response, and Next-Generation Ligand Design, *J. Am. Chem. Soc* 137, 14626–14639. [PubMed: 26491787]
19. Van Delden C, Pesci EC, Pearson JP, and Iglewski BH (1998) Starvation Selection Restores Elastase and Rhamnolipid Production in a *Pseudomonas aeruginosa* Quorum-Sensing Mutant, *Infect. Immun* 66, 4499–4502. [PubMed: 9712807]
20. Dekimpe V, and Déziel E (2009) Revisiting the quorum-sensing hierarchy in *Pseudomonas aeruginosa*: the transcriptional regulator RhlR regulates LasR-specific factors, *Microbiology* 155, 712–723. [PubMed: 19246742]
21. Bjarnsholt T, Jensen PO, Jakobsen TH, Phipps R, Nielsen AK, Rybtke MT, Tolker-Nielsen T, Givskov M, Hoiby N, and Ciofu O (2010) Quorum sensing and virulence of *Pseudomonas aeruginosa* during lung infection of cystic fibrosis patients, *PLoS ONE* 5, e10115. [PubMed: 20404933]
22. Folkesson A, Jelsbak L, Yang L, Johansen HK, Ciofu O, Høiby N, and Molin S (2012) Adaptation of *Pseudomonas aeruginosa* to the cystic fibrosis airway: an evolutionary perspective, *Nat. Rev. Microbiol* 10, 841–851. [PubMed: 23147702]
23. O'Loughlin CT, Miller LC, Siryaporn A, Drescher K, Semmelhack MF, and Bassler BL (2013) A quorum-sensing inhibitor blocks *Pseudomonas aeruginosa* virulence and biofilm formation, *Proc. Nat. Acad. Sci. USA* 110, 17981–17986. [PubMed: 24143808]
24. Mukherjee S, Moustafa D, Smith CD, Goldberg JB, and Bassler BL (2017) The RhlR quorum-sensing receptor controls *Pseudomonas aeruginosa* pathogenesis and biofilm development independently of its canonical homoserine lactone autoinducer, *PLoS Pathog.* 13, e1006504. [PubMed: 28715477]
25. Welsh MA, Eibergen NR, Moore JD, and Blackwell HE (2015) Small molecule disruption of quorum sensing cross-regulation in *Pseudomonas aeruginosa* causes major and unexpected alterations to virulence phenotypes, *J. Am. Chem. Soc* 137, 1510–1519. [PubMed: 25574853]
26. Smith KM, Bu Y, and Suga H (2003) Induction and Inhibition of *Pseudomonas aeruginosa* Quorum Sensing by Synthetic Autoinducer Analogs, *Chem. Biol* 10, 81–89. [PubMed: 12573701]
27. Jog GJ, Igarashi J, and Suga H (2006) Stereoisomers of *P. aeruginosa* autoinducer analog to probe the regulator binding site, *Chem. Biol* 13, 123–128. [PubMed: 16492560]
28. Eibergen NR, Moore JD, Mattmann ME, and Blackwell HE (2015) Potent and Selective Modulation of the RhlR Quorum Sensing Receptor by Using Non-native Ligands: An Emerging Target for Virulence Control in *Pseudomonas aeruginosa*, *ChemBioChem* 16, 2348–2356. [PubMed: 26460240]

29. Lindemann A, Pessi G, Schaefer AL, Mattmann ME, Christensen QH, Kessler A, Hennecke H, Blackwell HE, Greenberg EP, and Harwood CS (2011) Isovaleryl-homoserine lactone, an unusual branched-chain quorum-sensing signal from the soybean symbiont *Bradyrhizobium japonicum*, *Proc. Nat. Acad. Sci. USA* 108, 16765–16770. [PubMed: 21949379]
30. Smith KM, Bu Y, and Suga H (2003) Library Screening for Synthetic Agonists and Antagonists of a *Pseudomonas aeruginosa* Autoinducer, *Chem. Biol* 10, 563–571. [PubMed: 12837389]
31. Morkunas B, Galloway WR, Wright M, Ibbeson BM, Hodgkinson JT, O'Connell KM, Bartolucci N, Della Valle M, Welch M, and Spring DR (2012) Inhibition of the production of the *Pseudomonas aeruginosa* virulence factor pyocyanin in wild-type cells by quorum sensing autoinducer-mimics, *Org. Biomol. Chem* 10, 8452–8464. [PubMed: 23014532]
32. Ikeda T, Kajiyama K, Kita T, Takiguchi N, Kuroda A, Kato J, and Ohtake H (2001) The Synthesis of Optically Pure Enantiomers of N-Acyl-homoserine Lactone Autoinducers and Their Analogues, *Chem. Lett*, 314–315.
33. Gerdt Joseph P., McInnis Christine E., Schell Trevor L., Rossi Francis M., and Blackwell Helen E. (2014) Mutational Analysis of the Quorum-Sensing Receptor LasR Reveals Interactions that Govern Activation and Inhibition by Nonlactone Ligands, *Chem. Biol*, 1–9. [PubMed: 24439204]
34. McInnis CE, and Blackwell HE (2011) Thiolactone modulators of quorum sensing revealed through library design and screening, *Bioorganic Med. Chem* 19, 4820–4828.
35. Boursier ME, Manson DE, Combs JB, and Blackwell HE (2018) A comparative study of non-native N-acyl L-homoserine lactone analogs in two *Pseudomonas aeruginosa* quorum sensing receptors that share a common native ligand yet inversely regulate virulence, *Bioorganic Med. Chem*, in press (DOI: 10.1016/j.bmc.2018.05.018).
36. Janssens JCA, Metzger K, Daniels R, Ptacek D, Verhoeven T, Habel LW, Vanderleyden J, De Vos DE, and De Keersmaecker SCJ (2007) Synthesis of N-Acyl Homoserine Lactone Analogues Reveals Strong Activators of SdiA, the *Salmonella enterica* Serovar Typhimurium LuxR Homologue, *Appl. Environ. Microbiol* 73, 535–544. [PubMed: 17085703]
37. Glansdorp FG, Thomas GL, Lee JJK, Dutton JM, Salmond GPC, Welch M, and Spring DR (2004) Synthesis and stability of small molecule probes for *Pseudomonas aeruginosa* quorum sensing modulation, *Org. Biomol. Chem* 2, 3329–3336. [PubMed: 15534711]
38. Zhang R. g., Pappas KM, Brace JL, Miller PC, Oulmassov T, Molyneaux JM, Anderson JC, Bashkin JK, Winans SC, and Joachimiak A (2002) Structure of a bacterial quorum-sensing transcription factor complexed with pheromone and DNA, *Nature* 417, 971–974. [PubMed: 12087407]
39. Lindsay A, and Ahmer BMM (2005) Effect of sdiA on Biosensors of N-Acylhomoserine Lactones, *J. Bacteriol* 187, 5054–5058. [PubMed: 15995228]
40. Lee J-H, Lequette Y, and Greenberg EP (2006) Activity of purified QscR, a *Pseudomonas aeruginosa* orphan quorum-sensing transcription factor, *Mol. Microbiol* 59, 602–609. [PubMed: 16390453]
41. Pearson JP, Pesci EC, and Iglewski BH (1997) Roles of *Pseudomonas aeruginosa* las and rhl quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes, *J. Bacteriol* 179, 5756–5767. [PubMed: 9294432]
42. de Kievit TR, Gillis R, Marx S, Brown C, and Iglewski BH (2001) Quorum-Sensing Genes in *Pseudomonas aeruginosa* Biofilms: Their Role and Expression Patterns, *Appl. Environ. Microbiol* 67, 1865–1873. [PubMed: 11282644]
43. O'Reilly MC, and Blackwell HE (2016) Structure-Based Design and Biological Evaluation of Triphenyl Scaffold-Based Hybrid Compounds as Hydrolytically Stable Modulators of a LuxR-Type Quorum Sensing Receptor, *ACS Infect. Dis* 2, 32–38. [PubMed: 26807436]

Figure 1:
Simplified scheme of the *P. aeruginosa* QS network. Black arrows represent positive feedback (autoinducer synthesis/transcriptional regulation/receptor binding, etc.). Flat red arrows represent negative regulation.

Figure 2:

Structures of RhIR native ligand **1** (BHL), two previously reported RhIR modulators used as controls (**2**, **3**), and a focused library of compounds designed to probe SARs of RhIR (**4–33**). Compounds **21** and **22** were previously characterized and included for comparison in the current study (*28*).

Critical components for activity

Figure 3:

Summary of key SAR trends for RhlR activators as revealed in this study. Red moieties are vital for activation. Blue moieties improve agonism beyond levels achieved by the native ligand BHL. The amide shown in green is critical for both activation and inhibition.

Table 1:Primary RhIR assay data for BHL analog library in *E. coli*.^a

Compound	% activation ^b	% inhibition ^c	Compound	% activation ^b	% inhibition ^c
1 (BHL)	51	–	18	32	–
2 (D8)	54	–	19	70	–
3 (S4)	76	–	20	2	24
4	25	–	21 (S5)	34	–
5	25	–	22 (B8)^d	1	31
6	62	–	23	1	55
7	86	–	24	0	6
8	55	–	25	2	11
9	1	25	26	71	–
10	5	7	27	1	35
11	28	–	28	35	–
12	41	–	29	0	45
13	21	–	30	0	20
14	52	–	31	2	12
15	45	–	32	0	–
16	34	–	33	0	57
17	79	–			

^aSee Methods for details. SEM of n = 3 trials did not exceed $\pm 10\%$.^bLibrary compounds screened at 10 μM . RhIR activity measured relative to BHL at 1 mM.^cLibrary compounds screened at 1 mM in the presence of 10 μM BHL. Values expressed relative to activation of 10 μM BHL alone. – = RhIR activation observed that was $\leq 10\%$ to that of 10 μM BHL.

Table 2:

EC₅₀ values for selected AHLs in RhIR in *E. coli* and *P. aeruginosa*.^a

Compound	<i>E. coli</i>		<i>P. aeruginosa</i>	
	EC ₅₀ (μM) ^b	95% CI (μM)	EC ₅₀ (μM) ^b	95% CI (μM)
1 (BHL)	8.95	5.86 – 13.7	8.08	6.09 – 10.7
2 (D8)	7.93	6.28 – 10.02	–	–
3 (S4)	1.58	1.32 – 1.90	1.22	1.03 – 1.45
5	10.83	6.59 – 17.80	–	–
6	4.89	3.67 – 6.53	–	–
7	1.02	0.67 – 1.55	1.42	1.08 – 1.86
8	7.77	5.61 – 10.8	–	–
14	6.93	5.52 – 8.71	–	–
17	1.78	1.37 – 2.31	1.41	1.14 – 1.74
19	2.76	2.23 – 3.42	–	–
26	4.87	3.46 – 6.84	3.82	2.57 – 5.66
28	27.4	16.1 – 46.6	14.3	8.76 – 23.5

^aSee Methods for details. Assays performed in triplicate; 95% confidence intervals (CIs) calculated from the SEM of n = 3 trials.

^bFor the full agonism traces, see Figures S1 and S2 for *E. coli* and *P. aeruginosa*, respectively. Additional trace characterization listed in Table S4; discussions of dose-response curve shapes are in Notes S1 and S2.