Traumatic brain injury associated with dementia risk among people with type 1 diabetes

Paola Gilsanz, ScD, Kathleen Albers, MPH, Michal Schnaider Beer, PhD, Andrew J. Karter, PhD, Charles P. Quesenberry, Jr., PhD, and Rachel A. Whitmer, PhD

Neurology® 2018;91:e1611-e1618. doi:10.1212/WNL.0000000000006391

Correspondence

P. Gilsanz
Paola.Gilsanz@kp.org

Abstract

Objective

To examine the association between traumatic brain injury (TBI) and dementia risk among a cohort of middle-aged and elderly individuals with type 1 diabetes (T1D).

Methods

We evaluated 4,049 members of an integrated health care system with T1D ≥ 50 years old between January 1, 1996, and September 30, 2015. Dementia and TBI diagnoses throughout the study period were abstracted from medical records. Cox proportional hazards models estimated associations between time-dependent TBI and dementia adjusting for demographics, HbA1c, nephropathy, neuropathy, stroke, peripheral artery disease, depression, and dysglycemic events. Fine and Gray regression models evaluated the association between baseline TBI and dementia risk accounting for competing risk of death.

Results

A total of 178 individuals (4.4%) experienced a TBI and 212 (5.2%) developed dementia. In fully adjusted models, TBI was associated with 3.6 times the dementia risk (hazard ratio [HR] 3.64; 95% confidence interval [CI] 2.34, 5.68). When accounting for the competing risk of death, TBI was associated with almost 3 times the risk of dementia (HR 2.91; 95% CI 1.29, 5.68).

Conclusion

This study demonstrates a marked increase in risk of dementia associated with TBI among middle-aged and elderly people with T1D. Given the complexity of self-care for individuals with T1D, and the comorbidities that predispose them to trauma and falls, future work is needed on interventions protecting brain health in this vulnerable population, which is now living to old age.

RELATED ARTICLE

Comment

Type 1 diabetes, traumatic brain injury, and dementia: Interactions or just bad luck?

[Page 781](#)

MORE ONLINE

CME Course

NPub.org/cmelist

From the Kaiser Permanente Division of Research (P.G., K.A., A.J.K., C.P.Q., R.A.W.), Oakland, CA; Department of Epidemiology and Biostatistics (P.G., R.A.W.), University of California, San Francisco; Department of Public Health Sciences (R.A.W.), University of California, Davis; Icahn School of Medicine at Mount Sinai (M.S.B.), New York, NY; The Joseph Sagol Neuroscience Center (M.S.B.), Sheba Medical Center, Ramat Gan, Israel; and Department of Epidemiology (A.J.K.), University of Washington, Seattle.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Glossary

CDC = Centers for Disease Control and Prevention; **CI** = confidence interval; **HbA1c** = glycosylated hemoglobin; **HR** = hazard ratio; **ICD-9** = *International Classification of Diseases-9*; **KPNC** = Kaiser Permanente Northern California; **T1D** = type 1 diabetes; **TBI** = traumatic brain injury.

In 2010, over 2 million people in the United States sustained a traumatic brain injury (TBI),¹ and between 3.2 and 5.3 million Americans live with long-term disabilities due to TBI.^{1,2} In addition to potential physical and functional impairment,¹ TBIs can have long-term cognitive effects, from mild cognitive dysfunction to more serious chronic diseases such as dementia.³ Falls are the most common cause of TBIs across all age groups and contribute to 47% of TBI-related emergency department visits, hospitalizations, and deaths.⁴ Older individuals with diabetes are at an elevated risk of falls,⁵⁻⁷ possibly due to diabetic complications such as hypoglycemia,^{8,9} peripheral neuropathy,⁵ or diabetic retinopathy.⁸ The link between TBIs and dementia is of particular concern for people with diabetes because dementia is associated with worse diabetes self-management and treatment¹⁰ and increased likelihood of hypoglycemic events.¹¹ This may lead to a dangerous downward spiral diminishing cognitive abilities and overall health. Yet it remains unknown if TBIs among people with type 1 diabetes (T1D) is associated with dementia risk. As incidence of T1D is increasing among youth¹² and people with T1D increasingly live to an age conferring dementia risk,¹³ it is important to evaluate the effect of TBI on dementia risk in elderly T1D.

Methods

Study population

Individuals enrolled within the Kaiser Permanente Northern California (KPNC) health plan were ascertained for this project if they met the definition of having a diagnosis of T1D within their electronic medical health record. Overall, KPNC provides comprehensive medical services to a diverse, multi-cultural population of over 4 million members within the Northern California Bay Area, Sacramento, and San Joaquin valleys. Members are representative of the greater Northern California population with some exceptions: KPNC members tend to be more educated and more affluent.¹⁴⁻¹⁶

Within the Kaiser Permanente Northern California Diabetes Registry, which is a compilation of all KPNC members with diabetes mellitus, individuals with T1D were ascertained using KPNC's electronic medical health records. The Registry, developed in 2003, identifies KPNC members with diabetes through a multiprong approach of reviewing pharmacy data, laboratory results, and diagnoses from both inpatient and outpatient records using ICD-9 codes.^{17,18}

Individuals were considered to have a T1D diagnosis if they met the following 3 criteria during a review of electronic

health records for all Diabetes Registry members: (1) at least 75% of their diagnostic codes indicate T1D; (2) filled an insulin prescription at any time during the study period; and (3) did not fill prescriptions of any hypoglycemic agents other than insulin.¹⁹

An individual's cohort entry date was the first day after January 1, 1996, that a member was at least 50 years old and diagnosed with T1D during the study period. A total of 4,049 individuals with T1D and without dementia at cohort entry were followed through September 30, 2015.

Standard protocol approvals, registrations, and patient consents

This study was approved by the KPNC internal review board. The requirement for patient informed consent was waived since analyses were conducted on preexisting data.

Traumatic brain injury

TBI was defined as ICD-9 codes from inpatient and outpatient settings for intracranial injury excluding those with skull fracture (850.x-854.x) and late effects of injuries to the nervous system (907.0) in electronic medical records. Both a history of TBI (i.e., TBI occurring between 1996 and cohort entry) as well as TBIs occurring during follow-up were included in analyses. If an individual experienced multiple TBIs, only the first TBI was included. Based off a Centers for Disease Control and Prevention (CDC) Morbidity and Mortality Weekly Report, a broader definition of TBI also included skull fractures (ICD-9 codes: 800, 801, 803, 804), injury to optic nerve and pathways (ICD-9 codes: 950.1, 950.2, 950.3), and head injuries not otherwise specified (ICD-9 code: 959.01).⁴

Dementia diagnosis

Electronic health records were utilized to identify dementia diagnoses between January 1, 1996, and September 30, 2015, from both inpatient and outpatient visits using the following ICD-9 codes: Alzheimer disease (331.0), vascular dementia (290.4x), and other/nonspecific dementia (290.0, 290.1x, 290.2x, 290.3, 294.1, 294.2x, and 294.8). Dementia has been successfully identified using ICD-9 codes in previous studies in this population²⁰⁻²⁶ and a similar set of ICD-9 codes had a sensitivity of 77% and a specificity of 95% compared with a consensus diagnosis of dementia utilizing medical records review, physical examination, structured interviews, and a neuropsychiatric battery.²⁷

Covariates

Demographic data, including age, sex, and self-reported race and ethnicity, were extracted from KPNC health plan

membership databases. Baseline glycosylated hemoglobin (HbA1c) was obtained from the Kaiser Permanente laboratory measurement most proximal to baseline. The following baseline diabetes complications were ascertained from the KPNC electronic health record between January 1, 1996, and cohort entry using ICD-9 code definitions: nephropathy (ICD-9 codes: 250.4x, 585x, 583.81x, 581.81x), neuropathy (ICD-9 codes: 354.x, 355.x), stroke (ICD-9 codes: 431.x, 430.x, 433.x1, 434.x, 435.x excluding 435.2x), peripheral artery disease (ICD-9 codes: 440.x, 441.x, 442.0, 442.3, 443.81, 443.9), depression (ICD-9 codes: 296.2x, 296.3x, 298.0x, 300.4x, 309.0x, 309.28, 311), and severe hypoglycemic (ICD-9 codes: 251.0, 251.1, 251.2) or hyperglycemic (ICD-9 codes: 250.1x, 250.2x, 249.2x) events. Missing indicators were used for those missing data on HbA1c (n = 307) or race (n = 113); there were no other missing data.

Statistical analysis

We examined the distribution of characteristics of members with and without TBI using χ^2 and *t* tests.

Cox proportional hazards models with age as the time scale estimated the association between narrow and broader definitions of TBI and dementia risk. Covariates were added in 5 groups: demographics (age, race, sex), microvascular complications (neuropathy, nephropathy), macrovascular disease (stroke, peripheral artery disease), mental health (depression), and glycemic control (HbA1c, dysglycemic events). Possible effect modification by sex, stroke, or severe dysglycemic events was examined through interactions terms. Members were followed until dementia diagnosis, death, lapse in KPNC membership, or the end of the study period (September 30, 2015). Deaths were ascertained from KPNC hospital records, California death certificates, and Social Security Administration datasets.

To diminish the likelihood of reverse causation, we examined the risk of dementia diagnosis occurring more than 1 year after TBI diagnosis compared to individuals with no TBI diagnosis. Covariates were added to the Cox proportional hazards models (age as time scale) in the manner described above.

We conducted 2 analyses examining the association between a history of TBI at cohort entry and dementia accounting for the competing risk of death. First, the Fine and Gray regression model was implemented to estimate the HR of dementia accounting for the competing risk of death.²⁸ Second, we estimated the 5- to 30-year cumulative incidence of dementia in 5-year increments conditional on survival free of dementia until age 50. These estimates were obtained using the Practical Incidence Estimator macro, which incorporates information on age-specific death rates.²⁹

All statistical analyses were conducted on SAS statistical software version 9.3 (SAS Institute, Cary, NC). A *p* value below 0.05 was considered statistically significant.

Data availability

Data are available to qualified investigators from the Kaiser Permanente Institutional Review Board Institutional Data Access/Ethics Committee for the purposes of replicating procedures and results.

Results

Overall, the mean age at entry was 57.0 years (SD 9.2 years), 53% of participants were male, and 22% were nonwhite (table 1). Four percent of the sample (n = 178) experienced a TBI, 9% (n = 348) experienced a TBI/head injury, and 5% (n = 212) developed dementia. Participants were followed for an average of 5.8 years (SD 5.3 years). Among the 23 individuals who developed dementia after a TBI, there was an average of 4.1 years between TBI and dementia date (SD 4.4; range 0.003–15.8 years). The time from TBI to dementia diagnosis was within 1 year for 8 individuals (34.8%), between 1 and 3 years for 5 individuals (21.7%), between 4 and 8 years for 6 individuals (26.1%), and between 8 and 16 years for 4 individuals (17.4%). Compared to those without TBI, those with TBI were more likely to have baseline neuropathy, stroke, and depression. By the end of follow-up, 21% of the sample had died, 25% had a lapse in health plan membership greater than 90 days, and 49% survived dementia-free and remained members.

Adjusting for demographics, people with TBI had 3.8 times the risk of developing dementia than individuals without TBI (hazard ratio [HR] 3.77; 95% confidence interval [CI] 2.43, 5.85; table 2). This association remained strong after adjusting for demographics, microvascular and macrovascular complications, depression, HbA1c, and severe dysglycemic events (HR 3.64; 95% CI 2.34, 5.68). There was no evidence of effect modification by sex (TBI * sex interaction term *p* value = 0.58), baseline stroke (TBI * stroke interaction term *p* value = 0.90), or severe hypoglycemia (TBI * hypoglycemia interaction term *p* value = 0.50). There was evidence of the effect of TBI being worse among people with severe hyperglycemia (TBI * hyperglycemia interaction term *p* value = 0.003; HR_{TBI, hyperglycemia} 4.38, 95% CI 1.18, 16.24; HR_{TBI, no hyperglycemia} 3.57, 95% CI 2.23, 5.71); however, these results should be interpreted with caution due to small numbers.

TBI continued to be associated with dementia when excluding dementia cases occurring within 1 year of TBI diagnosis. Individuals who experienced a TBI had more than double the risk of dementia more than 1 year after TBI diagnosis compared to individuals without TBI (table 3).

The broader definition of TBI was also associated with dementia risk in models adjusting for demographics (HR 2.91; 95% CI 2.01, 4.20; table 4) and in fully adjusted models (HR 3.03; 95% CI 2.09, 4.39). There was no evidence that the relationship between the broader definition of TBI and dementia was modified by sex (TBI * sex interaction term *p* value = 0.29), baseline stroke (TBI * stroke interaction term *p*

Table 1 Baseline characteristics by traumatic brain injury (TBI) during study period

	Overall, n (%) or mean (SD)	No TBI, n (%) or mean (SD)	TBI, n (%) or mean (SD)	<i>p</i> Value
Total	4,049 (100)	3,871 (95.6)	178 (4.4)	
Dementia	212 (5.2)	189 (4.9)	23 (12.9)	<0.001
Age, y, mean (SD)				
At entry	57.0 (9.2)	57.0 (9.3)	55.8 (7.9)	0.06
At dementia diagnosis	63.4 (9.0)	72.4 (9.5)	65.9 (10.3)	<0.01
Sex				0.75
Female	1913 (47.3)	1831 (47.3)	82 (46.1)	
Male	2,136 (52.8)	2040 (52.7)	96 (53.9)	
Race and ethnicity				0.06
White	3,142 (77.6)	3,006 (77.7)	136 (76.4)	
Black	220 (5.4)	208 (5.0)	12 (6.7)	
Hispanic	247 (6.1)	237 (5.6)	10 (5.6)	
Asian	160 (4.0)	153 (3.8)	7 (3.9)	
Other	167 (4.1)	154 (3.9)	13 (7.3)	
Missing	113 (2.8)	113 (2.7)	0 (0)	
Comorbidities and complications				
Nephropathy	661 (16.3)	628 (16.2)	33 (18.5)	0.41
Neuropathy	530 (13.1)	489 (12.6)	41 (23.0)	<0.001
Stroke	170 (4.2)	152 (3.9)	18 (10.1)	<0.001
Peripheral artery disease	286 (7.0)	269 (7.0)	17 (9.6)	0.19
Depression	696 (17.2)	649 (16.8)	47 (26.4)	<0.01
Hypoglycemic event	487 (12.3)	459 (11.9)	28 (15.7)	0.12
Hyperglycemic event	267 (6.6)	254 (6.6)	13 (7.3)	0.70
HbA1c, mean (SD)	8.0 (1.7)	8.0 (1.7)	8.1 (1.9)	0.88

Abbreviation: HbA1c = glycosylated hemoglobin.

Percentages are shown as column percentages. *p* Values obtained from χ^2 tests (for binary or categorical variables) and *t* tests (for continuous variables).

value = 0.43), severe hypoglycemia (TBI * hypoglycemia interaction term *p* value = 0.73), or severe hyperglycemia (TBI * hyperglycemia interaction term *p* value = 0.31).

In Fine and Gray regression models adjusted for demographics and accounting for the competing risk of death, a history of TBI at cohort entry (narrow definition; *n* = 59) was associated with more than triple the risk of dementia (HR 3.50; 95% CI 1.58, 6.65; table 5). The association persisted after further controlling for microvascular and macrovascular complications, depression, HbA1c, and dysglycemic events and TBI continued to be associated with almost a 3-fold increase in dementia risk (HR 2.91; 95% CI 1.29, 5.68).

The estimates of cumulative incidence of dementia, which account for death rates, were consistently higher for

individuals with a history of TBI at cohort entry. However, the CIs were very wide due to the small number of dementia cases and encompass the CIs of those without a history of TBI at cohort entry (table 6, figure). The 15-year cumulative incidence for people with TBI was 22.38% (95% CI 0.29%, 41.15%) compared to 2.90% (95% CI 1.85%, 3.79%) among people without TBI.

Discussion

In a large cohort of people with T1D at least 50 years old, individuals who experienced TBI had 3.6 times the risk of developing dementia compared to individuals without TBI even after adjusting for demographics, depression, microvascular and macrovascular complications, HbA1c, and severe

Table 2 Traumatic brain injury and risk of dementia in individuals with type 1 diabetes

Adjusted for	HR (95% CI)
Demographics	3.77 (2.43, 5.85)
Demographics and microvascular complications	3.68 (2.37, 5.70)
Demographics and macrovascular complications	3.43 (2.20, 5.34)
Demographics and depression	3.67 (2.37, 5.68)
Demographics and HbA1c	3.95 (2.54, 6.13)
Demographics and dysglycemic events	3.80 (2.45, 5.89)
Demographics, microvascular and macrovascular complications, depression, HbA1c, and dysglycemic events	3.64 (2.34, 5.68)

Abbreviations: CI = confidence interval; HbA1c = glycosylated hemoglobin; HR = hazard ratio.
HRs estimated from Cox proportional hazards models with age as time scale. Demographics include age, sex, and race/ethnicity. Microvascular complications include neuropathy and nephropathy. Macrovascular complications include stroke and peripheral artery disease.

dysglycemic events. A broader definition of TBI based off work by the CDC was associated with approximately a threefold risk of dementia. In models accounting for the competing risk of death, a history of TBI at cohort entry was associated with almost a threefold dementia risk adjusting for demographics, depression, microvascular and macrovascular complications, HbA1c, and severe dysglycemic events. The 15-year cumulative incidence of dementia among individuals with TBI history at cohort entry was almost 8 times that of people without TBI (2.9% compared to 22.4%). However, the CIs overlap likely due to small numbers and the difference in cumulative incidence rates should be interpreted with caution. Given the large social and economic burden of dementia, coupled with the lack of disease-modifying treatments, it is

Table 3 Risk of dementia at least 1 year after traumatic brain injury diagnosis among individuals with type 1 diabetes

Adjusted for	HR (95% CI)
Demographics	2.45 (1.44, 4.17)
Demographics and microvascular complications	2.39 (1.40, 4.06)
Demographics and macrovascular complications	2.24 (1.31, 3.82)
Demographics and depression	2.38 (1.40, 4.04)
Demographics and HbA1c	2.55 (1.50, 4.33)
Demographics and dysglycemic events	2.46 (1.50, 4.33)
Demographics, microvascular and macrovascular complications, depression, HbA1c, and dysglycemic events	2.35 (1.38, 4.02)

Abbreviations: CI = confidence interval; HbA1c = glycosylated hemoglobin; HR = hazard ratio.

Table 4 Hazard ratio (HR) of broader definition of traumatic brain injury and risk of dementia among individuals with type 1 diabetes

Adjusted for	HR (95% CI)
Demographics	2.91 (2.01, 4.20)
Demographics and microvascular complications	2.83 (1.95, 4.09)
Demographics and macrovascular complications	2.79 (1.93, 4.04)
Demographics and depression	2.82 (1.95, 4.08)
Demographics and HbA1c	3.19 (2.21, 4.61)
Demographics and dysglycemic events	2.91 (2.02, 4.21)
Demographics, micro- and macrovascular complications, depression, HbA1c, and dysglycemic events	3.03 (2.09, 4.39)

Abbreviations: CI = confidence interval; HbA1c = glycosylated hemoglobin; HR = hazard ratio.
HRs estimated from Cox proportional hazards models with age as time scale. Microvascular complications include neuropathy and nephropathy. Macrovascular complications include stroke and peripheral artery disease.

critical that the link between TBI and dementia among people with T1D be further explored.

Our results suggest that the harmful effect of TBI on dementia risk may be greater among individuals with T1D than the general population. Studies examining the association between TBIs and dementia among the general population have had mixed results.^{3,30–32} A recent meta-analysis including 32 studies found that head injuries in the general population were associated with a 63% increased risk of dementia and a 51%

Table 5 Risk of dementia associated with baseline traumatic brain injury accounting for competing risk of death among individuals with type 1 diabetes

Adjusted for	HR (95% CI)
Demographics	3.50 (1.58, 6.65)
Demographics and microvascular complications	3.39 (1.53, 6.46)
Demographics and macrovascular complications	2.90 (1.29, 5.62)
Demographics and depression	3.32 (1.49, 6.37)
Demographics and HbA1c	3.59 (1.62, 6.83)
Demographics and dysglycemic events	3.50 (1.58, 6.65)
Demographics, microvascular and macrovascular complications, depression, HbA1c, and dysglycemic events	2.91 (1.29, 5.68)

Abbreviations: CI = confidence interval; HbA1c = glycosylated hemoglobin; HR = hazard ratio.
HRs estimated from Fine and Gray regression models with time since cohort entry as time scale. Demographics include age at cohort entry, sex, and race/ethnicity. Microvascular complications include neuropathy and nephropathy. Macrovascular complications include stroke and peripheral artery disease.

Table 6 Cumulative incidence of dementia by history of traumatic brain injury (TBI) at cohort entry conditional on dementia-free survival until age 50

	No history of TBI cumulative incidence (95% CI)	History of TBI cumulative incidence (95% CI)
5 years risk	0.60 (0.22, 0.96)	4.87 (0.00, 11.54)
10 years risk	1.71 (0.99, 2.36)	10.40 (0.00, 22.72)
15 years risk	2.90 (1.85, 3.79)	22.38 (0.29, 41.15)
20 years risk	7.26 (5.14, 8.81)	26.87 (0.00, 46.72)
25 years risk	12.29 (8.95, 14.36)	33.12 (0.00, 55.46)
30 years risk	16.14 (11.59, 18.49)	36.34 (5.40, 59.29)

Abbreviation: CI = confidence interval.

increased risk of Alzheimer disease.³² In a study including more than 3 million individuals, a mild case of TBI was associated with 63% greater risk of dementia compared to individuals without TBI, while those with severe TBI had double the dementia risk, and those with multiple TBI had 2.8 times the risk of dementia.³³ Consistent with some,³³ but not all prior studies,³¹ we found a strong association between TBI and dementia even after excluding dementia cases occurring within the first year after TBI diagnosis when possible bias due to reverse causation is greatest. However, it is challenging to compare our results with other studies given possible differences in age, TBI severity, and time since TBI. Future studies including both people with and without T1D in their sample will be better positioned to directly compare the magnitude of the effect of TBI on dementia risk across the 2 groups.

Figure Cumulative incidence of dementia by baseline traumatic brain injury (TBI) status conditional on dementia-free survival until age 50

TBIs trigger many neurotoxic events, ranging in timing from acute (e.g., axonal shearing, vasospasm ischemia) to secondary (e.g., inflammation, oxidative stress, apoptosis) to chronic (e.g., increased levels of amyloid precursor protein, hyperphosphorylation of tau).³⁴ It has been hypothesized that several of these events could elevate risk of dementia including inflammation, oxidative stress, and the production of β -amyloid.³⁵ TBIs also trigger perturbations in brain glucose metabolism resulting in an increased demand for glucose, especially in the first 48 hours.^{36,37} It has been estimated that in the general population, roughly 40% of patients with TBI experience hyperglycemia, a risk factor for oxidative stress.³⁷

Cognitive reserve (i.e., resilience to neuropathology) may be a key role in mitigating the effects of TBI on dementia risk. It has been hypothesized that TBIs occurring at older ages may be more strongly associated with dementia risk than those occurring at younger ages due to declines in cognitive reserves associated with aging.³⁵ Similarly, individuals with T1D may already have greater levels of cerebrovascular damage (i.e., less cognitive reserve), leaving them more susceptible to the harmful effects of TBIs. For example, repeated exposure to hyperglycemia has been associated with gray matter density loss.^{38,39} Animal models suggest that those with diabetes experienced greater levels of chronic oxidative stress that worsened their neurologic dysfunction after brain injury than those without diabetes.⁴⁰

The primary strength of this study is the uniquely large cohort of over 4,000 people with T1D ages 50 and above with information on a wide range of microvascular and macrovascular complications as well as baseline HbA1c. Another strength is the high level of membership stability, which enabled us to follow this cohort up to 19 years. However, our study is not exempt from limitations. We only have information on diabetes beginning in 1996 and therefore are unable to examine if longer duration of T1D is associated with a greater risk of TBI. Another limitation of the study is that dementia diagnoses were not confirmed via medical record review. However, a previous study utilizing a similar set of diagnostic codes showed 77% sensitivity and 95% specificity compared with a consensus diagnosis of dementia utilizing medical records review, structured interviews, clinical evaluation, and various cognitive tests.²⁷ Due to small numbers, we are unable to explore if the association between TBI and dementia varies by time since TBI or type of brain injury. We cannot rule out the possibility of reverse causation as the mean number of years between TBI and dementia diagnosis was 4.1 years. We are unable to examine the direct effect of TBI on brain health due to a lack of brain imaging. We also do not have information regarding the cause or severity of the TBI. Additional studies in other large-scale diabetes databases should be conducted to see if these findings are replicated and assess for possible effect modification by diabetes duration, time since TBI, or brain injury type.

This study demonstrates dramatic increases in risk of dementia among people with T1D who experience a TBI independent of

micro complications and macro complications, depression, severe dysglycemic events, and HbA1c levels. Hypoglycemia and proprioception dysfunction associated with peripheral neuropathy put patients with T1D at uniquely elevated risk of TBI, and thus ultimately increase the likelihood of dementia. Dementia, in turn, worsens diabetes self-management and treatment¹⁰ and may result in increased likelihood of hypoglycemic events⁴¹ and neuropathic damage. This could trigger a dangerous downward spiral diminishing cognitive abilities, overall health, and safety. Relatedly, individuals with comorbid dementia and diabetes have reduced levels of activities of daily living and substantial reliance on assistance for personal care.⁴² Given the limited treatment options once a TBI has occurred and the strong relationship between TBI and dementia, it is important for future research to identify modifiable risk factors for TBI and explore possible interventions aimed at reducing the risk of TBI among people with T1D and the general population. Lower levels of cognitive reserve may place individuals with T1D at greater risk of the deleterious effects of TBI, which may explain the greater magnitude of dementia associated with TBI in this sample than in prior studies. Given the complexity of self-care for elderly individuals with T1D, and the comorbidities that predispose them to trauma and falls, future work is needed on preventive measures for protecting brain health in this vulnerable population, which is now living to old age.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Author contributions

R.A. Whitmer and P. Gilsanz contributed to the conception and design of the study. P. Gilsanz was responsible for the statistical analysis. All coauthors interpreted the data and contributed significantly to the manuscript.

Study funding

This work was supported by grant R01 AG047500 (PI: Dr. Whitmer). Dr. Gilsanz is supported by the UCSF Training for Research on Aging and Chronic Disease (T32 AG049663). Dr. Karter is supported by grants R01 DK103721 (PI: Dr. Karter) and R01 DK081796 (PI: Dr. Karter) and the Health Delivery Systems Center for Diabetes Translational Research (P30 DK092924).

Publication history

Received by *Neurology* February 13, 2018. Accepted in final form June 8, 2018.

References

- Centers for Disease Control and Prevention. Report to Congress on Traumatic Brain Injury in the United States. Atlanta, GA: Centers for Disease Control and Prevention; 2015.
- Centers for Disease Control and Prevention. CDC grand rounds: reducing severe traumatic brain injury in the United States. *MMWR Morb Mortal Wkly Rep* 2013;62:549–552.
- Gardner RC, Burke JF, Nettiksimmons J, Kaup A, Barnes DE, Yaffe K. Dementia risk after traumatic brain injury vs nonbrain trauma: the role of age and severity. *JAMA Neurol* 2014;71:1490–1497.
- Taylor CA, Bell JM, Breiding MJ, Xu L. Traumatic brain injury-related emergency department visits, hospitalizations, and deaths: United States, 2007 and 2013. *Morb Mortal Wkly Rep Surveill Summ* 2017;66:1–16.
- Schwartz AV, Hillier TA, Sellmeyer DE, et al. Older women with diabetes have a higher risk of falls: a prospective study. *Diabetes Care* 2002;25:1749–1754.
- Yau RK, Strotmeyer ES, Resnick HE, et al. Diabetes and risk of hospitalized fall injury among older adults. *Diabetes Care* 2013;36:3985–3991.
- Gregg EW, Beckles GL, Williamson DF, et al. Diabetes and physical disability among older U.S. adults. *Diabetes Care* 2000;23:1272–1277.
- Schwartz AV, Vittinghoff E, Sellmeyer DE, et al. Diabetes-related complications, glycemic control, and falls in older adults. *Diabetes Care* 2008;31:391–396.
- Malabu UH, Vangaveti VN, Kennedy RL. Disease burden evaluation of fall-related events in the elderly due to hypoglycemia and other diabetic complications: a clinical review. *Clin Epidemiol* 2014;6:287–294.
- Bunn F, Goodman C, Malone JR, et al. Managing diabetes in people with dementia: protocol for a realist review. *Syst Rev* 2016;5:5.
- Kirkman MS, Briscoe VJ, Clark N, et al. Diabetes in older adults: a consensus report. *J Am Geriatr Soc* 2012;60:2342–2356.
- Rogers MAM, Kim C, Banerjee T, Lee JM. Fluctuations in the incidence of type 1 diabetes in the United States from 2001 to 2015: a longitudinal study. *BMC Med* 2017;15:199.
- Miller RG, Secrest AM, Sharma RK, Songer TJ, Orchard TJ. Improvements in the life expectancy of type 1 diabetes: The Pittsburgh Epidemiology of Diabetes Complications study cohort. *Diabetes* 2012;61:2987–2992.
- Gordon NP, Kaplan GA. Some evidence refuting the HMO “favorable selection” hypothesis: the case of Kaiser Permanente. *Adv Health Econ Health Serv Res* 1991;12:19–39.
- Krieger N. Overcoming the absence of socioeconomic data in medical records: validation and application of a census-based methodology. *Am J Public Health* 1992;82:703–710.
- Gordan NP. Similarity of the Kaiser Permanente Senior Member Population in Northern California to the Non-Kaiser Permanente Covered and General Population of Seniors in Northern California: Statistics from the 2009 California Health Interview Survey. Oakland, CA: Kaiser Permanente Northern California Division of Research; 2012.
- Karter AJ, Schillinger D, Adams AS, et al. Elevated rates of diabetes in Pacific Islanders and Asian subgroups: the Diabetes Study of Northern California (DISTANCE). *Diabetes Care* 2013;36:574–579.
- Karter AJ, Ferrara A, Liu JY, Moffet HH, Ackerson LM, Selby JV. Ethnic disparities in diabetic complications in an insured population. *JAMA* 2002;287:2519–2527.
- Klompas M, Eggleston E, McVetta J, Lazarus R, Li L, Platt R. Automated detection and classification of type 1 versus type 2 diabetes using electronic health record data. *Diabetes Care* 2013;36:914–921.
- Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA. Inequalities in dementia incidence between six racial and ethnic groups over 14 years. *Alzheimers Dement* 2016;12:216–224.
- Exalto LG, Biessels GJ, Karter AJ, et al. Severe diabetic retinal disease and dementia risk in type 2 diabetes. *J Alzheimers Dis* 2014;42:S109–S117.
- Exalto LG, Biessels GJ, Karter AJ, et al. Risk score for prediction of 10 year dementia risk in individuals with type 2 diabetes: a cohort study. *Lancet Diabetes Endocrinol* 2013;1:183–190.
- Mayeda ER, Karter AJ, Huang ES, Moffet HH, Haan MN, Whitmer RA. Racial/ethnic differences in dementia risk among older type 2 diabetic patients: the Diabetes and Aging Study. *Diabetes Care* 2014;37:1009–1015.
- Whitmer RA, Karter AJ, Yaffe K, Quesenberry CP, Jr., Selby JV. Hypoglycemic episodes and risk of dementia in older patients with type 2 diabetes mellitus. *JAMA* 2009;301:1565–1572.
- Whitmer RA, Sidney S, Selby J, Johnston SC, Yaffe K. Midlife cardiovascular risk factors and risk of dementia in late life. *Neurology* 2005;64:277–281.
- Whitmer RA, Gustafson DR, Barrett-Connor E, Haan MN, Gunderson EP, Yaffe K. Central obesity and increased risk of dementia more than three decades later. *Neurology* 2008;71:1057–1064.
- Katon WJ, Lin EHB, Williams LH, et al. Comorbid depression is associated with an increased risk of dementia diagnosis in patients with diabetes: a prospective cohort study. *J Gen Intern Med* 2010;25:423–429.
- Lau B, Cole SR, Gange SJ. Competing risk regression models for epidemiologic data. *Am J Epidemiol* 2009;170:244–256.
- Beiser A, D’Agostino RB, Seshadri S, Sullivan LM, Wolf PA. Computing estimates of incidence, including lifetime risk: Alzheimer’s disease in the Framingham Study: the Practical Incidence Estimators (PIE) macro. *Stat Med* 2000;19:1495–1522.
- Raj R, Kaprio J, Korja M, Mikkonen ED, Jousilahti P, Siironen J. Risk of hospitalization with neurodegenerative disease after moderate-to-severe traumatic brain injury in the working-age population: a retrospective cohort study using the Finnish national health registries. *PLOS Med* 2017;14:e1002316.
- Perry DC, Sturm VE, Peterson MJ, et al. Association of traumatic brain injury with subsequent neurological and psychiatric disease: a meta-analysis. *J Neurosurg* 2016;124:S11–S26.
- Li Y, Li Y, Li X, et al. Head injury as a risk factor for dementia and Alzheimer’s disease: a systematic review and meta-analysis of 32 observational studies. *PLOS One* 2017;12:e0169650.
- Nordstrom A, Nordstrom P. Traumatic brain injury and the risk of dementia diagnosis: a nationwide cohort study. *Plos Med* 2018;15:e1002496.
- Logsdon AF, Lucke-Wold BP, Turner RC, Huber JD, Rosen CL, Simpkins JW. Role of microvascular disruption in brain damage from traumatic brain injury. *Compr Physiol* 2015;5:1147–1160.

35. Mendez MF. What is the relationship of traumatic brain injury to dementia? *J Alzheimers Dis* 2017;57:667–681.
36. Jalloh I, Carpenter KL, Helmy A, Carpenter TA, Menon DK, Hutchinson PJ. Glucose metabolism following human traumatic brain injury: methods of assessment and pathophysiological findings. *Metab Brain Dis* 2015;30:615–632.
37. Godoy DA, Behrouz R, Di Napoli M. Glucose control in acute brain injury: does it matter? *Curr Opin Crit Care* 2016;22:120–127.
38. Wessels AM, Simsek S, Remijnse PL, et al. Voxel-based morphometry demonstrates reduced grey matter density on brain MRI in patients with diabetic retinopathy. *Diabetologia* 2006;49:2474–2480.
39. Musen G, Lyoo IK, Sparks CR, et al. Effects of type 1 diabetes on gray matter density as measured by voxel-based morphometry. *Diabetes* 2006;55:326–333.
40. Elangovan V, Kohen R, Shohami E. Neurological recovery from closed head injury is impaired in diabetic rats. *J Neurotrauma* 2000;17:1013–1027.
41. Yaffe K, Falvey CM, Hamilton N, et al. Association between hypoglycemia and dementia in a biracial cohort of older adults with diabetes mellitus. *JAMA Intern Med* 2013;173:1300–1306.
42. Sinclair AJ, Girling AJ, Bayer AJ. Cognitive dysfunction in older subjects with diabetes mellitus: impact on diabetes self-management and use of care services: All Wales Research into Elderly (AWARE) Study. *Diabetes Res Clin Pract* 2000;50:203–212.