Published in final edited form as:

ACS Chem Neurosci. 2018 October 17; 9(10): 2307–2330. doi:10.1021/acscchemneuro.7b00528.

The DARK Side of Total Synthesis: Strategies and Tactics in Psychoactive Drug Production

Schuyler A. Chambers[†], Jenna M. DeSousa[†], Eric D. Huseman[†], and Steven D. Townsend^{*,†,‡}

[†]Department of Chemistry, Vanderbilt University, 7330 Stevenson Center, Nashville, Tennessee 37235, United States

[‡]Institute of Chemical Biology, Vanderbilt University, 896 Preston Research Building, Nashville, Tennessee 37232, United States

Abstract

Humankind has used and abused psychoactive drugs for millennia. Formally, a psychoactive drug is any agent that alters cognition and mood. The term “psychotropic drug” is neutral and describes the entire class of substrates, licit and illicit, of interest to governmental drug policy. While these drugs are prescribed for issues ranging from pain management to anxiety, they are also used recreationally. In fact, the current opioid epidemic is the deadliest drug crisis in American history. While the topic is highly politicized with racial, gender, and socioeconomic elements, there is no denying the toll drug mis- and overuse is taking on this country. Overdose, fueled by opioids, is the leading cause of death for Americans under 50 years of age, killing ca. 64,000 people in 2016. From a chemistry standpoint, the question is in what ways, if any, did organic chemists contribute to this problem? In this targeted review, we provide brief historical accounts of the main classes of psychoactive drugs and discuss several foundational total syntheses that ultimately provide the groundwork for producing these molecules in academic, industrial, and clandestine settings.

Graphical abstract

*Corresponding Author steven.d.townsend@vanderbilt.edu.

Author Contributions

S.D.T. conceptualized the review. All authors wrote, reviewed, and approved the final version of the manuscript.

Notes

The authors declare no competing financial interest.

Keywords

Opiate; opioid; psychoactive; stimulant; benzodiazepines; hallucinogens

1. INTRODUCTION

Psychoactive drugs affect cognition by influencing the synaptic operation of neurotransmitters in the central nervous system (CNS).¹⁻⁴ This action occurs through one of three general mechanisms:⁵ (1) as agonists that mimic the operation of a neurotransmitter; (2) as antagonists that block the action of a neurotransmitter; or (3) by blocking the reuptake of neuro-transmitters at the synapse. In this regard, psychoactive drugs mimic or enhance naturally occurring states of consciousness; for example, sleeping pills promote drowsiness while benzodiazepines create a state of relaxation. Long-term use, whether medically or recreationally, often leads to tolerance, requiring the user to increase a drug's dosage in order to achieve the baseline effect. As drug use increases, the user may also develop dependence in one of two forms. The first is psychological dependence, in which the user feels the urge to use the drug but abstinence poses no risk. The second form is physical dependence where serious physical and mental effects develop when a drug is withdrawn. In the United States millions of people use psychoactive drugs everyday (Table 1).⁶ While misuse is prominent, approximately one in three Americans are prescribed an opiate/opioid for pain management every year,⁷ a testament to the importance of licit drug use for maintaining human health and wellness.

2. SCOPE OF THIS REVIEW

2.1. The United States Overdose Epidemic.

While the goal of this survey is to review synthetic strategies used to produce the carbon frameworks of psychoactive drugs, it is undeniable that the scholarly exercises completed by organic chemists have guided illicit drug production. Thus, we would be neglectful to not briefly discuss the "DARK" and highly politicized nature of the topic. Virtually all known civilizations have used psychoactive drugs in an effort to self-medicate, anesthetize from emotional or physical trauma, or for recreational and religious purposes. In the United

States, aggressive marketing of painkillers has led to a generation of individuals with opioid use disorders and a death toll closing in on 200 people per day. In both the second world war and Vietnam war, the Department of Defense employed “speed” or amphetamine to empower soldiers on the frontlines and sedatives and neuroleptics to prevent mental breakdowns from combat stress.⁸ The crack-cocaine epidemic of the 1980s and early 1990s wreaked havoc among low-income, urban Americans. Similarly, “backyard”-produced methamphetamine reached its pinnacle among poorer rural Americans in the 1990s and early 21st century. One difference between those crises and the current iteration of the American drug epidemic is that opioid use disorder has its genesis at the pharmacy counter and crosses broad demographic groups.

2.2. Chemistry’s Role in the Crisis.

From a chemistry perspective, the current epidemic can likely be traced back to the introduction of OxyContin. “Oxy” is a semisynthetic opioid that shares structural homology with morphine, *vide infra*. Strong analgesics of this type have historically been reserved to manage pain associated with cancer therapy or surgery. During the mid-1990s oxycodone was marketed to treat chronic pain. As the use of opioids increased, pill distribution spread from primary-care practitioners to neighborhood dealers. Pills were also sold in large quantities from poorly regulated pain treatment centers colloquially known as “pill mills”. The problem appeared so rapidly that the American Public Health Association published a report in 2009 describing the prescription drug crisis titled “The promotion and marketing of OxyContin: Commercial triumph, public health tragedy”.⁹ Ultimately, the closing of pill mills combined with an inability to treat addiction initiated a dark chapter in psychoactive drug use: the resurgence of heroin and the emergence of fentanyl. This has resulted in nearly 1000 deaths per week due to drug overdose, two-thirds of which are opioid related fatalities. The problem is so great that the illicit drug trade is currently the second most lucrative industry in the world, after weapons.¹⁰

While the most problematic players in the drug crisis, and the largest section of this review, are opium alkaloids, they are not the only burdensome psychoactive drugs. In fact, deciding which agents are defined as drugs, how drug supply and use is regulated, and how society responds to drug users are hotly debated issues. The fact that the two drugs that cause the most impairment and damage in the world tobacco and alcohol are both legal in most countries illustrates this point. As does the fact that caffeine, the world’s most popular stimulant, is legal and unregulated worldwide. What role does organic chemistry play in the field of psychoactive drug synthesis and use? As one reads through the annals of published efforts, it is clear that legendary figures in the field, ranging from Larry Overman¹¹ and Phil Magnus,¹² to Kathy Parker¹³ and James White,¹⁴ have trained scores of students using these molecules as training tools. While the approaches are scholarly in nature, one cannot help but recognize the dichotomy: academic lessons assist in fueling both licit and clandestine drug use.

Given the level of intrigue that accompanies this class of molecules, several comprehensive accounts have been written to describe both the synthesis and the reactivity of these substrates.^{15–18} The gap that this review fills is to compare and contrast syntheses in

academic, industrial, and clandestine settings. Historically, academic pursuits have focused on demonstrating novel bond forming strategies and assembly tactics whereas industry has viewed their efforts as exercises in scale and efficiency. Meanwhile, the underworld of synthesis relies upon operational simplicity and material availability to make their syntheses practical. As a result, the de novo total syntheses of the academic world are commonly translated into large-scale semisyntheses in industrial settings that are further simplified for illicit manufacturing in makeshift clandestine laboratories. With this in mind, we chose to take a similar approach in the construction of this review and have highlighted various academic and industrial syntheses of the chosen psychoactive compounds before delving into their illicit synthesis as appropriate. It is our hope that this review will provide the reader with a foundation for delving deeper into the psychoactive drug literature and give context to how scholarly exercises are translated to clandestine drug synthesis.

3. THE OPIATES

3.1. Historical Perspective.

Opium extracts have been used for millennia. The earliest reference to opium production, in which the inhabitants of southern Mesopotamia cultivated *Papaver somniferum*, the opium poppy, predates 4000 B.C.¹⁹ Known as Hul Gil, or the joy plant, the art of poppy culling was passed from the Sumerians to the Assyrians, Babylonians, and the ancient Egyptians. Eventually, its cultivation spread along the Silk Road from the Mediterranean to the Far East where the British sale of opium in China was the catalyst for the Opium Wars of the mid-1800s.²⁰ By way of definition, opiates are defined as any compounds, natural or semisynthetic, that share structural homology with opium alkaloids. We use the term opioid to refer to substances that elicit similar CNS effects but do not share structural homology with naturally occurring poppy alkaloids.²¹ Regardless of their structural classification, opiates and opioids are large components of the illicit drug market, generating revenue in excess of \$55 billion per year.⁷ While the pharmacology of these molecules will be covered in detail in other parts of the DARK special issue, we briefly mention here that opiates and opioids act on the CNS to produce effects ranging from analgesia to sedation.

3.2. Structure.

The opiates (Figure 1) mimic the activity of the endorphins, the body's natural pain relievers. Structurally, the opiates feature a pentacyclic scaffold containing a phenolic A ring, a bridging bicyclic piperidine D ring, and a dihydrofuran like E ring (Figure 1C). While masked, most receptor active opiates and opioids share structural homology. These similarities are known as the "morphine rule" in which compounds generally incorporate into their carbon framework (1) a tertiary nitrogen with a small alkyl substituent; (2) a quaternary carbon; (3) a phenyl ring or its equivalent attached to the quaternary carbon; and (4) an ethyl linker between the quaternary carbon and the tertiary nitrogen. This complex architecture has served as the motivation for synthetic efforts spanning Gates' synthesis of morphine in 1952 to the work of Fukuyama in 2017, *vide infra*.

3.3. Natural Opiates.

By weight the opium poppy *P. somniferum* contains ca. 8–19% morphine (1), 1–5% codeine (2), and 1–5% thebaine (4). Morphine and codeine are pharmaceutical agents, with morphine serving as an analgesic and codeine as a cough suppressant. Contrarily, oripavine (3) and thebaine are starting materials for semisynthetic opiates.

3.3.1. Morphine.—First isolated by Sertürner in the early 1800s,²² morphine (1) was widely prescribed because it provides greater pain relief capacity than processed opium. Morphine is one of the most effective analgesics and is the standard against which new pain relievers are measured. Though the molecule's structure, first proposed by Robinson²³ and later confirmed by Gates (total synthesis)²⁴ and Hodgkin (X-ray crystallography),²⁵ has inspired several syntheses, there is currently no practical source of morphine, either by chemical synthesis or through fermentation, that can compete with isolation. Nevertheless, synthetic efforts toward morphine have pushed the field of organic synthesis forward and have been appropriately recounted.^{26–28} Consequently, this review would be remiss to attempt a full recapitulation of morphine's rich synthetic history; instead, we aim to paint a broad picture and would direct the reader to the literature for a more detailed exhibition.

The review begins with the Gates' total synthesis of morphine (Scheme 1a).^{24,29–32} Starting from dienophile **14**, cycloaddition with butadiene provided racemic Diels–Alder adduct (±)-**15** in moderate yield, best represented as its enol tautomer (±)-**16**.²⁹ Cu₂Cr₂O₅ mediated reductive hydrogenation was accompanied by cyclization to give (±)-**17**, closing what would ultimately emerge as the piperidine D ring. Six further transformations, including a chiral resolution with L- (+)-dibenzoyl tartaric acid, gave enantiopure tetracycle (–)-**18**. At this junction, two major obstacles remained: (1) inversion of the C-14 stereocenter and (2) closure of the dihydrofuran-like ring. Thus, treatment of (–)-**18** with two equivalents of Br₂ followed by the addition of 2,4-dinitrophenyl hydrazine (NH₂NHAr) produced an α,β -unsaturated hydrazone which underwent epimerization at C-14 to give the thermodynamically favored β -hydrogen substituted product. Hydrolysis of the hydrazone and chemoselective catalytic hydrogenation gave (–)-**19** which was exposed to excess Br₂ followed by 2,4-dinitrophenyl hydrazine to simultaneously close the E ring and install the requisite C7–C8 unsaturation. Hydrolysis of the hydrazone gave enone (–)-**20** which was converted to (–)-codeine (2) after reduction of the enone (in a 1,2 fashion) and aryl bromide. Finally, exposure to BBr₃ under modified Rapoport conditions gave (–)-morphine (1).

In 2014, the Gaunt lab completed a formal synthesis of (–)-morphine (Scheme 1b).³³ The effort began with a seven step sequence highlighted by a Noyori transfer hydrogenation to give chiral aldehyde (–)-**22** from isovanillin (21). A PhI(OAc)₂ mediated, intramolecular oxidative coupling was followed by Michael addition to give enone (+)-**24**. Next, the enone was elaborated to Boc-protected amine (–)-**25**. Luche reduction gave the corresponding allylic alcohol, which was heated under microwave irradiation with HCl to fragment the cyclic enol ether and deprotect the amine. Dehydrative ring closure, reductive amination, and carbamate protection gave (–)-**28**, completing the formal synthesis of (–)-morphine (1).

A third, noteworthy synthesis of morphine was reported by the Fukuyama lab in 2017 (Scheme 1c).³⁴ The synthesis began with the transformation of 7-methoxy-2-tetralone into

chiral bicycle (+)-**29** using the d'Angelo approach.³⁵ Elimination of the tertiary alcohol accompanied acid mediated ring closure. Next, hydroselenation of the resulting olefin and oxidative elimination of the selenoxide gave vinyl tertiary alcohol (+)-**30**. A retro-aldol/aldol sequence provided tetracycle **31** which was transformed into diene (–)-**32** in five further steps. At this point Diels–Alder cycloaddition with singlet oxygen gave endoperoxide **33** which then underwent selective fragmentation to enone (+)-**34**. Elimination of the tertiary alcohol was followed by amine liberation and subsequent 1,6-conjugate addition to give a 1:1.4 ratio of neopinone to codeinone which converged to pure codeinone after acid treatment. Reduction of the enone gave (–)-codeine (**2**) and finally (–)-morphine (**1**) after treatment with BBr₃.

3.3.2. Codeine.—Codeine (**2**) is prescribed for the relief of moderate pain and cough suppression. It has less analgesic ability than morphine and is usually taken orally. The recommended dose is 15–60 mg every 4–6 h, not exceeding 360 mg/day. Given its availability, codeine is commonly abused. When cough syrup is mixed with soft drinks, the brews are referred to as “lean” and “sizzurp”, among other names. Originating in Houston, Texas, lean gained popularity within southern hip-hop culture and has been associated with a number of deaths. Users risk overdose once consumption approaches 500 mg/day. Synthetically, codeine serves as a key intermediate in most morphine syntheses.

Our summary begins with Trost's 2002 synthesis of (–)-codeine (Scheme 2a).³⁶ Corey–Fuchs olefination of chiral cyanoaldehyde (–)-**35**³⁷ and regioselective dehalogenation gave (Z)-alkene (–)-**36**. Next, an intramolecular Heck reaction was used to close the B ring. SeO₂ oxidation of the resulting tetracycle (+)-**37** occurred preferentially at C₆ to give a mixture of alcohol and ketone (–)-**38** oxidation products that converged after a Dess–Martin periodinane oxidation. Next, the cyanoketone was reduced to the corresponding hydroimine. Subsequent amine formation occurred in the presence of methyl amine and reduction then gave (–)-**39**. Irradiation of precursor (–)-**39** and LDA in THF afforded (–)-codeine (**2**) by way of an intramolecular hydroamination. Irradiation was critical as treatment with LDA alone did not result in cyclization.

The Stork lab published a synthesis of (±)-codeine using a Diels–Alder cycloaddition to establish the B and C rings in a single move (Scheme 2b).³⁸ The synthesis began by exposing (*E*)-1-methoxybut-1-en-3-yne to Schwartz's reagent followed by the addition of aldehyde **41** to give (±)-**42**. Heating this compound in a sealed tube in the presence of Et₃N produced the Diels–Alder adduct (±)-**43** as a 4:1 mixture of diastereomers. After 10 steps, amine (±)-**44** underwent 6-exo-tet closure to give (±)-**45**. Protection of the amine was followed by oxidation and LiAlH₄ reduction to give (±)-codeine (**2**).

3.4. Licit Semisynthetic Opiates.

While codeine and morphine are important analgesics, a number of semisynthetic opiates are also important therapeutics.

3.4.1. Hydromorphone.—(–)-Hydromorphone (**5**, Dilaudid) is a potent semisynthetic opiate. Structurally, it is identical to morphine with additional oxygenation at the C₆

position. As a result, it is more water-soluble than morphine and displays increased efficacy (Table 2).³⁹

Recently, the Hudlicky group published a second generation, chemoenzymatic formal synthesis of ent-hydromorphone (Scheme 3).⁴⁰ Beginning with (2-bromoethyl)benzene (**46**), enzymatic dihydroxylation provided chiral diene (+)-**47** which could be elaborated to phenol (+)-**48** in seven additional steps. In a series of key transformations, oxidative phenol dearomatization was followed by Diels–Alder cycloaddition to establish the B and E rings in a single step. While Hudlicky's first generation synthesis used a Pb(OAc)₄ oxidation and subsequent heating to give the desired Diels–Alder adduct in 50% yield,⁴¹ the second generation approach replaced Pb(OAc)₄ with an iodine(III) oxidant. In the event, treatment of (+)-**48** with PhI(OAc)₂ gave ortho-quinone like intermediate **49**. After workup and heating in toluene **49** underwent intramolecular Diels–Alder cycloaddition to give (+)-**50**. Two challenges remained: (1) adjustment of the A ring phenol and the C₆ carbon to their proper oxidation states and (2) closure of the piperidine D ring. Rearomatization followed by addition of TFA provided a phenolic amine that was tosylated to give bis-tosylate (–)-**51**. At this stage, the first and second-generation syntheses intercepted, thus completing the formal synthesis of ent-hydromorphone (*ent*-5)

3.4.2. Hydrocodone.—Hydrocodone (7) is commonly found in combination with acetaminophen in the brand named analgesic Vicodin. Synthetically, it serves as an intermediate in the synthesis of codeine and morphine.^{42,43}

The survey begins with Rice's racemic effort which is regarded as the approach most amenable to industrial scale up (Scheme 4A).⁴⁴ The synthesis began with condensation of carboxylic acid **52** and amine **53**. The amide, thus obtained, was treated with POCl₃ to initiate a Friedel–Crafts like acylation event completed by the reduction of the intermediate imine.⁴⁵ Selective Birch reduction of the less electron rich C ring was followed by N-formylation to give formamide (±)-**55**. Methyl enol ether hydrolysis and concomitant ketalization set the stage for C₁ electrophilic aromatic bromination. Acidic hydrolysis of the acetal gave cyclization precursor (±)-**56**. Exposure of (±)-**56** to NH₄F·HF in TfOH provided cyclization product (±)-**57**. The C₁ bromide in the preceding step is critical to obtaining the desired cyclization product as previous studies indicate a high propensity for *para* cyclization at the expense of the desired *ortho* closure.^{46,47} Next, hydrolysis of (±)-**57** revealed the free secondary amine. α -bromination was followed by a 5-exo-tet ring closure to establish the E ring. Finally, reductive amination was accompanied by aryl bromide removal to give (±)-hydrocodone (7).

Mulzer and Trauner published a synthesis of (–)-hydrocodone (Scheme 4b).^{48,49} Their effort began with aldol condensation of (±)-**58** and methyl formate. After Robinson annulation with MVK, treatment with KOH cleaved the formate via a retro-Claisen reaction. Next, the racemic enone (±)-**59** was resolved using chiral chromatography to give (–)-**59**. 1,4-addition of a Grignard-copper(I) species was followed by trapping with TMSCl to give a transient silyl enol ether. This species was reacted with NBS to give α -bromo ketone (–)-**60**. Heating of (–)-**60** at 140 °C led to closure of the E ring. Following ketalization, a hydroboration, quenched with basic H₂O₂ installed a distal primary alcohol. Subsequent dechlorination with

Raney nickel gave intermediate (–)-**61**. After amine installation via Mitsunobu reaction with protected methylamine, a point of unsaturation was introduced via radical benzylic bromination and elimination. Dissolving metal reduction liberated the amine and set in motion reductive closure of the D ring. Ketal hydrolysis gave (–)-hydrocodone (**7**).

3.4.3. Oxycodone.—Oxycodone (**8**) is structurally identical to hydrocodone (**7**) sans C₁₄ oxygenation. Industrially, (**8**) is accessed through semisynthesis from (–)-thebaine (**4**) through a redox sequence (Scheme 5).^{50,51}

The Fukuyama lab reported the first total synthesis of oxycodone that circumvents thebaine as an intermediate (Scheme 6).⁵² Starting from aryl bromide **65**, a Fagnou palladium catalyzed direct acylation closed the B-ring to give tricycle **66**. A further three step sequence provided phenol **67**. In a key-step, oxidative dearomatization in the presence of PhI(OAc)₂ proceeded through oxonium intermediate **68** to give C₁₄ hydroxylated product **69**. Next, chemoselective hydrogenation with Wilkinson's catalyst provided enone **70**. At this stage, acylation of the C₁₄ alcohol with methyl malonyl chloride followed by exposure to Cs₂CO₃ resulted in intramolecular Michael addition to form the benzylic quaternary carbon. Subsequent Krapcho decarboxylation and deprotection generated phenol **71**. After five additional steps amide **72** was treated with PhI(OAc)₂ to induce an oxidative Hofmann rearrangement. The amine, resulting from aqueous hydrolysis of the intermediate isocyanate, ring opened the neighboring lactone to reveal the C₁₄ alcohol and establish the piperidine D ring. A final two step sequence converted **73** to (–)-oxycodone (**8**).

3.4.4. Buprenorphine and Naloxone.—Buprenorphine (**10**, Subutex) and naloxone (**9**, Narcan) are semisynthetic opiates used to treat drug dependence or overdose, respectively. Buprenorphine is a partial μ -receptor agonist often prescribed as an alternative to methadone to treat opioid dependence. It has less abuse potential than methadone, a full μ -receptor agonist,⁵³ and is also sold as a combination therapy with naloxone under the brand name Suboxone to further deter abuse. Naloxone is a μ -receptor antagonist that is United States Food and Drug Administration (FDA) approved for emergency treatment of known or suspected opioid and opiate overdose.⁵⁴ It is interesting to note that while naloxone is a receptor antagonist the majority of opiates are receptor agonists.

Synthetically, buprenorphine is made through semisynthesis from known opiate precursors. Starting from (–)-oripavine (**3**), amine alkylation and subsequent thiol mediated N-demethylation gave diene (–)-**74** (Scheme 7).⁵⁵ An endo selective Diels–Alder cycloaddition with MVK followed by Grignard addition and hydrogenation provide (–)-buprenorphine (**10**). The observed diastereoselectivity of the Grignard addition can be rationalized by invoking model (**75**) in which the oxygens of the carbonyl and proximal methyl ether coordinate to magnesium, locking the carbonyl in place for top-side attack.

Similarly, the Hudlicky group recently disclosed a semisynthesis of (–)-naloxone (**9**) from (–)-oxymorphone (**6**) (Scheme 8).⁵⁶ The synthesis commenced with acetate protection of the phenolic alcohol followed by amine oxidation. Next, dehydration with the Burgess reagent gave imine **77**. While disfavored, 5-endo-trig ring closure at this stage would provide oxazolidine **79**. One could argue that an alternative mechanism featuring a favored 5-exo-tet

cyclization (78) is more likely. In either case, oxazolidine 79 is next protected as its dimethyl acetal to allow for clean oxazolidine opening with a vinyl Grignard reagent. Hydrolysis of the acetal protecting group reveals (–)-naloxone (9).

3.5. Illicit Semisynthetic Opiates.

While most opiate abuse is enabled through prescriptions, some are illicitly synthesized for recreational use.

3.5.1. Heroin.—First synthesized from morphine in 1874, the Bayer Company of Germany introduced heroin for medical use in 1898. Physicians remained unaware of its addictive potential for years, and, in 1903, heroin abuse had risen to alarming levels in the United States. Heroin was made federally illegal in 1924. Later, the Counter-Narcotics Police of Afghanistan (CNPA) offered German authorities the opportunity to observe and document the illicit synthesis of heroin (Scheme 9).⁵⁷ The synthesis began with the isolation of morphine base from raw opium (82). Treatment of raw opium with lime (calcium oxide) and hot water resulted in a biphasic mixture in which the morphine base (1) resided in the lower aqueous layer. Siphoning of the aqueous layer into a fresh barrel followed by treatment with solid ammonium chloride resulted in the precipitation of morphine base. Next, the solid was treated with approximately 5 equiv of Ac_2O to convert morphine freebase to heroin. Addition of Na_2CO_3 neutralized the AcOH formed during the quench, precipitating brown heroin base in the process. Purification over activated carbon and precipitation with aq. NH_3 gave white heroin which was treated with HCl and acetone to give heroin hydrochloride ($11 \cdot \text{HCl}$).

3.5.2. Krokodil.—(–)-Desomorphine (12) or krokodil (from the Russian word meaning crocodile) became popular in Russia around 2003 and is a opioid synthesized from codeine and other readily obtained materials.⁵⁸ Over the past 15 years krokodil has slowly spread across Europe. It has also been sensationalized in a number of media reports as a drug that “turns the user into a zombie”. Indeed, krokodil can leave abusers disfigured via skin discoloration that results from cutaneous infection, gangrene, and necrosis.¹ The synthesis of krokodil begins with extraction of codeine containing tablets using alkaline pipe cleaner, gasoline, and HCl (Scheme 10).⁵⁹ The resulting hydrochloride salt ($2 \cdot \text{HCl}$) is then reduced with I_2 and red phosphorus through a Nagai Type reaction.

4. THE OPIOIDS

4.1. Historical Perspective.

The term opioids was originally coined in the 1960s by George H. Acheson to “refer to any chemical with morphine-like properties.” These compounds were originally synthesized to develop both less addictive and more powerful analgesics. From this research, compounds like fentanyl (84) emerged. While all of the opioids are addictive, some are larger targets for abuse such as Demerol (83) and methadone (85), while others are less abused due to their high potency and increased likelihood for overdose.

4.2. Structure.

Based on the morphine rule, opioids have similar (or stronger) effects as the opiates due to their masked structural homology, *vide supra*. Thus, each molecule features a tertiary amine linked to an aryl ring (Figure 2).

4.3. Demerol.

Demerol (83, also known as meperidine or pethidine) was first synthesized in 1939 and advertised as a “less addictive form of morphine” throughout the mid-1900s. However, a high level of dependence develops after initial use. Celebrities ranging from Elvis Presley to Michael Jackson have infamously suffered from Demerol addiction. In 1939, the Eisleb lab patented a synthesis of piperidine compounds, where di(β -halogenalkyl)-amine 87 is treated with benzyl cyanide. Cyclization is accompanied by nitrile hydrolysis to produce Demerol 83 (Scheme 11).⁶⁰ Many of the current industrial syntheses use this method.

While lacking the practicality of Eisleb, Hite’s 1959 synthesis, featuring a quasi-Favorskii rearrangement, provides an interesting alternative (Scheme 12).⁶¹ Starting with monochloroketone 89, available in six steps from isonicotinic acid, treatment with NaOH induced the desired rearrangement, albeit in modest yield as the reaction’s minor product. Further inspection reveals that either α -phenyl or α -hydroxyl migration can occur (transition states 90 and 92, respectively). Although not directly applicable, comparison of the A-values for the phenyl (3.0 kcal mol⁻¹) and hydroxyl (1.0 kcal mol⁻¹) substituents prove instructive in evaluating the relative amount of diaxial-like strain felt by each group when positioned antiperiplanar to the exiting chloride. Returning to the synthesis, Fischer esterification of 91 gives Demerol (83).

4.4. Fentanyl.

In 1953, Paul Janssen set out to develop a new analgesic that would improve upon morphine and Demerol.⁶² Hypothesizing that increased lipophilicity would increase CNS penetration and potency, Janssen and co-workers began modifying Demerol.⁶³ These efforts led to the synthesis of fentanyl (84) through a five step process involving reductive amination, acylation, and amine deprotection/alkylation (Scheme 13).⁶⁴ Though first brought to the market as an intravenous analgesic, fentanyl soon became a common anesthetic for cardiac and vascular surgeries. Later developments allowed for administration via transdermal patches, buccal lozenges (lollipops), and intranasal sprays.⁶³ Fentanyl is a synthetic analgesic that is 50–100 times more potent than morphine. Given its powerful pain-relieving effects, and its ease of synthesis, fentanyl is often mixed with heroin and other drugs. In many parts of the United States and Canada, fentanyl has all but replaced heroin, unbeknownst to the user.⁶⁵

4.5. Methadone.

In 1938, Bockmühl and Erhart synthesized compound VA 10820 that exhibited stronger analgesic effects than Demerol. Later named methadone (85), this compound is prescribed to treat opiate addiction. As methadone itself is highly addictive, many users cycle through the use of both opiates and opioids in an attempt to break dependence. The original synthesis of methadone (85) starts with racemic 1-dimethylamino-2-propanol (97) (Scheme 14).

Reaction with sodium diphenyl acetonitrile produces amino-nitrile ($\pm$)-99 via intermediate ($\pm$)-98. A Grignard addition concludes the synthesis to produce ($\pm$)-methadone (85).

5. THE BENZODIAZEPINES

5.1. Historical Perspective.

Research to expand the ring of quinazoline-3-oxides in the late 1950s led to the development of the benzodiazepines. Leo Sternbach investigated the heterocycles of benzheptoxdiazines, known in the German literature in the late 1800s as acylindazoles, in order to replace the use of the highly addictive barbiturates.⁶⁶ During his research, Sternbach made a library of compounds that were not assayed until a colleague rediscovered them ca. 2 years later. The molecules showed remarkable activity as sedatives and sparked a major development campaign into this new class of compounds.⁶⁷ A series of substituted benzodiazepines were synthesized and marketed as the antianxiety medications readily prescribed today. Others, like flunitrazepam (**104**), affect cognitive function and promote anterograde amnesia, leading to abuse as the “date rape” drug commonly known as a “Roofie”. After some time, it would be demonstrated that the benzodiazepines were as addictive as the barbiturates they hoped to replace. Nonetheless, the benzodiazepines offered a new chemical scaffold that medicinal chemists continue to manipulate today.

5.2. Structure.

Benzodiazepines are named according to their fusion of the benzene and diazepine ring systems (Figure 3). Benzodiazepine drugs are most commonly 1,4-substituted and can be differentiated based on modifications at the C_{2'}, C_{4'}, and C₇ positions.

5.3. Valium.

After the discovery of the first benzodiazepine (**100**, Librium), diazepam (**101**, Valium) was synthesized and demonstrated to be more potent. Valium would ultimately become the top selling drug in the United States from 1968 to 1982 but eventually received backlash as negative effects on cognition started to surface.⁶⁸ Beginning with *ortho*-aminobenzophenone (**108**), quinazoline-3-oxide (**105**) was generated in three steps.^{69,70} After treatment with NaOH, **105** underwent ring expansion to give the cyclic 7-membered lactam **107** which could be elaborated to Valium (**101**) in two steps (Scheme 15). Subsequent optimization reduced the synthesis from six steps to two (Scheme 16).^{66,68} This route began with condensation of *ortho*-aminobenzophenone (**108**) with glycine ethyl ester to establish the benzodiazepine ring system. Amide methylation completes the synthesis.

The Gates lab achieved the first academic synthesis of Valium in 1980 (Scheme 17).⁷¹ Isatoic anhydride **109** was converted to 1,4-benzodiazepine-2,5-dione **110** via nucleophilic attack of the amine and subsequent decarboxylation and cyclization. Acetylation and Grignard addition resulted in aminobenzophenone derivative **111** which yielded Valium (**101**) upon deacetylation and treatment with NaHSO₄.

5.4. Xanax.

The Upjohn Company patented the first synthesis of alprazolam (**103**, Xanax) in 1971 (Scheme 18a).^{72,73} Starting with quinoline derivative **112**, treatment with hydrazine gave the corresponding 2-hydrazinyl quinoline (not pictured) via an S_NAr (nucleophilic aromatic substitution) mechanism. Installation of the 1,2,4-triazole was realized after reaction with (EtO)₃CCH₃ to give **113**. Treatment with O₃ followed by oxidative workup with aq. NaI and AcOH gave ketone **114** in good yield. Mechanistically (Scheme 18b), one can envision that collapse of secondary ozonide (±)-**115** could occur after iodide attack to yield aldehyde **116** and an equivalent of hydroiodite (IO⁻). Reduction of this +1 iodine species could occur in situ under the influence of iodide and acid to give molecular iodine (I₂) which could in turn serve to oxidize **116**. After deprotonation of **117**, resonance stabilized acylium cation **118** is generated. Cation quenching with water would give carboxylic acid **120** which, following heterocycle protonation ((±)-**121**), can undergo decarboxylation to yield final product **114**. Returning to the synthesis, hydroxymethylation of **114** with paraformaldehyde, formation of the chloride, and subsequent displacement with NH₃ provided an amine that cyclizes after condensation to give Xanax (**103**).

6. THE STIMULANTS

6.1. Historical Perspective.

Stimulants promote wakefulness and decrease fatigue. Compounds in this category range from caffeine to cocaine, amphetamine, ephedrine, and MDMA (ecstasy).

6.2. Structure.

Cocaine (**122**) is a bicyclic tropinone alkaloid, which represents one of the more complex stimulant structures (Figure 4). Amphetamine, ephedrine, methamphetamine, and methylphenidate are classified as phenethylamines. MDMA is a phenethylamine derivative containing a 3,4-methylenedioxy bridge.

6.3. Cocaine.

The euphoric effects that result from chewing the leaves of the *Erythroxylum coca* plant have been known to humankind for millenia.⁷⁴ However, it was not until the mid-19th century that the coca-alkaloid cocaine (**122**) was isolated from this natural source.^{75,76} Soon after, cocaine gained popularity in the United States as a local anesthetic, a cure for opiate use disorder, and an additive to cigarettes and soda. At the turn of the century, cocaine's addictive nature emerged, and the United States responded with the 1914 Harrison Act to restrict the drug's sale and distribution. This largely ended the country's first cocaine epidemic.

From a synthetic perspective, cocaine's compact bicyclic structure and biological activity has long attracted attention. The Willstätter lab completed the first synthesis of (–)-cocaine (**122**) in 1903 (Scheme 20).^{77,78} This was a landmark moment in organic synthesis, as it involved the formation of the tropinone core and set a new precedent for the complexity that chemical synthesis could reach. The synthesis began with the transformation of cycloheptanone into cycloheptatriene (**130**). Treatment of **130** with Br₂ and (CH₃)₂NH

followed by chemoselective reduction provided unsaturated amine **131**. After dibromination of the remaining olefin, intramolecular bromide displacement provided bicyclic bromide salt ($\pm$)-**133**, presumably through a transition state similar to **132**. Bromide elimination and amine monodemethylation gave olefin($\pm$)-**134** which was subjected to bromination, hydrolysis, and oxidation to complete the synthesis of tropinone (**135**). Treatment of tropinone with Na in EtOH in the presence of CO₂ gave the α -carboxylic acid which was converted to methyl ester ($\pm$)-**136** under acidic conditions. Reduction with Na(Hg) and subsequent benzoylation afforded ($\pm$)-cocaine (**122**). A chiral resolution with D-tartaric acid allowed for the isolation of naturally occurring (–)-cocaine (**122**).

While trying to elucidate the biosynthetic pathways responsible for the synthesis of the tropinone scaffold, Sir Robert Robinson demonstrated a simplified synthesis of tropinone (**135**) in 1917 (Scheme 19).^{79,80} In a one-pot procedure, aqueous methylamine was added to a solution of succinaldehyde (**137**) and calcium acetonedicarboxylate (**138**). Following condensation of methylamine with aldehyde **137**, consecutive Mannich reactions gave tropinone (**135**). This reaction laid the groundwork for Robinson's proposal for the biosynthetic pathway of cocaine and similar alkaloids, an area of research that eventually earned him the Nobel Prize in Chemistry in 1947.

Unsurprisingly, cocaine's intricate bicyclic structure has continued to attract the interest of synthetic chemists long after the seminal efforts of Willstätter and Robinson. Rappaport and co-workers reported an enantioselective synthesis of (–)-cocaine (**122**) in 1998 (Scheme 21).⁸¹ Using the chiral pool as its source of asymmetry, the synthesis began with the Eschenmoser sulfide contraction of racemic dibenzyl malate (**139**) and D-glutamic acid derived chiral thiolactam (–)-**140**. Treatment of ($\pm$)-**139** with Tf₂O afforded the corresponding α -triflate which was displaced after addition of the thiolactam. Addition of PPh₃ as a sulfur scavenger followed by the introduction of a tertiary amine base completed the one pot reaction sequence to give vinyl amine (+)-**141** as an inconsequential mixture of *E* and *Z* isomers. Catalytic hydrogenation of (+)-**141** resulted in debenzylation and was accompanied by decarboxylation of the resulting vinylic acid carboxylic acid affording a chiral amino acid which, upon reprotection of the liberated carboxylic acid and amine functionalities, was transformed into (+)-**142**. KHMDS mediated Dieckmann cyclization gave bicyclic β -keto ester (+)-**143**. Nucleophilic decarboxylation followed by BamfordStevens reduction gave tropene intermediate (+)-**144** which underwent an *exo* selective [3 + 2] dipolar cycloaddition to give 4,5-dihydroisoxazole (–)-**145**. Ester hydrolysis followed by thermal decarboxylation and concomitant cleavage of the N–O bond gave β -hydroxyl cyanide (+)-**146** which was elaborated in six further steps to (–)-cocaine (**122**).

In regard to its illicit use, cocaine reemerged as a highly abused drug in the United States in the latter half of the 20th century. It is primarily consumed in one of two forms: powder cocaine and crack cocaine (Scheme 22). Powder cocaine, better chemically defined as the HCl salt of (–)-cocaine (**122**·HCl), is used via injection or insufflation. While this is the form in which cocaine is most commonly isolated, the free amine commonly known as crack cocaine is far more addictive. Crack cocaine reaches the brain quicker than the hydrochloride salt and produces a more intense high. The name “crack” is derived from the

sound this form of cocaine makes when burned. This form of cocaine is illicitly made from the cocaine salt (Scheme 22). In this process, powder cocaine (**122**-HCl) is treated with aq. NaHCO₃. After heating, the free base (**122**) separates from the water into an oil that can be solidified.

6.4. Phenethylamines.

The phenethylamines represent one of the most widely abused and illicitly produced classes of drugs today. The structural backbone of the class is seen in amphetamine (**123**), which features a single stereocenter and a key primary amine (Figure 4). Their structural similarity makes them easy to interconvert and alter. In this section, we describe general synthetic techniques used to produce these compounds and detail their illicit production.

6.4.1. Amphetamine.—Amphetamine (**123**) is a stimulant that increases wakefulness and focus while decreasing fatigue and appetite. Physicians often prescribe amphetamine as a 1:1:1:1 mixture, by weight, of four salts (Figure 5) to treat attention deficit hyperactivity disorder (ADHD). This mixture, sold under the brand name Adderall, and other stimulants prescribed to treat ADHD, such as (+)-lisdexamfetamine (Vyvanse, **124**) and (+)-methylphenidate (Focalin, Ritalin, Concerta, **128**), are often abused by college students as “study drugs”.

Synthetically, Edeleanu first prepared amphetamine (**123**) in 1887 and subsequently adapted the synthesis to access both ephedrine (**125**) and methamphetamine (**127**).⁸² However, interest in developing these compounds for use as drugs did not emerge until the early 1960s, at which point several syntheses made racemic amphetamine widely available. For example, in 1968, Belovsky and co-workers demonstrated a simple synthesis of (±)-amphetamine (**123**) from phenylacetone (**151**) and formamide (Scheme 23).⁸³ The synthesis began with the condensation of phenylacetone (**151**) and formamide under Leuckhart conditions to give (±)-**152**. Subsequent hydrolysis yielded (±)-amphetamine (**123**), completing the synthesis in two simple transformations. A variety of other traditional syntheses of racemic amphetamine exist and use key transformations such as the Ritter reaction, Friedel–Crafts acylation, Knoevenagel condensation, Curtius rearrangement and Wolff rearrangement.⁸⁴

While effective as a racemate, studies with enantiopure amphetamine have revealed that (+)-amphetamine demonstrates higher potency in the CNS while (–)-amphetamine has a larger effect on cardiovascular activity. As such, many have put forth efforts to access amphetamine in enantiopure form either through chiral resolution or enantioselective synthesis. These synthetic efforts engender chirality through the use of chiral starting materials, asymmetric hydrogenations, or enzymatic transformations.⁸⁴ Perhaps most prominent of these methods is Yamaguchi’s use of chiral aziridines (Scheme 24).⁸⁵ Beginning with chiral building block L-alinol (**153**), phosphonate protection followed by mesylation provided **154**. Next, this intermediate underwent base promoted intramolecular displacement to give chiral aziridine (**155**). Treatment of the aziridine with a mixture of PhMgBr and catalytic CuCl resulted in nucleophilic attack at the less substituted carbon.

Finally, acid mediated removal of the *N*-phosphonate revealed enantiopure (+)-amphetamine (**123**). A number of variations on this method have been reported.⁸⁶

6.4.2. (Pseudo)ephedrine.—Indiscriminate hydroxylation of the benzylic position of amphetamine yields two enantiomeric sets of diastereomers (Figure 6). The *erythro* diastereomers constitute the ephedrines (**125**) while the *threo* diastereomers comprise the pseudoephedrines (**126**). Medicinally, the *erythro* and *threo* diastereomers demonstrate different activities. Ephedrine stimulates the central nervous system, promoting wakefulness and decreased appetite, while pseudoephedrine acts on the peripheral nervous system and is used as a nasal decongestant sold under the brand name Sudafed.

6.4.3. Methamphetamine.—Mono-*N*-methylation of amphetamine (**123**) yields methamphetamine (**127**), a commonly abused drug with street names such as “meth” and “crank”. Due to the relative ease with which it can be synthesized, many clandestine laboratories manufacture this highly addictive drug. However, such synthetic work is not without danger as methamphetamine production frequently leads to “lab” explosions and house fires. The impact of such accidents as well as the socioeconomic ramification of methamphetamine addiction make plain the motivation behind the controlled nature of pseudoephedrine (**126**) containing medications, methylating reagents, and reducing agents.

Synthetically, methylation of amphetamine (**123**) with dimethyl sulfate can generate methamphetamine (**127**); however, the high propensity for these conditions to yield dimethylamphetamine makes this method less appealing.⁸⁷ The mono-*N*-methylated ephedrine (**125**) and pseudoephedrine (**126**) (jointly referred to as (pseudo)ephedrine) are more common starting materials as various routes exist for the reduction of (pseudo)ephedrine to methamphetamine (**127**). The most common strategy employs HI (Nagai route) or Birch reduction.

In recent years, a modification of the Birch reduction referred to as “shake and bake” has become the method of choice for illicit methamphetamine synthesis (Scheme 25). This method usually starts with (–)-pseudoephedrine containing decongestants. The pills are crushed and subsequently added to a plastic bottle. NH_4NO_3 , Li^0 , solid NaOH , and xylenes, chemicals available to the layperson from the household goods highlighted in red, are added. Next, iterative cycles of shaking and venting are used. After gas evolution ceases, the reaction is filtered and acidified with gaseous HCl to give methamphetamine (**127**) as its hydrochloride salt.

6.5. MDMA.

Commonly referred to as “ecstasy” or “Molly”, MDMA (**129**) is a strong stimulant that prevents synaptic reuptake of serotonin, dopamine, and norepinephrine. It is so effective that, when used repeatedly, it can deplete the number of neurotransmitters available in the brain, producing a mental and physical crash that results in long-term depression. MDMA also affects the temperature-regulating mechanisms of the brain with users commonly dying from hyperthermia and hyponatremia (depletion of sodium content). Merck patented the first commercial MDMA (**129**) synthesis in 1912 (Scheme 26).⁸⁸ Beginning with safrole (**156**),

Markovnikov hydrobromination of the terminal olefin gives the corresponding secondary bromide. Finkelstein reaction provides a secondary iodide that is displaced with methylamine to give (±)-MDMA (**129**).

The Shulze lab recently published an MDMA synthesis that features a Curtius rearrangement (Scheme 27).⁸⁹ Treatment of helional (**157**) with sodium propionate and propionic anhydride resulted in a Perkin reaction to give unsaturated carboxylic acid **158**. Olefin hydrogenation followed by acyl chloride synthesis gave (±)-**159**. At this point, generation of the acyl azide and subsequent Curtius rearrangement provided an isocyanate. Exposure of this intermediate to *t*-BuOK gave the corresponding Boc carbamate. N-Methylation followed by acid mediated Boc deprotection provided (±)-MDMA (**129**) which was isolated as its oxalate salt.

In the hands of the clandestine chemist, illicit MDMA synthesis often begins with distillation of safrole (**156**) from the more readily available sassafras oil (Scheme 28). Subsequent Wacker–Tsuji oxidation using PdCl₂ (available as a photograph developing agent) yields the ketone MDP2P (**161**). Subsequent reductive amination with methylamine utilizing any one of a variety of reducing agents (e.g., aluminum amalgam, available from the treatment of aluminum foil with a mercury(I) salt) provides (±)-MDMA (**129**). An interesting variation on this reaction bypasses the need for the highly regulated methylamine, instead using the more readily available nitromethane to generate methylamine in situ upon addition to the reaction's reducing environment.

7. THE HALLUCINOGENS

7.1. Historical Perspective.

Naturally occurring hallucinogens have been used for thousands of years as recreational drugs. Consumed for their spiritual and cognitive effects, many of these compounds structurally mimic neurotransmitters such as dopamine (**162**) and serotonin (**163**) (Figure 7) and have significant impacts on the CNS with users often claiming to have “spiritual awakenings” under their influence.

In response, a number of countries, have challenged their legality over the last century. In particular, during the mid-1960s counterculture movements in the United States, many Americans campaigned for the decriminalization of (+)-lysergic acid diethylamide (**164**, LSD).

7.2. Structure.

As noted above, many hallucinogenic compounds share structural homology with either dopamine (**162**) or serotonin (**163**) (Figures 7 and 8). This is most readily seen in the structures of mescaline (**167**) and *N,N*-dimethyltryptamine (**168**, herein referred to as DMT), respectively. However, careful examination of the structure of LSD (**164**) also reveals a resemblance to serotonin. On the other hand, ketamine (**165**) and phencyclidine (**166**, herein referred to as PCP) do not resemble either of the aforementioned neurotransmitters yet still elicit hallucinogenic effects.

7.3. LSD.

In the midst of a synthetic effort directed at accessing various amide analogues of lysergic acid, Albert Hofmann unwittingly constructed a derivative that would quickly permeate the world as a hallucinogen: LSD (**164**). However, it was not until 5 years later that Hofmann would accidentally ingest a small amount of LSD and discover the psychedelic properties of this semisynthetic ergot alkaloid.⁹⁰ While initially used by the medical community, LSD soon became a prominent illegal recreational drug.⁹¹ Structurally, LSD shares a tetracyclic skeleton with clinically relevant ergot alkaloids such as ergometrine (**170**) and ergotamine (**171**) (Figure 9).⁹²

Industrial access to clinically relevant ergot alkaloids relies on direct isolation from fermentation or field cultivation.^{93, 94} Similarly, one can imagine LSD resulting from the amination of lysergic acid (**169**) or arising from the hydrolysis of drugs such as ergometrine (**170**) or ergotamine (**171**). Consequently, the syntheses covered in the following section revolve around the synthesis of lysergic acid as a formal synthesis of LSD.

Woodward and co-workers published the first total synthesis of (±)-lysergic acid (**169**) in 1954 (Scheme 29).^{95,96} Beginning from known dihydroindole derivative (±)-**172**, acyl chlorination followed by AlCl₃ mediated Friedel–Crafts acylation closed the C ring to furnish (±)-**173**, a 2,2-*α*-dihydro derivative of Uhle's ketone. Ketone (±)-**173** was treated with pyridinium tribromide to afford the corresponding *α*-bromide which could be displaced with methylamino ketal **174**. Subsequent acid hydrolysis of the ketal and benzamide protecting groups revealed diketone (±)-**175** which, upon treatment with NaOMe, underwent aldol condensation to close the D ring and give (±)-**176**. Six subsequent transformations completed the first total synthesis of (±)-lysergic acid **169**.

Fukuyama's 2013 asymmetric synthesis of (+)-lysergic acid was equally instructive (Scheme 30).⁹⁷ Chiral *β*-hydroxyl hydrazide (–)-**177**, the product of Evans aldol condensation and subsequent auxiliary cleavage with hydrazine, was treated with *t*-BuONO to generate acyl azide **178**. In situ Curtius rearrangement afforded isocyanate **179** which underwent intramolecular trapping to produce oxazolidinone (–)-**180**. After three transformations, ring closing metathesis installed the A ring, giving (–)-**182**. Next, a Heck cyclization closed the B ring to give (–)-**184** after *β*-hydride elimination. Functional group interconversions over the course of 11 steps transformed this compound into the final product, (+)-lysergic acid (**169**). It is worth noting the authors' judicious control of the *β*-hydrogen stereochemistry in intermediate **183**; the syn positioning of this hydrogen and the palladium species enabled *β*-hydride elimination, demonstrating a firm command of mechanistic understanding.

In a similar vein, Fujii and Ohno published a 2011 formal synthesis of (+)-lysergic acid (**169**) featuring an amino-palladation cascade that established the C and D rings in a single step (Scheme 31).⁹⁸ Starting with enyne **186**, available in four steps from 4-bromoindole (**185**), Shi epoxidation utilizing catalyst **187** afforded epoxide **188** which was taken forward as a mixture of diastereomers. Silyl protection of the primary alcohol followed by Zn(OTf)₂ mediated reductive oxirane opening yielded ynol **189**. At this stage, the authors established the allene needed for the amino-palladation cascade via a Movassaghi modified Myers

allenation;⁹⁹ stereoselective formation of **192** occurred, presumably, through the concerted Alder-Ene reaction of intermediate **191**. With the allene in hand, TBAF deprotection of the primary alcohol and subsequent treatment with Pd(Ph₃)₄ and K₂CO₃, cyclization conditions previously established by the authors,¹⁰⁰ yielded tetracycle **194**. The amino-palladation precursor and the resulting product (**194**) intersected with a previously disclosed synthesis of (+)-lysergic acid by the authors.¹⁰⁰

7.4. Ketamine.

A molecule of many uses, ketamine (**165**) has long served as an anesthetic, analgesic, and recreational drug known on the street by the name “Special K.”¹⁰¹ ParkeDavis and Company disclosed the seminal synthesis of (±)-ketamine in a 1956 patent (Scheme 32).¹⁰² The synthesis began with *o*-chlorobenzonitrile (**195**) which yielded ketone **196** after treatment with a cyclopentyl Grignard reagent. Subsequent α -bromination followed by imine formation and bromide hydrolysis gave α -hydroxyl imine **197**, the precursor for the synthesis’ key pinacol-like rearrangement. In the event, heating **197** in decalin resulted in [1,2] migration to give (±)-ketamine (**165**).

More recently, Zhang and co-workers reported a racemic ketamine synthesis that started from ketone (±)-**198** (Scheme 33).¹⁰³ Direct α -nitration mediated by ceric ammonium nitrate (CAN) and Cu(OAc)₂ provided α -nitro ketone (±)-**199**. Nitro reduction and subsequent reductive amination with aqueous formaldehyde yielded (±)-ketamine (**165**).

While both of these syntheses provide facile access to (±)-ketamine, research has demonstrated that (–)-ketamine exhibits greater potency than the corresponding (+)-ketamine antipode in regards to anesthesia and analgesia.¹⁰⁴ As such, methods for accessing (–)-ketamine are gaining importance. Kiyooka recently published the first¹⁰⁵ and second¹⁰⁶ asymmetric syntheses of (–)-ketamine (**165**).

Kiyooka’s second synthesis begins with an enantioselective Mukaiyama aldol condensation mediated by chiral oxazaborolidinone **201** with α,β -unsaturated aldehyde (*P/M*)-**200** (Scheme 34). The resulting enantioenriched (86% ee) β -hydroxyl ester (*P/M*)-**202** was subjected to LiAlH₄ reduction and selective benzylation of the resulting primary alcohol. Acylation of the secondary alcohol with trichloroacetyl isocyanate and subsequent hydrolysis gave carbamate (*P/M*)-**203**. After dehydration, the carbamate was converted to isocyanate (–)-**205**. Mechanistically, this reaction occurs via dehydration of the carbamate to form intermediate allyl cyanate (*P/M*)-**204** which undergoes a stereoselective [3,3]-sigmatropic rearrangement giving the isocyanate product with 1,3-chirality transfer. Finally, reduction of the isocyanate (**205**) to give the corresponding methylamine was followed by reductive ozonolysis to give enantioenriched (–)-ketamine (**165**) in 87% ee.

7.5. PCP.

While Kötzt and Merkel reported the first synthesis of PCP (**166**) in 1926,¹⁰⁷ the compound was not fully investigated for its biological properties until the late 1950s when it was made available as an anesthetic (Sernyl).¹⁰⁸ Research demonstrated that the compound elicited the

desired anesthetic effects but was also a strong hallucinogen. This promptly led to its discontinued use in 1965. The hallucinogenic effects, however, fueled abuse of PCP.

Traditional syntheses of PCP are based on common cyclohexanone precursors, making them optimal for conversion into clandestine syntheses.^{109, 110} One commonly used procedure is exemplified by Parke-Davis and Company's 1965 synthesis of PCP (Scheme 35). Beginning from cyclohexanone (**206**), reaction with aq. NaHSO₄ provided an addition product that was treated with aqueous piperidine and KCN to give PCC (**207**). PhMgBr addition to the imine intermediate, presumably formed in situ from **207**, provides PCP (**166**).

An alternative synthesis from Pons began with conversion of tertiary alcohol **208** to the corresponding tertiary azide (Scheme 36).¹¹¹ Presumably, this substitution proceeds through an S_N1 manifold via a benzylic carbocation. Reduction of the azide provided a tertiary amine, which was elaborated to PCP (**166**) in good yield after dialkylation with 1,5-dibromopentane.

Clandestine syntheses of PCP often proceed according to routes similar to that depicted in Scheme 34 and can be performed on large scales. One such manifestation is termed the "The Bucket Method" (Scheme 37).¹⁰⁹ Following instructions seized from a clandestine PCP laboratory by the United States Drug Enforcement Administration (DEA), the starting materials are first divided into two buckets, Bucket A and Bucket B; to Bucket A are added cyclohexanone (**206**) and Na₂S₂O₅ (**209**, which can generate NaHSO₄ in situ when exposed to water), while piperidine (**210**) and an alkali cyanide salt are added to Bucket B. Addition of the contents of Bucket A to those of Bucket B (or vice versa) yields masked iminium species PCC (**207**). In a third bucket, Bucket C, bromobenzene (**211**) is added to magnesium turnings to generate PhMgBr. This Grignard reagent is then added to a solution of PCC in naphtha to give PCP (**166**). Because such processes can readily produce large batches of PCP, the DEA stringently monitors several related compounds, such as piperidine and potassium cyanide.

7.6. Mescaline.

Native Americans in Central and North America have used mescaline (**167**), the active component in the peyote cactus, for thousands of years in spiritual ceremonies. The "buttons" of the cactus are often dried and eaten or soaked in water to drink. Mescaline (**167**) is believed to mimic the neurotransmitter dopamine (**162**) (vide supra) and causes intense hallucinations and altered states of consciousness. Mescaline was first synthesized by Späth and co-workers in 1919 (Scheme 38).¹¹² Beginning with eudesmic acid (**212**), treatment with SOCl₂ gave the corresponding acyl chloride. Reduction under Rosenmund's conditions gave aldehyde **213** which could be elaborated to nitrostyrene **214** through a Henry reaction with nitromethane. With the molecule's carbon skeleton established, successive reductions with Zn in AcOH and then with Na(Hg) provided mescaline (**167**). In 1950, Morin and co-workers reported a similar synthesis using LiAlH₄ to convert the nitrostyrene intermediate **214** directly into the primary amine in higher yields.¹¹³

Tsao published a complementary mescaline synthesis in 1951 (Scheme 39).¹¹⁴ Beginning with eudesmic acid (**212**), a three step sequence of esterification, reduction, and chlorination

yielded benzyl chloride **215**. Addition of the remaining aminomethylene unit was accomplished through Kolbe nitrile synthesis and LiAlH_4 reduction.

7.7. DMT.

An entheogen is any psychoactive substance that induces a spiritual experience. Ayahuasca is a spiritual medicinal brew traditionally used in ceremonies by peoples of the Amazon river basin. DMT (**168**) is the active component in ayahuasca.¹¹⁵ Its biosynthesis occurs within plants via the shikimate pathway.¹¹⁶ This pathway is also responsible for the biosynthesis of the amino acid tryptophan as well as various indole alkaloid precursors. The direct precursor of DMT is tryptamine (**216**), which was first chemically synthesized by Shapiro and co-workers in 1956 (Scheme 40).¹¹⁷ The synthesis begins with an intermolecular Michael addition between diethyl malonate (**217**) and acrylonitrile (**218**) to give a product that cyclizes after Raney nickel reduction to yield ($\pm$)-**219**. A Japp–Klingemann reaction with benzenediazonium chloride gives hydrazine **220** which undergoes cyclization when exposed to polyphosphoric acid to give ($\pm$)-**221**. Amide hydrolysis and subsequent decarboxylation yielded tryptamine (**216**).

Conversion of tryptamine (**216**) to DMT (**168**) occurs under standard methylating conditions or through reductive amination (Scheme 41).¹¹⁸

8. CONCLUSION

A number of key lessons were learned during the assembly of this document. First, the field of psychoactive drug synthesis is rich with foundational strategies and tactics to prepare both polycyclic and linear alkaloids. While the seminal Gates synthesis was completed 65 years ago, the field remains vibrant with countless players distributed worldwide. Academic syntheses, however, are primarily instructional. Even in an era where synthetic approaches are arguably the shortest they have ever been, the use of total synthesis in matching the efficiency of direct isolation or semisynthesis is futile (at least in the case of the complex opium or coca alkaloids). Perhaps in time an “ideal” four to six-step synthesis will emerge that challenges plant isolation.

Second, it is clear that the early exercises of our synthetic education are showcased throughout psychoactive drug production. Whether it is the ability to produce a 5- or 6-membered ring (the latter via Diels–Alder technology) or introduce tertiary amines through reductive amination, basic organic chemistry maneuvers are critical to the production of psychoactive drugs, be it in the lab or in the garage.

A final point is rather distal in nature and focuses on the future of psychoactive drug production. Currently, the United States is overwhelmed by opioid use disorder. Since 2000, benzodiazepine, painkiller, and heroin deaths have all risen to record levels. While there are a number of social strategies in place to combat the problem, perhaps the solution lies at the hands of synthetic chemists and our ability to develop compounds that address pain and suffering while minimizing addiction.

ACKNOWLEDGMENTS

We'd like to thank Dr. Katherine Chong for critical feedback and the reviewers for their suggestions.

Funding

S.D.T. would like to acknowledge Vanderbilt University and the Institute of Chemical Biology for financial support. S.A.C. acknowledges support from the Vanderbilt Chemical Biology Interface (CBI) training program (T32 GM065086). E.D.H. acknowledges support from the Vanderbilt Chemical Biology of Infectious Diseases (CBID) training program (T32 AI112541).

ABBREVIATIONS

BSNR	based on starting material not recovered
AcOH	acetic acid
Br₂	bromine
Cu₂CrO₅	copper chromite
CuCl	copper chloride
LiAlH₄	lithium aluminum hydride
BBr₃	boron tribromide
PhI(OAc)₂	(diacetoxyiodo)benzene
HCl	hydro-chloric acid
TFA	trifluoroacetic acid
Cs₂CO₃	cesium carbonate
Et₃N	triethylamine
SeO₂	selenium dioxide
NaBH₄	sodium borohydride
Pb(OAc)₄	lead tetraacetate
POCl₃	phosphoryl chloride
NaBH₃CN	sodium cyanoborohy-dride
Tf₂O	triflic anhydride
NH₄F·HF	ammonium fluoride
KOH	potassium hydroxide
NaOH	sodium hydroxide
TMSCl	chlorotrimethylsilane

Ac₂O	acetic anhydride
NBS	N-bromosuccinimide
H₂O₂	hydrogen peroxide
MVK	methyl vinyl ketone
MOM	methoxymethyl
Na₂CO₃	sodium carbonate
NaHSO₄	sodium bisulfate
O₃	ozone
NH₃	ammonia
aq	aqueous
(EtO)₃CCH₃	triethyl orthoformate
(CH₃O)₂SO₂	dimethyl sulfate
PhMgBr	phenylmagnesium bromide
NH₄NO₃	ammonium nitrate
Li⁰	lithium zero
t-BuOK	potassium t-butoxide
NaI	sodium iodide
I₂	iodine
Na⁰	sodium zero
EtOH	ethanol
CO₂	carbon dioxide
Na(Hg)	sodium amalgam
PPh₃	triphenyl phosphine
KHMDS	potassium bis(trimethylsilyl)amide
NaHCO₃	sodium bicarbonate
SOCl₂	thionyl chloride
AlCl₃	aluminum trichloride
NaOMe	sodium methoxide
Zn(OTf)₂	zinc(II)-triflate

TBAF	tetrabutylammonium fluoride
K₂CO₃	potassium carbonate
KCN	potassium cyanide

REFERENCES

- (1). Fava GA (1999) The practitioner's guide to psychoactive drugs, edition 4. Psychother. Psychosom 68, 226–226.
- (2). McCann UD, Lowe KA, and Ricaurte GA (1997) Long-lasting effects of recreational drugs of abuse on the central nervous system. Neuroscientist 3, 399–411.
- (3). Saal D, Dong Y, Bonci A, and Malenka RC (2003) Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. Neuron 37, 577–582. [PubMed: 12597856]
- (4). Carlsson A (1967) Basic Actions of Psychoactive Drugs. Int. J. Neurol 6, 27. [PubMed: 4392241]
- (5). Qazi S, Caberlin M, and Nigam N (2007) Mechanism of psychoactive drug action in the brain: Simulation modeling of GABA(A) receptor interactions at non-equilibrium conditions. Curr. Pharm. Des 13, 1437–1455. [PubMed: 17504166]
- (6). (2016) In Facing Addiction in America: The Surgeon General's Report on Alcohol, Drugs, and Health, Washington, DC.
- (7). Center for Behavioral Health Statistics and Quality (2016) Key substance use and mental health indicators in the United States: Results from the 2015 National Survey on Drug Use and Health, HHS Publication No. SMA 16–4984, NSDUH Series H-51, SAMHSA, Rockville, MD.
- (8). Reitan T (2016) Commitment without confinement. Outpatient compulsory care for substance abuse, and severe mental disorder in Sweden. Int. J. Law Psychiatry 45, 60–69. [PubMed: 26912456]
- (9). Van Zee A (2009) The Promotion and Marketing of OxyContin: Commercial Triumph, Public Health Tragedy. Am. J. Public Health 99, 221–227. [PubMed: 18799767]
- (10). Mejia D, and Restrepo P (2016) The economics of the war on illegal drug production and trafficking. J. Econ Behav Organ 126, 255–275.
- (11). Hong CY, Kado N, and Overman LE (1993) Asymmetric synthesis of either enantiomer of opium alkaloids and morphinans. Total synthesis of (–)- and (+)-dihydrocodeinone and (–)- and (+)-morphine. J. Am. Chem. Soc 115, 11028–11029.
- (12). Magnus P, Sane N, Fauber BP, and Lynch V (2009) Concise syntheses of (–)-galanthamine and (±)-codeine via intra-molecular alkylation of a phenol derivative. J. Am. Chem. Soc 131, 16045–16047. [PubMed: 19835379]
- (13). Parker KA, and Fokas D (2006) Enantioselective synthesis of (–)-dihydrocodeinone: a short formal synthesis of (–)-morphine. J. Org. Chem 71, 449–455. [PubMed: 16408951]
- (14). Blakemore PR, and White JD (2002) Morphine, the Proteus of organic molecules. Chem. Commun. (Cambridge, U. K.), 1159–1168.
- (15). Reed JW, and Hudlicky T (2015) The quest for a practical synthesis of morphine alkaloids and their derivatives by chemo-enzymatic methods. Acc. Chem. Res 48, 674–687. [PubMed: 25730681]
- (16). Rinner U, and Hudlicky T (2011) Synthesis of morphine alkaloids and derivatives. Top. Curr. Chem 309, 33–66.
- (17). Chida N (2010) Recent advances in the synthesis of morphine and related alkaloids. Top. Curr. Chem 299, 1–28.
- (18). Zografos AL (2016) From biosynthesis to total synthesis: strategies and tactics for natural products, Wiley, Hoboken, NJ.
- (19). Macht DI (1915) The history of opium and some of its preparations and alkaloids. JAMA 64, 477–481.
- (20). Lovell J (2015) The Opium War: Drugs, Dreams, and the Making of Modern China, The Overlook Press, New York.

- (21). Wikler A, Martin WR, Pescor FT, and Eades CG (1963) Factors regulating oral consumption of an opioid (Etonitazene) by morphine-addicted rats. *Psychopharmacologia* 5, 55–76. [PubMed: 14082382]
- (22). Sertürner FW (1817) Über das Morphinum, eine neue salzfähige Grundlage, und die Mekonsäure, als Hauptbestandteile des Opiums. *Ann. Phys* 55, 56–89.
- (23). Gulland JM, and Robinson R (1925) Constitution of Codeine and Thebaine. *Mem. Proc. Manchester Lit. Philos. Soc* 69, 79–86.
- (24). Gates M, and Tschudi G (1952) The Synthesis of Morphine. *J. Am. Chem. Soc* 74, 1109–1110.
- (25). Mackay M, and Hodgkin DC (1955) A crystallographic examination of the structure of morphine. *J. Chem. Soc*, 3261.
- (26). Li Q, and Zhang H (2017) Research Progress on the Synthesis of Morphine Alkaloids. *Youji Huaxue* 37, 1629–1652.
- (27). Goto K (1949) Advances in the Syntheses of Morphine Alkaloids. *Yakugaku Zasshi* 69, 107–113.
- (28). Rinner U, and Hudlicky T (2011) Synthesis of morphine alkaloids and derivatives. *Top. Curr. Chem* 309, 33–66.
- (29). Gates M (1950) The Synthesis of Ring Systems Related to Morphine. III. 5,6-Dimethoxy-4-cyanomethyl-1,2-naphthoquinone and its Condensation with Dienes. *J. Am. Chem. Soc* 72, 228–234.
- (30). Gates M, and Tschudi G (1956) The Synthesis of Morphine. *J. Am. Chem. Soc* 78, 1380–1393.
- (31). Gates M, Woodward RB, Newhall WF, and Knzli R (1950) The Synthesis of Ring Systems Related to Morphine. IV. N-Methylisomorphinan*. *J. Am. Chem. Soc* 72, 1141–1146.
- (32). Gates M, and Tschudi G (1950) The Synthesis of Racemic β -6-Dihydrodesoxycodine Methyl Ether. *J. Am. Chem. Soc* 72, 4839–4840.
- (33). Tissot M, Phipps RJ, Lucas C, Leon RM, Pace RD, Ngouansavanh T, and Gaunt MJ (2014) Gram-scale enantioselective formal synthesis of morphine through an ortho-para oxidative phenolic coupling strategy. *Angew. Chem., Int. Ed* 53, 13498–13501.
- (34). Umihara H, Yokoshima S, Inoue M, and Fukuyama T (2017) Total Synthesis of (–)-Morphine. *Chem. - Eur. J* 23, 6993–6995. [PubMed: 28378531]
- (35). Sdassi H, Revial G, Pfau M, and d'Angelo J (1990) Enantioselective approach to morphinans. *Tetrahedron Lett* 31, 875–878.
- (36). Trost BM, and Tang W (2002) Enantioselective synthesis of (–)-codeine and (–)-morphine. *J. Am. Chem. Soc* 124, 14542–14543. [PubMed: 12465957]
- (37). Trost BM, and Tang W (2002) An Efficient Enantioselective Synthesis of (–)-Galanthamine. *Angew. Chem., Int. Ed* 41, 2795–2797.
- (38). Stork G, Yamashita A, Adams J, Schulte GR, Chesworth R, Miyazaki Y, and Farmer JJ (2009) Regiospecific and stereoselective syntheses of ($\pm$) morphine, codeine, and thebaine via a highly stereocontrolled intramolecular 4 + 2 cycloaddition leading to a phenanthrofurane system. *J. Am. Chem. Soc* 131, 11402–11406. [PubMed: 19624126]
- (39). Yan C, and Theodorescu D (2015) One step closer to targeting RAS. *Cell Cycle* 14, 287–288. [PubMed: 25591050]
- (40). Rycek L, Hayward JJ, Latif MA, Tanko J, Simionescu R, and Hudlicky T (2016) Chemoenzymatic Total Synthesis of Hydromorphone by an Oxidative Dearomatization/Intramolecular [4+2] Cycloaddition Sequence: A Second-Generation Approach. *J. Org. Chem* 81, 10930. [PubMed: 27690149]
- (41). Varghese V, and Hudlicky T (2014) Short Chemoenzymatic Total Synthesis of (–)-Hydromorphone: An Oxidative Dearomatization/Intramolecular [4 + 2] Cycloaddition/Amination Sequence. *Angew. Chem., Int. Ed* 53, 4355–4358.
- (42). Iijima I, Rice KC, and Silverton JV (1977) Studies in the (+)-Morphinan Series I. An Alternative Conversion of (+)-Dihydrocodeinone into (+)-Codeine. *Heterocycles* 6, 1157–1165.
- (43). Rice KC (1977) A rapid, high-yield conversion of codeine to morphine. *J. Med. Chem* 20, 164–165. [PubMed: 833816]

- (44). Rice KC (1980) Synthetic Opium Alkaloids and Derivatives. A Short Total Synthesis of (A)-Dihydrothebainone, (f)-Dihydroco-deinone, and (*)-Nordihydrocodeinone as an Approach to a Practical Synthesis of Morphine, Codeine, and Congeners. *J. Org. Chem* 45, 3135–3137.
- (45). Rice KC, and Bossi A (1980) Expedient Synthesis of Racemic and Optically Active N-Norreticuline and N-Substituted and 6'-Bromo-N-norreticuline. *J. Org. Chem* 45, 592–601.
- (46). Grewe R, and Friedrichsen W (1967) Die Cyclisierung von Octahydro-isochinolinderivaten durch Morphinan-Ringschluss. Synthese des Dihydrothebainons. *Chem. Ber* 100, 1550–1558. [PubMed: 5593839]
- (47). Morrison GC, Waite RO, and Shavel JJ (1967) An Alternate Route in the Synthesis of Morphine. *Tetrahedron Lett* 8, 4055–4056.
- (48). Mulzer J, Duerner G, and Trauner D (1996) Formal Total Synthesis of (–)-Morphine. *Angew. Chem., Int. Ed. Engl* 35, 2830–2832.
- (49). Mulzer J, and Trauner D (1999) Practical Synthesis of (–)-Morphine. *Chirality* 11, 475–482.
- (50). Kok GB, and Scammells PJ (2012) Improved synthesis of 14-hydroxy opioid pharmaceuticals and intermediates. *RSC Adv* 2, 11318–11325.
- (51). Krassnig R, Hederer C, and Schmidhammer H (1996) Optimization of the synthesis of oxycodone and 5-methyloxycodone. *Arch. Pharm* 329, 325–326.
- (52). Kimishima A, Umihara H, Mizoguchi A, Yokoshima S, and Fukuyama T (2014) Synthesis of (–)-Oxycodone. *Org. Lett* 16, 6244–6247. [PubMed: 25423610]
- (53). Whelan PJ, and Remski K (2012) Buprenorphine vs methadone treatment: A review of evidence in both developed and developing worlds. *J. Neurosci Rural Pract* 3, 45–50. [PubMed: 22346191]
- (54). Kerensky T, and Walley AY (2017) Opioid overdose prevention and naloxone rescue kits: what we know and what we don't know. *Addict. Sci. Clin. Pract* 12, 4. [PubMed: 28061909]
- (55). Theodor WB, and Roux L (2016) Improved method of manufacturing buprenorphine and analogues thereof from oripavine, Patent WO 2016/078833A1.
- (56). Endoma-Arias MAA, Cox DP, and Hudlicky T (2013) General Method of Synthesis for Naloxone, Naltrexone, Nalbuphine, and Nalbuphine by the Reaction of Grignard Reagents with an Oxazolidine Derived from Oxymorphone. *Adv. Synth. Catal* 355, 1869–1873.
- (57). Zerell U, Ahrens B, and Gerz P (2005) Documentation of a heroin manufacturing process in Afghanistan. *Bull. Narc* 57, 11–31. [PubMed: 21338014]
- (58). Shelton M, Ramirez-Fort MK, Lee KC, and Ladizinski B (2015) Krokodil: from Russia with love. *JAMA Dermatol* 151, 32. [PubMed: 25587682]
- (59). Alves EA, Soares JX, Afonso CM, Grund JP, Agonia AS, Cravo SM, Netto AD, Carvalho F, and Dinis-Oliveira RJ (2015) The harmful chemistry behind "krokodil": Street-like synthesis and product analysis. *Forensic Sci. Int* 257, 76–82. [PubMed: 26282512]
- (60). Otto E (1939) Piperidine compounds and a process of preparing them, US Patent US 2167351A.
- (61). Smissman EE, and Hite G (1959) The Quasi-Favorskii Rearrangement. I. The Preparation of Demerol and β -Pethidine. *J. Am. Chem. Soc* 81, 1201–1203.
- (62). Black J (2005) A Personal Perspective on Dr. Paul Janssen. *J. Med. Chem* 48, 1687–1688. [PubMed: 15771410]
- (63). Stanley TH (2014) The fentanyl story. *J. Pain* 15, 1215–1226. [PubMed: 25441689]
- (64). Janssen PAJ (1965) 1-Aralkyl-4-(n-aryl-carbonyl amino)-piperidines and related compounds, US Patent US 3164600A.
- (65). Aschenbrenner DS (2017) Fentanyl's Role in Opioid Overdose Deaths. *Am. J. Nurs* 117, 21–23.
- (66). Sternbach LH (1971) 1,4-benzodiazepines. Chemistry and some aspects of the structure-activity relationship. *Angew. Chem., Int. Ed. Engl* 10, 34–43. [PubMed: 4993720]
- (67). Sternbach LH (1979) The benzodiazepine story. *J. Med. Chem* 22, 1–7. [PubMed: 34039]
- (68). Calcaterra NE, and Barrow JC (2014) Classics in chemical neuroscience: diazepam (valium). *ACS Chem. Neurosci* 5, 253–260. [PubMed: 24552479]
- (69). Stempel A, Reeder E, and Sternbach LH (1965) Quinazolines and 1,4-Benzodiazepines. XXVII. Mechanism of Ring Enlargement of Quinazoline 3-Oxides with Alkali to 1,4-Benzodiazepine-2-1 4-Oxides. *J. Org. Chem* 30, 4267–4271. [PubMed: 4158669]
- (70). Sternbach LH (1959) 1,4-Benzodiazepine 4-Oxides, US Patent US2893992

- (71). Gates M (1980) New Synthesis of Diazepam. *J. Org. Chem* 45, 1675–16781.
- (72). Hester J (1973) Process for the production of triazolobenzo-diazepines and intermediates, US Patent US3709898A
- (73). Hester J (1973) 7-Chloro-1-methyl-5-phenyl-s-triazolo (4,3-a)quinolines, US Patent US3781289A.
- (74). Das G (1993) Cocaine abuse in North America: a milestone in history. *J. Clin. Pharmacol* 33, 296–310. [PubMed: 8473543]
- (75). Niemann A (1860) Ueber eine neue organische Base in den Cocablättern. *Arch. Pharm* 153, 129–155.
- (76). Gaedcke F (1855) Ueber das Erythroxylin, dargestellt aus den Blättern des in Siidamerika. kultivirten Strauches, Erythroxyton coca, Lam. *Arch. Pharm* 132, 141–150.
- (77). Willstätter R (1901) Synthesen in der Tropicgruppe. I. Synthese des Tropilidens. *Justus Liebigs Ann. Chem* 317, 204–265.
- (78). Humphrey AJ, and O'Hagan D (2001) Tropane alkaloid biosynthesis. A century old problem unresolved. *Nat. Prod. Rep* 18, 494–502. [PubMed: 11699882]
- (79). Robinson R (1917) LXIII. A synthesis of tropinone. *J. Chem. Soc., Trans* 111, 762–768.
- (80). Robinson R (1955) *The Structural Relations of Natural Products*, Clarendon Press, Oxford.
- (81). Lin R, Castells J, and Rapoport H (1998) Enantiospecific Synthesis of Natural (–)-Cocaine and Unnatural (+)-Cocaine from D-and L-Glutamic Acid. *J. Org. Chem* 63, 4069–4078.
- (82). Edeleano L (1887) Über einige derivate der Phenyl-methacrylsäure und der Phenylisobuttersäure. *Ber. Dtsch. Chem. Ges* 20, 616–622.
- Cervinka O, Kroupova E, and Belovsky O (1968) Asymmetric reactions. XXIX. Absolute configuration of ω -phenyl-2-alkylamines and their N-methyl derivatives. *Collect. Czech. Chem. Commun* 33, 3551–3557.
- (84). Allen A, and Ely R Synthetic Methods for Amphetamine A review, <http://www.nwafs.org/newsletters/SyntheticAmphetamine.pdf>.
- Manaka TNS, Desadee W, Yajima N, Kumamoto T, Watanabe T, Ishikawa T, Kawahata M, and Yamaguchi K (2007) Ring-Opening Reactions of 3-Aryl-1-benzylaziridine-2-carboxylates and Application to the Asymmetric Synthesis of an Amphetamine-Type Compound. *Helv. Chim. Acta* 90, 128–142.
- Meckler H, Gregg BT, and Yang J (2015) Synthesis of Chiral Amphetamine Derivatives by Stereospecific, Regioselective Cuprate Addition Reaction with Aziridine Phosphoramidate Compounds, US Patent 20150183810.
- (87). Kiefer EF (1972) A Rapid, Convenient Preparative Procedure for Phenethylamines. *J. Med. Chem* 15, 214.
- (88). Merck E (1914) Verfahren zur darstellung von alkyloxyaryl-, dialkyloxyaryl- und alkylenedioxyarylamino-propanen bzw. deren am stickstoff monoalkylierten derivaten, German patent 274,350.
- (89). Schulze M (2010) Synthesis of 2-Arylethylamines by the Curtius Rearrangement. *Synth. Commun* 40, 1461–1476.
- (90). Hofmann A (1996) LSD: Completely Personal. *MAPS*, 6.
- (91). Passie T, Halpern JH, Stichtenoth DO, Emrich HM, and Hintzen A (2008) The pharmacology of lysergic acid diethylamide: a review. *CNS Neurosci. Ther* 14, 295–314. [PubMed: 19040555]
- (92). Liu H, and Jia Y (2017) Ergot alkaloids: synthetic approaches to lysergic acid and clavine alkaloids. *Nat. Prod. Rep* 34, 411–432. [PubMed: 28300233]
- (93). Hanosová H, Koprna R, Valík J, Knoppová L, Fiebort I, Dzurová L, and Galuszka P. (2015) Improving field production of ergot alkaloids by application of gametocide on rye host plants. *New Biotechnol* 32, 739–746.
- (94). Schiff PLJ (2006) Ergot and Its Alkaloids. *Am. J. Pharm. Educ* 70, 98. [PubMed: 17149427]
- (95). Kornfeld EC, Fornfeld EJ, Kline GB, Mann MJ, Jones RG, and Woodward RB (1954) The Total Synthesis of Lysergic Acid and Engrovine. *J. Am. Chem. Soc* 76, 5256–5257.
- (96). Kornfeld EC, Fornfeld EJ, Kline GB, Mann MJ, Morrison DE, Jones RG, and Woodward RB (1956) The Total Synthesis of Lysergic Acid. *J. Am. Chem. Soc* 78, 3087–3114.

- (97). Liu Q, Zhang YA, Xu P, and Jia Y (2013) Total synthesis of (+)-lysergic acid. *J. Org. Chem* 78, 10885–10893. [PubMed: 24111583]
- (98). Iwata A, Inuki S, Oishi S, Fujii N, and Ohno H (2011) Formal total synthesis of (+)-lysergic acid via zinc(II)-mediated regioselective ring-opening reduction of 2-alkynyl-3-indolyloxirane. *J. Org. Chem* 76, 5506–5512. [PubMed: 21598967]
- (99). Movassaghi M, and Ahmad OK (2007) N-Isopropylidene-N'-2-nitrobenzenesulfonyl Hydrazine, a Reagent for Reduction of Alcohols via the Corresponding Monoalkyl Diazenes. *J. Org. Chem* 72, 1831–1841.
- (100). Inuki S, Iwata A, Oishi S, Fujii N, and Ohno H (2011) Enantioselective total synthesis of (+)-lysergic acid, (+)-lysergol, and (+)-isolysergol by palladium-catalyzed domino cyclization of allenes bearing amino and bromoindolyl groups. *J. Org. Chem* 76, 2072–2083. [PubMed: 21361331]
- (101). Tyler MW, Yourish HB, Ionescu DF, and Haggarty SJ (2017) Classics in Chemical Neuroscience: Ketamine. *ACS Chem. Neurosci* 8, 1122–1134. [PubMed: 28418641]
- (102). Stevens CL (1966) Aminoketones and Methods For Their Production, US Patent US3254124.
- (103). Zhang ZQ, Chen T, and Zhang FM (2017) Copper-Assisted Direct Nitration of Cyclic Ketones with Ceric Ammonium Nitrate for the Synthesis of Tertiary alpha-Nitro-alpha-substituted Scaffolds. *Org. Lett* 19, 1124–1127. [PubMed: 28206766]
- (104). Weiskopf RB, Nau C, and Strichartz GR (2002) Drug Chirality in Anesthesia. *Anesthesiology* 97, 497–502. [PubMed: 12151942]
- (105). Yokoyama R, Matsumoto S, Nomura S, Higaki T, Yokoyama T, and Kiyooka S (2009) Enantioselective construction of nitrogen-substituted quaternary carbon centers adjacent to the carbonyl group in the cyclohexane ring: first asymmetric synthesis of anesthetic (S)-ketamine with high selectivity. *Tetrahedron* 65, 5181–5191.
- (106). Yokoyama T, Yokoyama R, Nomura S, Matsumoto S, Fujiyama R, and Kiyooka S-i. (2009) Synthesis of (S)-Ketamine via [1,3]-Chirality Transfer of a Stereocenter Created by Enantioselective Aldol Reaction. *Bull. Chem. Soc. Jpn* 82, 1528–1532.
- (107). Kotz A, and Merkel P (1926) Hydromatic alkalines. *J. Prakt. Chem* 113, 49–76.
- (108). Shulgin AT, and Mac Lean DE (1976) Illicit synthesis of phencyclidine (PCP) and several of its analogs. *Clin. Toxicol* 9, 553–560. [PubMed: 975751]
- (109). Allen AC, Robles J, Dovenski W, and Calderon S (1993) PCP: A review of synthetic methods for forensic clandestine investigation. *Forensic Sci. Int* 61, 85–100. [PubMed: 8307527]
- (110). Maddox VH, Godefroi EF, and Parcell RF (1965) The Synthesis of Phencyclidine and Other 1-Arylcyclohexylamines. *J. Med. Chem* 8, 230–235. [PubMed: 14332667]
- (111). Geneste P, Herrmann P, Kamenka JM, and Pons A (1975) Nouvelle voie d'accès aux isomères des phenyl-1-cyclohexyl-amines substitués au cyclohexane. *Bull. Soc. Chim. Fr* 7–8, 1619–1626.
- (112). Spath E (1919) Über die Anhalonium-Alkaloide: I. Anhalin und Mezcalin. *Monatsh. Chem* 40, 129–154.
- (113). Benington F, and Morin RD (1951) An Improved Synthesis of Mescaline. *J. Am. Chem. Soc* 73, 1353–1353.
- (114). Tsao MU (1951) A New Synthesis of Mescaline. *J. Am. Chem. Soc* 73, 5495–5496.
- (115). McKenna D, and Riba J (2016) New World Tryptamine Hallucinogens and the Neuroscience of Ayahuasca. *Curr. Top. Behav. Neurosci*, 1–29.
- (116). Facchinelli F, and Weber A (2011) The Metabolite Transporters of the Plastid Envelope: An Update. *Front. Plant Sci* 2, 50. [PubMed: 22645538]
- (117). Abramovitch RA, and Shapiro D (1956) 880. Tryptamines, carbolines, and related compounds. Part II. A convenient synthesis of tryptamines and β -carbolines. *J. Chem. Soc* 0, 4589–4592.
- (118). Brandt SD, Moore SA, Freeman S, and Kanu AB (2010) Characterization of the synthesis of N,N-dimethyltryptamine by reductive amination using gas chromatography ion trap mass spectrometry. *Drug Test. Anal* 2, 330–338. [PubMed: 20648523]

Figure 1.

Structures of various natural and semisynthetic opioid alkaloids. ^aHydrocodone is also referred to as dihydrocodeinone. ^bVicodin is a combination of hydrocodone and acetaminophen. ^cPercocet is a combination of oxycodone and acetaminophen. ^dSuboxone is a combination of naloxone and buprenorphine. Structural requirements of the so-called “morphine rule” are highlighted in pink.

meperidine (**83**)
Demerol

fentanyl (**84**)
Perc-A-Pop

(±)-methadone (**85**)
Dolophine

(±)-tramadol (**86**)
Ultram

Figure 2.
Structures of various opioids.

chlordiazepoxide (**100**)
Librium

diazepam (**101**)
Valium

(±)-lorazepam (**102**)
Ativan

alprazolam (**103**)
Xanax

flunitrazepam (**104**)
Rohypnol, Roofie

Figure 3.
Structures of various benzodiazepines.

Figure 4.

Structures of various stimulants. ^aBronkaid is combination of (-)-ephedrine and guaifenesin.

^bThe active ingredient in Ritalin and Concerta is (±)-methylphenidate.

amphetamine aspartate monohydrate (**147**) dextroamphetamine saccharate (**148**) amphetamine sulfate (**149**) dextroamphetamine sulfate (**150**)

Figure 5.
Structures of the four amphetamine salts that constitute Adderall.

erythro

(-)-ephedrine (125)

threo

(+)-pseudoephedrine (126)

Figure 6. Comparison of ephedrine and pseudoephedrine stereo-chemistry. The medically used enantiomer of each compound is depicted.

dopamine (**162**)

serotonin (**163**)

Figure 7.
Structures of dopamine and serotonin, two neuro-transmitters.

Figure 8.
Structures of various hallucinogens. ^aResemblance to serotonin is highlighted in red.

(+)-lysergic acid (**169**)

(+)-LSD (**164**)

ergometrine (**170**)

(-)-ergotamine (**171**)

Figure 9.
Structures of various ergot alkaloids.

Scheme 1.

Three Total Syntheses of (–)-Morphine by the Gates, Gaunt, and Fukuyama Labs

a. Trost's 2002 Synthesis of (-)-Codeine

b. Stork's 2009 Synthesis of (±)-Codiene.

Scheme 2.
Synthesis of Codeine by the Trost and Stork Labs

Scheme 3.
Hudlicky's Second Generation Chemoenzymatic Synthesis of *ent*-Hydromorphone

a. Rice's 1980 Synthesis of (±)-Hydrocodone.

b. Mulzer and Trauner's 1996 Synthesis of (-)-Hydrocodone.

Scheme 4.

Total Synthesis of Hydrocodone from the Rice and Mulzer Labs

Scheme 5.
Conversion of (-)-Thebaine to (-)-Oxycodone

Scheme 6.
Fukuyama's 2014 Total Synthesis of (-)-Oxycodone

Scheme 8.
Hudlicky's 2013 Synthesis of (-)-Naloxone

Step 1: Isolation of morphine base from raw opium

Step 2: Synthesis of heroin•HCl from morphine base.

Scheme 9.
Illicit Synthesis of Heroin from Raw Opium

Scheme 10.
Illicit Synthesis of Krokodil

Scheme 11.
Eisleb's 1939 Synthesis of Demerol

Scheme 12.
Hite's Synthesis of Demerol

Scheme 13.
Janssen Company's 1965 Synthesis of Fentanyl

Scheme 14.
Synthesis of ($\pm$)-Methadone

Scheme 15.
Sternbach's Synthesis of Valium

Scheme 16.
Synthesis of Valium

Scheme 17.
Gates's Synthesis of Valium

a. The Upjohn Company's Synthetic Route to Xanax

b. Proposed Mechanism for the Transformation of ($\pm$)-**115** to **114**

Scheme 18.
Upjohn Company's 1971 Synthesis of Xanax

Scheme 19.
Robinson's 1917 Synthesis of Tropinone

Scheme 20.
Willstätter's Synthesis of (-)-Cocaine from 1903

Scheme 21.
Rapoport's 1998 Synthesis of (-)-Cocaine

Scheme 22.
Conversion of Powder Cocaine to Crack Cocaine

Scheme 23.
Belovsky's Synthesis of (±)-Amphetamine

Scheme 24.
Yamaguchi's Synthesis of (+)-Amphetamine

Scheme 25.

Illicit Synthesis of Methamphetamine via the “Shake and Bake” Method

Scheme 26.
Merck's 1912 Synthesis of (±)-MDMA

Scheme 29.
Woodward's Approach to (±)-Lysergic Acid

Scheme 30.
Fukuyama's Synthesis of (±)-Lysergic Acid

Scheme 31.
Fujii and Ohno's Synthesis of (±)-Lysergic Acid

Scheme 32.
Parke-Davis and Company's 1956 Synthesis of (±)-Ketamine

Scheme 33.
Zhang's Synthesis of (±)-Ketamine

Scheme 34.

Kiyooka's Second Generation Synthesis of (-)-Ketamine

Scheme 35.
Parke-Davis and Company's 1965 Synthesis of PCP

Scheme 36.
Pons's Synthesis of PCP

Scheme 37.
Illicit Synthesis of PCP by the "Bucket" Method

Scheme 38.
Spath's 1919 Synthesis of Mescaline

Scheme 39.
Tsao's Synthesis of Mescaline

Scheme 41.
Conversion of Tryptamine to DMT

Table 1.Psychoactive Drug Addiction in America^a

substance	estimated number of Americans with substance use disorder (SUD)
tobacco	52 million
alcohol	17 million
marijuana	8.3 Million
pain killers ^b	3.8 Million
Cocaine	1.9 Million
hallucinogen ^c	1.2 Million
methamphetamine	900 K
Heroin	330 K

^aValues taken from ref 6.

^bOpiates, opioids, and benzodiazepines.

^cLSD, PCP, MDMA, mescaline, DMT, and ketamine.

Table 2.**Opioid Conversion Chart^a**

drug	parenteral (mg)	oral (mg)
morphine (1)	10	30
codeine (2)	100–130	200
hydromorphone (5)	1.5–2	7.5–8
oxymorphone (6)	1	10
hydrocodone (7)	NA ^b	30–45
oxycodone (8)	NA ^b	20–30

^aValues from ref 21^bNA = not available.