CORRESPONDENCE

The Treatment of Primary Arthritis of the Finger and Thumb Joint

by PD Dr. med. Christian Karl Spies, Prof. Dr. med. Martin Langer, Prof. Dr. med. Peter Hahn, Prof. Dr. med. Lars Peter Müller, and Prof. Dr. med. Frank Unglaub in issue 16/2018

Radiotherapy

The authors of the selective literature search (1) did not mention the option of radiotherapy. The effectiveness of radiotherapy in inflammatory degenerative disorders has been known for more than 100 years and is used often and successfully in Germany. In the meantime, S2 guidelines for radiotherapy in benign disorders have been published (2–3).

About 75% of patients with osteoarthritis of the small joints benefit from radiotherapy; the usual treatment is six fractions with single doses of 0.5–1 Gy up to a total dose of 3 Gy or 6 Gy. Even after five years, most patients benefit from sustained pain relief. The tumor risk is negligible in the mostly older patients. For the dose range that is used in radiotherapy of joint arthritis, the literature does not mention any tumor induction (3).

The results of 809 patients after radiotherapy for painful arthritis of the hand and finger joints have been published. A response to the treatment was documented in 63–75% of patients (1). In a recently published study, the effectiveness of radiotherapy in 101 joints of 84 patients with thumb carpometacarpal osteoarthritis (rhizarthrosis) was evaluated—which is a larger number of patients compared to the studies mentioned in the review article. After 12 months, 70% of cases responded to the treatment, and complete pain response was achieved in 26% (4). Unfortunately, no randomized controlled trials have been reported for the radiotherapy of painful arthritis versus the usual therapeutic alternatives or non-treatment.

DOI: 10.3238/arztebl.2018.0596a

References

1. Spies CK, Langer M, Hahn P, Müller LP, Unglaub F: The treatment of primary arthritis of the finger and thumb joint. *Dtsch Arztebl Int* 2018; 115: 269–75.
2. Ott OJ, Niewald M, Weitmann HD, et al.: German Cooperative Group on Radiotherapy for Benign Diseases (GCG-BD). DEGRO guidelines for the radiotherapy of non-malignant disorders. Part II: painful degenerative skeletal disorders. *Strahlenther Onkol* 2015; 191: 1–6.
3. Deutsche Gesellschaft für Radioonkologie: Strahlentherapie gutartiger Erkrankungen. Fachgruppenspezifische evidenzbasierte S2e-Leitlinie der Deutschen Gesellschaft für Radioonkologie (DEGRO). Version 1.0 vom 14.11.2013.
4. Kaltenborn A, Bulling E, Nitsche M, Carl U, Hermann RM: The field size matters: low dose external radiotherapy for thumb carpometacarpal osteoarthritis. *Strahlenther Onkol* 2016; 192: 582–8.

Prof. Dr. med. Michael Pinkawa

MediClin Robert Janker Klinik
Klinik für Strahlentherapie und Radioonkologie
MVZ MediClin Bonn GmbH, Medizinisches Versorgungszentrum für Strahlentherapie und Neurochirurgie
michael.pinkawa@mediclin.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Autologous Fat Transplantation

Spies et al. provided an appealing review article on an important topic (1). As they mentioned, resection arthroplasty in advanced

osteoarthritis of the basal joint of the thumb (rhizarthrosis) is the standard procedure. Many patients, however, request minimally invasive alternatives for the treatment of osteoarthritis of the basal joint of the thumb. In this context, intraarticular autologous fat transplantation for the treatment of osteoarthritis of the basal joint of the thumb as interposition arthroplasty would have deserved to be mentioned. After it was initially presented in 2014, a larger study showed in 50 patients that in Eaton and Littler stage 2 arthritis, the long-term results after 12 months are convincing and comparable to those of trapeziectomy—and that morbidity is minimal. Even though no randomized, controlled studies have been published to date, initial comparative studies versus cortisone injections (3) or trapeziectomy with tendon interposition (4) support the value to this alternative.

DOI: 10.3238/arztebl.2018.0596b

References

1. Spies CK, Langer M, Hahn P, Müller LP, Unglaub F: The treatment of primary arthritis of the finger and thumb joint. *Dtsch Arztebl Int* 2018; 115: 269–75.
2. Herold C, Rennekampff HO, Groddeck R, Allert S: Autologous fat transfer for thumb carpometacarpal joint osteoarthritis: a prospective study. *Plast Reconstr Surg* 2017; 140: 327–35.
3. Haas EM, Volkmer E, Giunta RE: Pilot study on the effects and benefits of autologous fat grafting in osteoarthritis of the CMC-1 joint compared to intraarticular cortisone injection: results after 3 months. *Handchir Mikrochir Plast Chir* 2017; 49: 288–96.
4. Erne HC, Cerny MK, Ehrli D, et al.: Autologous fat injection versus lundborg resection arthroplasty for the treatment of trapeziometacarpal joint osteoarthritis. *Plast Reconstr Surg* 2018; 141: 119–24.

Prof. Dr. med. Christian Herold

HPC Oldenburg und DIAKO Bremen
christianherold@gmx.de

Prof. Dr. med. Hans-Oliver Rennekampff

Plastische & Ästhetische Chirurgie, Verbrennungschirurgie
Klinik für Orthopädie, Unfall-, Hand-, und Wiederherstellungschirurgie
Klinikum Leverkusen

Dr. med. Sixtus Allert

Klinik für Plastische und Ästhetische Chirurgie – Handchirurgie
Sana Klinikum Hameln-Pyrmont, Hameln

Conflict of interest statement

Professor Herold and Professor Rennekampff received study support (third party funding) from Asclepius Medizintechnik. Professor Rennekampff is the guideline coordinator of the S2k guideline on autologous fat transplantation.

In 2012, the company human med (Schwerin) made a body jet for a study available to the Sana Klinikum Hameln Pyrmont hospital for three years, free of charge, which Professor Herold and Dr. Allert used for research.

In Reply:

First of all, we wish to express our warmest thanks to our correspondents Herold et al. and Pinkawa. Autologous fat transplantation can certainly constitute an interesting alternative, although its efficacy will need to be proved by means of medium-term to long-term study results. The cited references present results that are, on the one hand, based on a follow-up period of only three months (in the correspondence: Haas et al.), and, on

the other hand, an average of 12 months' follow-up, with individual observations at 24 months (in the correspondence: Herold et al. and Erne et al.). Resection arthroplasty with or without suspension for the treatment of CMC-1-Arthritis Eaton/Littler stages 2–4 has in the meantime become a sufficiently studied therapeutic approach in terms of evidence-based aspects, with good results in the long term (1, 2). Revision surgeries after trapeziectomy because of recurrent or persistent pain are therefore rare. Alternative methods will have to measure up to these well-proven surgical techniques.

Radiotherapy to treat arthritis of the small joints of the hand certainly constitutes a tempting non-surgical approach. But for this approach too, randomized, controlled trials with long-term follow-up periods are lacking. The precise application of ionizing radiation to the target organ cannot always be reliably delivered to the hand, with its many joints in close proximity. For the treatment of rhizarthrosis, Kaltenborn et al. defined the clinically identifiable painful region with an additionally arbitrarily selected safety margin of 1 cm, respectively, in each direction (3). If this radiation field is put in relation to the actually measured CMC-1 joint areas of an average of 16.03 mm at the mean sagittal diameter for the metacarpal joint surface area and of 11.96 mm for the surface area of the trapezium joint, a not insignificant and unnecessary risk of damage to the adjacent anatomical structures exists (4). Viewing the literature, radiotherapy for the treatment

of rhizarthrosis has been studied overwhelmingly. Scarce data exist regarding radiotherapy of the digital joints. From a surgical perspective, the substantial risk of tissue scarring owing to radiotherapy needs to be borne in mind, which might render a subsequent surgical procedure difficult.

DOI: 10.3238/arztebl.2018.0596c

References

1. Gangopadhyay S, McKenna H, Burke FD, Davis TRCC: Five- to 18-year follow-up for treatment of trapeziometacarpal osteoarthritis: a prospective comparison of excision, tendon interposition, and ligament reconstruction and tendon interposition. *J Hand Surg Am* 2012; 37: 411–7.
2. Spekrijse KR, Vermeulen GM, Kedilioglu MA, et al.: The effect of a bone tunnel during ligament reconstruction for trapeziometacarpal osteoarthritis: a 5-year follow-up. *J Hand Surg Am* 2015; 40: 2214–22.
3. Kaltenborn A, Bulling E, Nitsche M, Carl UM, Hermann RM: The field size matters: low dose external beam radiotherapy for thumb carpometacarpal osteoarthritis. *Strahlenther Onkol* 2016; 192: 582–8.
4. Leversedge FJ: Anatomy and Pathomechanics of the thumb. *Hand Clin* 2008; 24: 219–29.
5. Spies CK, Langer M, Hahn P, Müller LP, Unglaub F: The treatment of primary arthritis of the finger and thumb joint. *Dtsch Arztebl Int* 2018; 115: 269–75.

On behalf of the authors

PD Dr. med. Christian Karl Spies

Abteilung für Handchirurgie,
Vulpius Klinik, Bad Rappenau
christianspies27@gmail.com

Conflict of interest statement

The author declares that no conflict of interest exists.