Published in final edited form as:

J Am Chem Soc. 1996 March 13; 118(10): 2527–2528. doi:10.1021/ja9539148.

Synthesis of Enantiomerically Pure Anti-Aldols: A Highly Stereoselective Ester-Derived Titanium Enolate Aldol Reaction

Arun K. Ghosh and Masanobu Onishi

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607

The aldol reaction leading to enantioselective construction of carbon-carbon bonds has emerged as an extremely powerful method in organic synthesis. In this context, a number of excellent synthetic methods have been developed over the years.¹ The control of both relative and absolute acyclic stereochemistry in aldol reactions can now be achieved in a highly stereoselective manner with a high degree of predictability. Such control has become particularly sophisticated in the synthesis of various syn-aldol products and is used widely.² We have recently reported that commercially available optically active *cis*-1-amino-2-indanol-derived oxazolidinones are highly effective chiral auxiliaries for syn-aldol reactions.³ The corresponding enantioselective anti-aldol methodologies are currently an active area of research.⁴ Herein, we report the development of a convenient *cis*-1-arylsulfonamido-2-indanol-derived titanium ester enolate based aldol reaction to provide anti-aldol products with excellent diastereoselectivity and isolated yields. Optically active *cis*-1-arylsulfonamido-2-indanols are readily accessible by sulfonylation of commercially available, enantiomerically pure *cis*-1-amino-2-indanols. The optically pure anti-*R*-methyl- β -hydroxy acids are conveniently obtained after removal of the chiral auxiliary under mild saponification conditions, and the chiral auxiliary is recovered without loss of optical purity.

The 1*R*,2*S*-chiral sulfonamide **1** was prepared (Scheme 1) by reaction with commercially available⁵ optically active 1(*R*),2-(*S*)-*cis*-aminoindan-2-ol, *p*-toluenesulfonyl chloride (1 equiv), and triethylamine (3 equiv) in CH₂Cl₂ in the presence of a catalytic amount of DMAP at 23 °C for 12 h (85–92% yield). The acylation of hydroxy sulfonamide **1** with propionyl chloride (1.2 equiv) and triethylamine (3 equiv) in CH₂Cl₂ at 23 °C for 12 h afforded the propionate ester **2** (mp 106 °C; $\alpha_{\text{D}}^{23} +86^\circ$, *c* 0.98, CHCl₃) after silica gel chromatography (91% yield). The titanium enolate of **2** was generated by reaction with 1.2 equiv of TiCl₄ in CH₂Cl₂ at 0–23 °C for 15 min followed by addition of 4 equiv of *N*-ethyl-diisopropylamine at 23 °C and stirring of the resulting brown solution for 1 h.^{6,7} The ¹H-NMR (400 MHz) studies of the titanium enolate generated in a mixture of CDCl₃ and CH₂Cl₂, as described above, established that the enolization is complete under these conditions, providing a single enolate presumably with *Z*-geometry.⁸ The titanium enolate thus generated was reacted with 2 equiv of butyraldehyde or isobutyraldehyde at –78 to

Supporting Information Available: Experimental procedures and spectral data for aldols and their derived β -hydroxy acids (11 pages). This material is contained in many libraries on microfiche, immediately follows this article in the microfilm version of the journal, can be ordered from the ACS, and can be downloaded from the Internet; see any current masthead page for ordering information and Internet access instructions.

23 °C for several hours; interestingly, however, no aldol product was obtained, and the starting propionate ester **2** was recovered unchanged. However, the reaction of the above titanium enolate with various aldehydes precomplexed with TiCl₄ proceeded with good to excellent diastereoselectivities and isolated yields. The reactions are typically carried out by addition of the above titanium enolate to a solution of 2 equiv of aldehyde precomplexed with 2.4 equiv of TiCl₄ in CH₂Cl₂ at –78 °C followed by stirring the resulting mixture for 1 h and workup with aqueous NH₄Cl solution. The results are summarized in Table 1. The ester enolate aldol reaction with a variety of conjugated and nonconjugated aldehydes proceeded with excellent anti-diastereoselectivity except in the case of benzaldehyde, which did not exhibit any selectivity.⁹ The diastereomeric mixture ratio was determined by ¹H-NMR (400 MHz) as well as by HPLC analysis of the aldol products prior to chromatography. Aldol reaction with butyraldehyde (entry 3) afforded an anti-diastereoselectivity of 95:5. Both anti- and syn-isomers could be separated by silica gel chromatography (74% combined yield). The reaction with isovaleraldehyde proceeded with almost complete anti-diastereoselectivity (>99% de by HPLC; 400 MHz ¹H-NMR revealed only one diastereomer) with 97% isolated yield after silica gel chromatography. Similarly, the reaction with conjugated aldehydes also provided excellent anti-diastereoselectivity.¹⁰

In order to establish the relative stereochemistry, the antiisomer **3** derived from crotonaldehyde (entry 6) was converted to the isopropylidene derivative **5** by reduction with LAH in THF at 0 °C followed by exposure of the resulting diol to dimethoxypropane in CH₂Cl₂ in the presence of a catalytic amount of pyridinium *p*-toluenesulfonate. The ¹H-NMR (400 MHz) analysis of **5** firmly established the identity of the antiisomer in this chiral ester enolate based aldol reaction. Of particular interest, a coupling constant (*J*_{CD}) of 10.1 Hz was measured between the H_C and H_D protons of **5**.¹¹ The absolute configurations of the new asymmetric centers of **3** were assigned after removal of the chiral sulfonamides, conversion of the resulting anti- α -methyl- β -hydroxy acids to the corresponding methyl ester, and comparison of the optical rotations of the resulting **7** with literature values.^{4a} For example, treatment of the isobutyraldehyde-derived anti-aldol product (entry 4) with aqueous lithium hydroxide in THF at 23 °C for 2 h afforded the corresponding acid which was converted to optically pure methyl ester **7** (α _D²³ –14.2°, *c* 1.04, CHCl₃; lit.^{4a} α _D²³ –12.5°, *c* 1.04, CHCl₃) with an excess of diazomethane in diethyl ether. The chiral sulfonamide **1** was recovered (95% after silica gel chromatography) after saponification without loss of optical purity (α _D²³ –34°, *c* 1.5, CHCl₃). Similarly, the butyraldehyde-derived anti-aldol product was converted to the optically pure methyl ester **7** (α _D²³ –4°, *c* 2.0, CHCl₃; lit.^{4a} α _D²³ –2.5°, *c* 1.03, CHCl₃). The relative and absolute stereochemistries of the isobutyraldehyde-derived syn-aldol product **4** (entry 4) were established by comparison of ¹H-NMR (400 MHz) and optical rotation of **4** with an authentic sample prepared utilizing Evan's boron enolate based syn-aldol reaction.^{2a} The assignment of stereochemistry for other syn-aldol products was made on the basis of comparisons of ¹H-NMR spectra.

The high degree of stereoselection associated with this current asymmetric anti-aldol process could be rationalized by a Zimmerman-Traxler type transition state model **A** (Figure 1).¹² The model is derived based on the following assumptions: (1) the geometry of the titanium enolate is *Z*; (2) the titanium enolate is a seven-membered metallocycle with a chairlike

conformation; (3) a second titanium metal is involved in the transition state where it is chelated to indanyloxy oxygen as well as to the aldehyde carbonyl in a six-membered chairlike transition state. The involvement of two titanium metal atoms is supported by the fact that the titanium enolate derived from **2** does not react with aldehydes without precomplexation with TiCl_4 . The above transition state model accounts for the observed stereoselection except in the case of benzaldehyde, which has shown no selectivity. While this model explains much of the present stereoselection, the evidence of such a model requires further experimentation which is currently in progress.

In summary, the present chiral ester derived titanium enolate aldol reaction with various aldehydes represents a highly effective synthetic protocol for providing anti-aldol products with high levels of diastereo- and enantioselectivity. The generality of the current anti-selective aldol process has been demonstrated with nine different aldehydes. Since both enantiomers of *cis*-1-arylsulfonamido-2-indanol are readily prepared from the commercially available optically active *cis*-1-amino-2-indanols, the present anti-aldol methodology provides a convenient access to either anti-aldol enantiomer with high optical purity. Mechanistic investigations as well as synthetic applications of the current anti-aldol methodology are the subject of ongoing research in our laboratory.

Acknowledgment.

Financial support for this work was provided by the University of Illinois at Chicago. The authors thank Professor Steven Zimmerman and Donald Wink for helpful discussions. M.O. is a visiting scientist from Nihon Nohyaku Co. Ltd., Japan.

References

- (1). (a) Heathcock CH. In *Asymmetric Synthesis*; Morrison JD, Ed.; Academic Press: New York, 1984; Vol. 3, p 111. (b) Evans DA; Nelson JV; Taber TR *Top. Stereochem* 1982, 13, 1. (c) Masamune S; Choy W; Petersen JS; Sita LR *Angew. Chem., Int. Ed. Engl* 1985, 24, 1. (d) Braun M *Angew. Chem., Int. Ed. Engl* 1987, 26, 24. (e) Franklin AS; Paterson I *Contemp. Org. Synth* 1994, 317.
- (2). (a) Evans DA; Bartoli J; Shih TL. *J. Am. Chem. Soc* 1981, 103, 2127. (b) Evans DA *Aldrichimica Acta* 1982, 15, 23. (c) Paterson I; Lister MA; McClure CK *Tetrahedron Lett* 1986, 27, 4787. (d) Corey EJ; Imwinkelried R; Pikul S; Xiang YB *J. Am. Chem. Soc* 1989, 111, 5493. (e) Oppolzer W; Blagg J; Rodriguez I; Walther EJ *Am. Chem. Soc* 1990, 112, 2767. (f) Roder H; Helmchen G; Peters E-M; Peters K; von Schnering H-G *Angew. Chem., Int. Ed. Engl* 1984, 23, 898. (g) Sankhavasi W; Yamamoto M; Kohmoto S; Yamada K *Bull. Chem. Soc. Jpn.* 1991, 64, 1425. (h) Drewes SE; Malissar DGS; Roos GH P. *Chem. Ber.* 1991, 124, 2913.
- (3). Ghosh AK; Duong TT; McKee SPJ *Chem. Soc., Chem. Commun* 1992, 1673.
- (4). (a) Meyers AI; Yamamoto Y *Tetrahedron* 1984, 40, 2309. (b) Helmchen G; Leikauf U; Tauffer-Knopfel I *Angew. Chem., Int. Ed. Engl* 1985, 24, 874. (c) Gennari C; Bernardi A; Colombo L; Scolastico CJ *Am. Chem. Soc* 1985, 107, 5812. (d) Palazzi C; Colombo L; Gennari C *Tetrahedron Lett.* 1986, 27, 1735. (e) Oppolzer W; Marco-Contelles J; *Helv. Chim. Acta* 1986, 69, 1699. (f) Masamune S; Sato T; Kim BM; Wollman TA *J. Am. Chem. Soc.* 1986, 108, 8279. (g) Danda H; Hansen MM; Heathcock CH *J. Org. Chem.* 1990, 55, 173. (h) Corey EJ; Kim SS *J. Am. Chem. Soc* 1990, 112, 4976. (i) Corey EJ; Kim SS *Tetrahedron Lett* 1990, 31, 3715. (j) Meyers AG; Widdowson KL *J. Am. Chem. Soc* 1990, 112, 9672. (k) Duthaler RO; Herold P; Helfer S-W; Riediker M *Helv. Chim. Acta* 1990, 73, 659. (l) Walker MA; Heathcock CH *J. Org. Chem.* 1991, 56, 5747. (m) Oppolzer W; Lienard P *Tetrahedron Lett.* 1993, 34, 4321.
- (m) Gennari C; Moresca D; Vieth S; Vulpetti A *Angew. Chem., Int. Ed. Engl* 1993, 32, 1618.
- (n) Paterson I; Wren SPJ *Chem. Soc., Chem. Commun* 1993, 1790 and references cited therein.

- (5). Available from Aldrich Chemical Co., Milwaukee, WI and Sepracor Inc., Marlborough, MA 01752.
- (6). While alkyl esters are known^{7a} to be not enolizable with $\text{TiCl}_4/\text{Et}_3\text{N}$, formation of titanium enolate from esters through internal chelation with the sulfonamido group has been reported; see: Xiang Y; Olivier E; Ouimet N; Tetrahedron Lett. 1992, 33, 457.
- (7). Evans DA; Urpi F; Somers TC; Clark JS; Bilodeau MT J. Am. Chem. Soc 1990, 112, 8215. Evans DA; Rieger DL; Bilodeau MT; Urpi FJ Am. Chem. Soc 1991, 113, 1047. Bonner MP; Thornton ER J. Am. Chem. Soc. 1991, 113, 1299. Siegel C; Thornton ER J. Am. Chem. Soc 1989, 111, 5722. Siegel C; Thornton ER Tetrahedron Lett 1986, 27, 457.
- (8). ^1H -NMR (CDCl_3 , 400 MHz): δ 7.78 (d, 2 H, $J = 8.3$ Hz), 7.16–7.35 (m, 6 H), 5.55 (d, 1 H, $J = 5.7$ Hz), 4.67 (q, 1 H, $J = 7$ Hz), 4.28 (m, 1 H), 3.05–3.14 (m, 2 H), 2.50 (s, 3 H), 1.51 (d, 3 H, $J = 7.0$ Hz).
- (9). When the aldol reaction was carried out with a 1:1 mixture of benzaldehyde and isovaleraldehyde, the ^1H -NMR of crude products revealed the presence of a 65:35 ratio of anti/syn-diastereomers of benzaldehyde and a single anti-isomer of isovaleraldehyde as seen in entry 5.
- (10). All new compounds gave satisfactory spectroscopic and analytical results.
- (11). ^1H -NMR (CDCl_3 , 400 MHz): δ 5.73 (dq, 1 H, $J = 6.5, 15.3$ Hz, HF), 5.38 (ddd, 1 H, $J = 1.3, 7.7, 15.3$ Hz, HE), 3.84 (dd, 1 H, $J = 8.1, 10.1$ Hz, HD), 3.73 (dd, 1 H, $J = 5.1, 11.7$ Hz, HB), 3.54 (dd, 1 H, $J = 11.5, 11.5$ Hz, HA), 1.72 (dd, 3 H, $J = 1.3, 6.5$ Hz), 1.66 (m, 1 H, He), 1.47 (s, 3 H), 1.41 (s, 3 H), 0.7 (d, 3 H, $J = 6.8$ Hz).
- (12). Zimmerman HE; Traxler MD J. Am. Chem. Soc 1957, 79, 1920.

Figure1.

Scheme 1.

^a Key: (a) CH_3CH_2COCl , Et_3N , CH_2Cl_2 , 23 °C; (b) $TiCl_4$, iPr_2NEt , 23 °C then $RCHO$ and $TiCl_4$, CH_2Cl_2 , -78 °C; (c) $LiAlH_4$, THF, 0–23 °C; (d) $Me_2C(OMe)_2$, PPTS, CH_2Cl_2 , 23 °C; (e) $LiOH$, THF- H_2O , 23 °C; (f) CH_2N_2 , Et_2O , 23 °C.

Table 1.Aldol Reaction of Propionate Ester **2** with Representative Aldehydes

entry	aldehyde	% yield ^a	anti:syn (3/4) ^b
1	MeCHO	50 ^c	85:15
2	EtCHO	50	85:15
3	nPrCHO	74	95:5
4	iPrCHO	91	85:15
5	iBuCHO	97	>99:1
6	MeCHdCHCHO	41	95:5
7	PhCHO	85	45:55
8	PhCH ₂ CH ₂ CHO	44	96:4
9	PhCHdCHCHO	63	99:1

^aIsolated yield of both isomers after silica gel chromatography. Time = 1 h.^bDiastereomeric ratios were determined by ¹H-NMR and HPLC.^cIsolated anti-isomer only.