Received:
24 May 2017

Revised:
13 January 2018

Accepted:
30 January 2018

<https://doi.org/10.1259/bjr.20170389>

Cite this article as:

Koutoulidis V, Papanikolaou N, Mouloupoulos LA. Functional and molecular MRI of the bone marrow in multiple myeloma. *Br J Radiol* 2018; **91**: 20170389.

REVIEW ARTICLE

Functional and molecular MRI of the bone marrow in multiple myeloma

¹VASSILIS KOUTOULIDIS, MD, ²NICKOLAS PAPANIKOLAOU, PhD and ¹LIA A MOULOPOULOS, MD

¹First Department of Radiology, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece

²Computational Clinical Imaging Group, Centre for the Unknown, Champalimaud Foundation, Lisbon, Portugal

Address correspondence to: Dr Vassilis Koutoulidis

E-mail: vkoutoulidis@med.uoa.gr

ABSTRACT

MRI plays an important role in the management of patients with plasma cell neoplasms and has been recognized as a biomarker of malignancy in the novel criteria for the diagnosis of multiple myeloma. Functional and molecular MRI techniques such as diffusion-weighted imaging (spinal or whole body), intravoxel incoherent motion, and dynamic contrast enhanced MRI, provide additional information related to tumor cellularity and angiogenesis, which may have prognostic implications for patients with smoldering and symptomatic myeloma. These non-invasive functional techniques are also being evaluated as imaging biomarkers for response assessment in myeloma patients. The purpose of this article is to provide a comprehensive critical review on the current use and potential future applications of these advanced MRI techniques in multiple myeloma. In addition, we will address the technologies involved and describe the qualitative and quantitative characteristics of normal bone marrow with these techniques.

INTRODUCTION

Functional and molecular MRI techniques play an increasing role in the management of patients with multiple myeloma. Diffusion-weighted imaging (DWI) and dynamic contrast enhanced (DCE) MRI permit non-invasive evaluation of tumor-specific characteristics related to cellularity and angiogenesis and can thus improve diagnosis, prognostication, treatment planning and response assessment.^{1–3} Both diffusion and perfusion quantification can be provided from intravoxel incoherent motion (IVIM) modelling.⁴ Whole body MRI (WBMRI) has been recently recommended as a first line imaging modality by the International Myeloma Working Group.⁵ Whole body DWI (WBDWI) in particular, can provide a comprehensive assessment of disease burden before, during and after therapy.⁶ In this article, we will critically review the current use and potential future applications of these advanced MRI techniques in multiple myeloma. In addition, we will briefly address the technologies involved and describe the qualitative and quantitative characteristics of normal bone marrow with these techniques.

DIFFUSION-WEIGHTED IMAGING

DWI examines the self-diffusion of water molecules in the extracellular space, the intracellular space, and the intravascular space.⁷ In biological tissues, water diffusion is

not free, but impeded by cell membranes, macromolecules and intracellular organelles. Diffusion is more restricted in tissues with high cellular density, decreased extracellular space and intact cell membranes.^{8,9} An MRI pulse sequence sensitized to microscopic water mobility using strong gradient pulses can thus be utilized to provide insight into the complexity of the environment, which in turn can reveal information related to tissue microarchitecture. Apparent diffusion coefficient (ADC) is the most widely used diffusion-related quantitative biomarker, and can be estimated from two or more different *b*-values, an index used to control the sensitivity of DWI contrast to water mobility, measured in s mm^{-2} ^{10,11} (Figure 1). A more complex model, the IVIM model, takes into account flow phenomena that contribute to DWI contrast.¹² This model attempts to measure the diffusion signal contamination with the added term of “pseudo-diffusion” (D^*), representing signal loss resulting from other processes, most likely microperfusion of blood nutrients at the capillary level. Values for the true diffusion coefficient (*D*), pseudodiffusion coefficient (D^*), and microperfusion fraction (*f*) can, thus, be calculated. An alternative technique that attempts to quantify the microperfusion contribution to the diffusion signals is based on the acquisition of at least three *b*-values, a low *b*-value (*b*₀), a moderate *b*-value (between 150 and 200) and a high *b*-value (>800). According to this method, two separate

Figure 1. DWI of the normal bone marrow in a young adult. Diffusion-weighted images of the lumbosacral spine in a 21-year-old healthy female, with b -values (s mm^{-2}) 0 (a), 150 (b), 250 (c), 500 (d), and 750 (e). The mean ADC value of the five lumbar vertebral bodies calculated from the monoexponential ADC map generated by using all five b -values (f) was $0.421 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

ADC maps are generated by using the low and moderate b -values (ADC_{low}) and the moderate and high b -values (ADC_{high}). The former reflects diffusion and perfusion effects, while the latter reflects only diffusion effects. By calculating the relative percent change between ADC_{low} and ADC_{high} , the so-called “relative enhanced diffusivity”, one can quantify tissue vascularity based on the acquisition of a three b -value DWI sequence.¹³ (Figure 2).

Whole body MRI and whole body diffusion-weighted imaging

WBMRI is already incorporated in the clinical routine due to recent technological advances in the field of RF technology, which include integrated high-density phased array coils that can cover large anatomical areas, providing an adequate signal to noise ratio. Main areas of clinical applications of WBMRI comprise imaging metastatic disease, but also evaluation of patients with hematological malignancies like multiple myeloma and

lymphoma.^{6,14,15} WBDWI, in particular, can provide a comprehensive qualitative assessment of disease burden in multiple myeloma (MM), by visual inspection of bright areas on the high b -value images using inversion recovery based echo planar imaging diffusion sequences like DWIBS (diffusion-weighted whole body imaging with background body signal suppression). Moreover, quantitative methods of WBDWI, including calculation of total disease burden and marrow ADC analysis of the whole skeleton with segmentation techniques are used to monitor treatment response.^{6,16} Finally, Dixon-type sequences that generate four types of images (in-phase, opposed phase, fat only, water only) in relatively fast acquisition times are being increasingly incorporated in WBMRI protocols for MM. They may be valuable both for lesion detection as well as for response assessment due to their potential for fat quantification.^{17,18}

DWI of normal bone marrow

Understanding bone marrow composition and its dynamic changes is mandatory for correct interpretation of findings, both on conventional and advanced MRI sequences. Red (hematopoietic) marrow and yellow (fatty) marrow differ in cellular and chemical composition. Fetal bone marrow is entirely red, and following birth the process of red to yellow marrow conversion starts and continues through life, following an orderly pattern.¹⁹ Almost equal proportions of water and fat are found in the chemical composition of red marrow. Yellow marrow, which is made almost entirely of adipose cells with very few hematopoietic elements, contains a much larger proportion of fat with very little water.^{20,21} Moreover, red marrow has a rich vascular supply with a dense sinusoidal network, whereas fatty marrow vascularization is considerably decreased.²² Water diffusivity is very restricted in normal bone marrow, even more so in yellow marrow than in red marrow. Although there are probably multiple factors influencing the free diffusivity of water molecules in the bone marrow, the prominent presence of fat seems to play the most important role. The large adipose cells which are the major component of yellow marrow (as well as a significant component of red marrow) presumably impede the free movement of water molecules, because of increased extracellular space tortuosity and the hydrophobic nature of fat.^{23,24} Thus, ADCs of normal bone marrow are very low, and ADCs of yellow marrow are lower than those of red marrow. Accordingly, the steady increase of the fat component in marrow throughout life may help explain the decrease of normal bone marrow ADCs with aging, which has been noted in many studies²⁵ (Figure 3). The inverse correlation of marrow ADCs and marrow fat content has also been demonstrated in studies of osteoporosis.^{26,27} Lower ADCs of yellow marrow may also be related to reduced water proton density and decreased perfusion (the latter effect more pronounced if b_0 images are used for ADC calculations). Conversely, higher ADCs of red marrow may be explained by lower adipose content, higher water content and increased vascularization. In most studies, normal bone marrow ADCs fall below $0.6 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$ and can be as low as $0.15 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$, especially in elderly individuals with very prominent fatty marrow.^{3,24,28,29} On high b -value images, red marrow has higher signal intensity than yellow marrow, again probably related to its higher water content. Marrow infiltrating tumors, which, by definition, replace marrow fat cells (focally,

Figure 2. RED maps in three different patients with multiple myeloma presenting with three different MRI patterns. RED values of the diffuse MRI pattern are significantly higher relative to the other two patterns, indicating increased bone marrow vascularity. The focal area of involvement (arrow) in the focal MRI pattern, also shows increased RED value compared to the normal MRI pattern. RED, relative enhanced diffusivity.

diffusely or both) and display high cellularity, appear with high signal on high b -value images and with higher ADC values than normal marrow (Figure 4). In other words, bone marrow malignancies are characterized by increased diffusivity compared to the very low diffusivity of normal marrow.^{24,30} A narrow zone of overlap may exist between ADC values of marrow malignancies and ADC values of hypercellular red marrow, especially in very young subjects as well as in patients receiving hematopoietic growth factors, such as granulocyte-colony

stimulating factor. Padhani et al showed a non-linear, but rather polynomial relation between signal intensity on high b -value images and ADC values. They hypothesized that with increasing marrow cellularity due to tumor infiltration and fat cell replacement, signal intensity and ADCs increase, but that at some point, possibly when all fat cells are lost, further increases in marrow cellularity due to tumor, may result in ADC reductions.^{30,31} For a more detailed discussion on normal marrow ADCs and differentiation from ADCs of malignantly

Figure 3. DWI of the normal bone marrow in a healthy female with prominent fatty marrow. Diffusion-weighted images of the lumbosacral spine in a 69-year-old healthy female with b -values (s mm^{-2}) 0 (a), 150 (b), 250 (c), 500 (d), and 750 (e). The mean ADC value of the five lumbar vertebral bodies calculated from the monoexponential ADC map generated by using all five b -values was $0.249 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

Figure 4. DWI of tumor infiltrated marrow in a patient with multiple myeloma. Diffusion-weighted images with b -values (s mm^{-2}) 0 (a), 150 (b), 250 (c), 500 (d), and 750 (e) of the lumbosacral spine in a 37-year-old female with newly diagnosed myeloma (95% bone marrow plasmacytosis) and a diffuse pattern of involvement. The mean ADC value of the five lumbar vertebral bodies calculated from the monoexponential ADC map generated by using all five b -values was $0.692 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$. Also note diffusely elevated signal of the marrow on the high b -value images compared to Figures 1 and 3. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

infiltrated marrow, please see the section on DWI of multiple myeloma.

DYNAMIC CONTRAST-ENHANCED (DCE) MRI

DCE-MRI has been used to assess perfusion in various anatomical regions, including the bone marrow.³² It is usually performed with a combination of dynamic acquisition of T_1 weighted images before, during and after the intravenous injection of a paramagnetic contrast agent. The dynamics of the signal intensity in every pixel of the examined area contain tissue-specific information related to vascular pathophysiology. The simplest, non-quantitative, approach is to visually assess the shape of the time-signal intensity curves (TICs), and classify them into predefined tissue-specific types.³³ Perfusion can also be assessed using semi-quantitative and quantitative markers. Widely used semi-quantitative parameters derived from the TICs include maximum enhancement (E_{max}), time to peak, area under the curve, and washin and washout rates²⁰ (Figure 5). Finally, complex pharmacokinetic models (*e.g.* Tofts, extended Tofts, and Brix) are frequently used to provide objective, quantitative measurements of blood volume and vascular permeability based on DCE data analysis.³⁴

DCE-MRI of normal bone marrow

DCE-MRI enables the evaluation of bone marrow microcirculation in a non-invasive way. Its ability to simultaneously image a large volume of marrow overcomes the potential sample bias of conventional assessments of angiogenesis, such as trephine biopsy-based measurements of microvessel density (MVD). Moreover, it may provide functional information on marrow vascularity, such as blood flow and microvessel permeability.³¹ DCE analysis of normal bone marrow can be done qualitatively, by studying the shape of the TIC. Normal bone marrow has a

characteristic dynamic contrast profile with early enhancement, which peaks within 1 min from contrast injection.³⁵ In older subjects with prominent fatty marrow, the curve shows a relatively flat enhancement slope and very low maximal enhancement followed by a plateau or an almost imperceptible washout. In subjects with a higher proportion of red marrow, the curve displays a steeper enhancement slope, higher maximum enhancement and a washout effect^{36–38} (Figure 6). Studies evaluating normal marrow perfusion using semi-quantitative indices derived from TICs produced with the region of interest (ROI) method, have shown that maximum enhancement (E_{max})

Figure 5. DCE-MRI-derived time-signal intensity curve and the corresponding semi-quantitative markers in a model free analysis approach. DCE, dynamic contrast-enhanced, TTP, time to peak; AUC, area under the curve.

Figure 6. Fitted time-signal intensity curves of normal bone marrow from DCE-MRI exams of the lumbar spine in two different healthy individuals show a flat enhancement slope and very low maximal enhancement followed by a plateau (green curve) or mild washout (red curve). DCE, dynamic contrast-enhanced.

decreases significantly with age.^{36,39} A strong negative correlation between maximum enhancement and marrow fat content has also been found.^{26,36,37} Another study evaluated the effects of age and marrow fat on quantitative DCE-MRI parameters including the transfer constant (K_{trans}) and the efflux rate constant (K_{ep}). Breault et al found a statistically significant correlation between both age and fat fraction and the parameters K_{trans} and K_{ep}, but this correlation was much weaker than the one found between age and marrow fat and the semiquantitative parameters in the previous studies. It was postulated that a major contributor to this difference may have been the voxel-by-voxel analysis of the quantitative study (which excludes from analysis data pixels with insufficient enhancement, which are more likely to contain primarily yellow marrow), vs the whole ROI analysis of most semiquantitative studies (where the lack of enhancement in the fatty components produces a lower mean value for maximum enhancement).⁴⁰

MULTIPLE MYELOMA

MM is a clonal plasma cell neoplastic disorder characterized by a multistep progressive pathogenesis, with all cases developing from an asymptomatic monoclonal gammopathy, either monoclonal gammopathy of undetermined significance (MGUS, with an annual rate of progression to MM of 0.5–1%) or smoldering MM (SMM, with an annual rate of progression to MM of approximately 10% for the first 5 years after diagnosis).^{41,42} The diagnosis of both MGUS and SMM requires absence of hypercalcemia, renal failure, anemia and bone lesions (the so-called CRAB features), that can be attributed to the underlying plasma cell proliferative disease. In 2014, the International Myeloma Working Group (IMWG) revised the diagnostic criteria for MM to enable early diagnosis and prompt treatment for a subset of high-risk patients without evidence of end organ damage.⁴³ This decision was driven by evidence that smoldering myeloma is a biologically heterogeneous condition, with some SMM patients demonstrating a very low risk for progression, similar to that of MGUS, whereas, at the other end of the spectrum, some patients develop end organ damage within 2 years after

the initial diagnosis. The changes of the disease definition were made possible by major developments in treatment, as well as by considerable advances in imaging technology. Traditionally, the conventional radiographic skeletal survey was used in MM patients in order to determine the presence of bone lesions.⁴⁴ The updated IMWG criteria, acknowledging the emergence of more sensitive modern imaging techniques, states explicitly that the presence of one or more osteolytic lesions seen on CT (including low-dose whole body CT) or PET-CT does meet the criteria for bone disease in multiple myeloma, even when conventional radiographs are negative.⁴³ A very recent multicenter study of 212 patients lends support to these revised criteria, by showing not only that whole body CT is significantly more sensitive than the conventional radiographic skeletal survey for detecting myeloma-related osteolyses, but also that osteolytic lesions detected by whole body CT alone are prognostically significant for the progression of SMM into MM.⁴⁵

MRI has also been used extensively for the management of patients with plasma cell neoplasms with the following MRI patterns of involvement described in multiple studies: normal (*i.e.* normal appearing), focal, variegated, diffuse, combined focal and diffuse.^{44,46–49} The role of MRI in the initial assessment of MM has been recognized in the updated IMWG criteria, in which MRI is considered as one of three biomarkers of malignancy for the diagnosis of MM (along with clonal bone marrow plasma cells $\geq 60\%$ and serum free light chain ratio ≥ 100). Specifically, to identify SMM patients with a high probability of progression to MM within 2 years, the IMWG defines the presence of more than one focal lesion on MRI as establishing the diagnosis of MM.^{43,50,51} A diffuse MRI marrow infiltration pattern has also been associated with an increased risk of progression.^{50,52} The IMWG, nevertheless, did not recommend the inclusion of the diffuse MRI pattern in the updated definition of MM. One explanation may be that the number of patients with a diffuse MRI pattern, no lytic lesions or other CRAB features and absence of extreme bone marrow plasmacytosis (60%) is probably small and cannot be adequately analyzed, especially in single center studies. A further potential difficulty, is the absence of a consensual definition for the diffuse pattern of marrow infiltration on conventional MRI sequences.³ Therefore, regarding the diffuse MRI pattern, the IMWG underlines the need for additional data on the specific progression risk, in the absence of focal lesions and extreme bone marrow plasmacytosis.⁴³

The recommendations for the use of MRI in the aforementioned updated IMWG criteria for the diagnosis of multiple myeloma are based on conventional MRI studies. However, it is very important to note that functional and molecular MRI techniques—specifically DWI and DCE MRI—are mentioned and briefly described in a consensus statement on the role of MRI in the management of patients with MM that was published almost concurrently by the IMWG. The consensus panel states as a Grade A recommendation that MRI is the imaging gold-standard method for the detection of bone marrow involvement in MM, since it can detect early bone marrow infiltration, even in the absence of bone destruction. Of note, the whole body approach is proposed, although it is stressed that MRI of the spine and pelvis can detect

Figure 7. Whole body MRI in a 68-year-old male with multiple myeloma. Coronal inverted grayscale b 800 diffusion-weighted (a), T_1 weighted (b) and STIR (c) images show two lesions in the L1 vertebral body and in the diaphysis of the left femur (arrows). Note excellent conspicuity of lesions on the diffusion weighted image, including the small extraosseous soft tissue extension at L1 (arrow head). STIR, short tau inversion-recovery.

the majority of focal lesions in MM, and can be used in cases where WBMRI is not available.⁵

Diffusion-weighted MRI in multiple myeloma

A number of recent studies have underlined the added value of DWI in the management of patients with MM.^{3,6,24,53,54} DWI in MM can be applied as part of an axial skeleton (*i.e.* spinal alone or spinal and pelvic) MRI study or within a WB protocol. As stated earlier, normal bone marrow is characterized by very restricted water diffusivity, with low signal intensity on high b -value images. Qualitative assessment of high b -value images results in excellent lesion conspicuity, with focal MM lesions appearing as high signal foci against low signal normal marrow (by definition, lesions appear with low signal on the inverted grayscale images of a WBDWI study)⁵⁵ (Figure 7). Diffuse infiltration is also characterized by increased signal on high b -value images. Evaluation in these cases can be more difficult by visual assessment alone, due to the lack of a reliable qualitative internal reference standard and correlation with the ADC values is advised. It has been shown that myeloma-infiltrated marrow displays ADC values that differ significantly from those of bone marrow in normal subjects. Messiou et al studied 41 volunteers and 43 patients with metastatic (breast and prostate) and myelomatous involvement and performed axial DWI of the lumbar spine and pelvis with five b -values ranging from 0 to 750 s mm⁻². They found that an ADC threshold of 0.655×10^{-3} mm² s⁻¹ separated normal from abnormal marrow with a sensitivity and specificity of 90

and 93%, respectively. In this study, 11 patients with MM had a mean ADC of 1.235×10^{-3} mm² s⁻¹ with no separate analysis of values from focal MM lesions and diffuse myelomatous disease. The mean ADC of the age-matched healthy control group was 0.471×10^{-3} mm² s⁻¹.²⁴ In a different study by Padhani et al, the mean ADC of a mixed group of focal and diffuse myeloma lesions (calculated from b -values of 50 and 800 or 900 s mm⁻²) was 0.875×10^{-3} mm² s⁻¹.³⁰ Koutoulidis et al performed spinal MRI with DWI (b -values 0, 150, 250, 500, 750 s mm⁻²) in 99 newly-diagnosed MM patients and 16 healthy volunteers. They found that ADCs of MR imaging patterns in patients with MM differed significantly (normal MM pattern: 0.360×10^{-3} mm² s⁻¹, focal MM pattern: 1.046×10^{-3} mm² s⁻¹, diffuse MM pattern: 0.770×10^{-3} mm² s⁻¹). The mean ADC of normal bone marrow in the age-matched healthy control group was 0.325×10^{-3} mm² s⁻¹ (range, 0.152 – 0.542×10^{-3} mm² s⁻¹) and was significantly different from the ADCs of both the diffuse and the focal MM patterns, but not from those of the normal MM pattern.³ In a study by Bourillion et al, who performed IVIM DWI in 27 patients with MM, the mean ADC of 46 focal lesions, obtained by using only b -values of 0 and 800 s mm⁻², was 0.892×10^{-3} mm² s⁻¹.⁴

There are multiple sources of variance when measuring ADCs of normal and abnormal bone marrow, including factors related to the MR imaging protocol itself.⁵⁶ ADC calculations are particularly influenced by the choice of b -values as well as by other

technical factors, such as the method of fat suppression.³⁰ There is currently no clear consensus on the optimal number and scale of *b*-value selection for the evaluation of patients with plasma cell neoplasms. In some studies, the lowest *b*-value used is 0, whereas other investigators prefer a value of 50–100 s mm⁻², to limit perfusion contamination. Although ultrahigh *b*-values around 1400 s mm⁻² have been found to optimize contrast between normal and abnormal marrow, they can produce images with poor signal and increased distortion.²⁴ In most DWI studies of MM, the *b*-value of the highest diffusion-sensitizing gradient is set between 750 and 1000 s mm⁻², in line with expected ADCs from myeloma-infiltrated marrow.^{3,6,16,24,30,53,54,57} Moreover, some authors use multiple *b*-values in their MRI protocol for MM evaluation, while others typically use only two. It would, therefore, be commendable that expert recommendations for standardizing the use of bone marrow DWI (both spinal and WB) in plasma cell dyscrasias are developed. The recently published recommendations for the acquisition, interpretation, and reporting of WB MRI scans (including DWI) for use in males with advanced prostate cancer are a good example of this growing need for standardization of advanced imaging techniques in oncology.⁵⁸

Even considering these caveats, the above-mentioned ADCs are quite consistent across multiple studies and underline the fact that DWI is a relatively robust technique with great potential for the evaluation of MM patients. First, DWI increases lesion conspicuity with qualitative assessment, and can further increase diagnostic confidence for focal lesions with the use of ADC measurements. The IMWG currently recommends that SMM patients with equivocal focal lesions on conventional MRI should repeat the exam after 3 to 6 months, and in cases of MRI progression, should be considered as symptomatic patients who need therapy.⁵ DWI, especially with ADC calculations, may assist characterization in cases of equivocal findings on other pulse sequences and this potential additive value of DWI could possibly lead to more SMM patients meeting criteria for symptomatic disease requiring treatment. Future studies should investigate the important question, whether patients with a positive MRI based on the DWI sequence are at increased risk for earlier progression. In symptomatic MM, it has been shown that the number of focal lesions detected on conventional MRI has prognostic implications and again the value of the increased sensitivity of DWI, especially as part of a WB protocol, in this setting merits further assessment.⁵⁹ Second, DWI can play a role in the better definition of the diffuse MRI pattern of involvement in MM. The diffuse pattern is mostly diagnosed with the use of qualitative criteria on morphologic sequences, but there is no consensus on these criteria across studies.^{46–48,60–63} This is probably one of the reasons for discrepant results in the literature as to which MRI pattern confers a worse prognosis in patients with newly diagnosed MM.^{64–66} Moreover, the diffuse pattern (as opposed to the focal pattern) has not been incorporated in the updated criteria for the diagnosis of symptomatic myeloma.⁴³ WBDWI has been shown to be more sensitive than fluorodeoxyglucose PET-CT in detecting diffuse disease, by visually assessing the inverted grayscale maximum intensity projection images with *b*-value 800–900 s mm⁻².⁵⁷ Quantitative DWI with ADC

measurements may help define the diffuse pattern more objectively, facilitating comparison of results. In a recent study using spinal DWI with multiple *b*-values, an ADC greater than $0.597 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$ showed 96% sensitivity and 100% specificity for diagnosis of a diffuse MRI pattern (vs a normal MRI pattern) in patients with newly diagnosed MM, using morphological MRI criteria as a reference for pattern assignment.³ Similarly, Bourillon et al performed IVIM DWI in patients with MM and found that those with diffuse marrow involvement (defined in their study as maximal bone marrow enhancement above 100% on WBDCE MRI) had significantly different ADC compared to patients with maximal bone marrow enhancement below 100% (0.673 vs $0.543 \text{ mm}^2 \text{ s}^{-1}$ respectively). In the same study, IVIM parameter molecular diffusion coefficient *D* was also significantly higher in the diffuse marrow involvement group.⁴ Third, DWI could play an important role in therapy response assessment. PET-CT is the only imaging modality incorporated in the most recent IMWG consensus criteria for response and minimal residual disease assessment.⁶⁷ This is based on studies showing that among patients with a complete response, those with a positive PET scan had an inferior outcome.⁶⁸

MRI findings of response to therapy include decrease in size and number of focal lesions, appearance of a fatty halo with high *T*₁ signal at the periphery of focal lesions, and conversion of a diffuse pattern to a normal, focal or variegated pattern.⁶⁹ Morphological MRI criteria are, however, not adequate for response assessment since nonviable focal marrow lesions may persist after treatment even in good responders and no broad consensus exists on the value of these criteria.⁷⁰ These drawbacks of conventional MRI in assessing response in myeloma patients can potentially be overcome by a functional technique such as DWI, which can detect treatment-induced changes in cellularity. Several studies of different tumor types have shown that ADC values increase in good responders, and that this increase can occur as early as 1–4 weeks after treatment initiation.⁵⁶ These changes of ADCs reflect increases in the diffusivity of water molecules as a result of tumor cell death and increased extracellular spaces.³¹ In the case of MM, changes of ADCs following treatment are more complicated due to the particular cellular composition and dynamic nature of bone marrow, and depend highly on the timing of the examination and on the pre-treatment pattern of involvement (normal, focal, diffuse or combined). Responding focal lesions will exhibit an initial increase in ADCs from baseline, due to tumor necrosis, which can be appreciated very early following initiation of treatment (Figure 8). This is followed by a decrease in ADC values, due to reappearance of fatty marrow and, in some instances, possibly some degree of sclerosis.^{31,71} Rarely, some necrotic focal lesions may not exhibit total fatty marrow conversion and will display persistent, very elevated ADC values (and very high signal on high *b*-value images due to *T*₂ shine through). In the case of a diffuse or normal-appearing pre-treatment MRI pattern, ADC changes are more unpredictable and probably very time-dependent, since multiple factors can influence ADC values either way. Thus, in responders, the increase in ADCs caused by tumor cell death, edema and red marrow reversion during the phase of hematopoietic recovery (further enhanced in the case of administration of hematopoietic growth

Figure 8. b 750 image (a) and ADC map (b) from a spinal DWI exam of a 65-year-old male with newly diagnosed multiple myeloma. A focal lesion of T12 appears with high signal on the high b -value image and had an ADC value of $0.990 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$ (arrows). Corresponding images (c, d) from a second DWI exam performed 4 weeks after initiation of treatment with VCD (bortezomib, cyclophosphamide, dexamethasone). The ADC value of the T12 focal lesion (arrow) increased to $1.460 \times 10^{-3} \text{ mm}^2 \text{ s}^{-1}$, indicative of response. International Myeloma Working Group response status at 95 days post-autologous stem cell transplantation was very good partial response (VGPR). ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

factors) is counteracted by the reappearance of fatty marrow. In a study of 20 MM patients, responders showed an initial increase in ADCs at 4 to 6 weeks post-treatment, followed by a decrease at 20 weeks.⁵⁴ Considering the above, the whole body approach seems compelling, since it can combine information on ADCs from areas with diffuse infiltration as well as focal lesions, in effect assessing quantitatively the tumor burden before and after therapy. Giles *et al* used ADC histograms derived from a semi-automated segmentation technique with WBDWI of MM patients, in order to quantify disease throughout the skeleton before and at a median of 13 weeks after treatment. They found that the mean ADC increased in 95% of responders and decreased in all non-responders with a 3.3% increase identifying response with 90% sensitivity and 100% specificity. The authors also reported good results of response assessment with qualitative analysis of the high b -value images, using standardized observer scores.⁶ In a very recent study, Latifoltojar *et al* evaluated various WBMRI quantitative parameters for response assessment in 21 MM patients, including ADC and signal fat fraction, the latter derived from signal intensity of focal lesions measured on fat-only and water-only Dixon images.⁷² Both ADC and fat fraction significantly increased in myeloma lesions responding to treatment, but the authors found that lesion fat fraction was the best discriminator of myeloma treatment response. Although these encouraging results need to be validated in more prospective studies with larger cohorts, they seem to suggest that a WBMRI protocol incorporating both DWI and Dixon imaging may provide valuable complementary information for treatment assessment in MM within reasonable scanning times. Of note, the second WBMRI in this study was performed after the end of the second cycle of chemotherapy, which highlights the fact that no consensus currently exists on the most optimal time point for the repeat MRI. Evaluation of reproducibility across centers

and different vendors, standardization of technique, and implementation in large prospective studies are needed for WBMRI and WBDWI in particular, to gain acceptance as an imaging biomarker for response in MM (Figure 9). Direct comparison with PET-CT should be a priority, since the latter is currently

Figure 9. Coronal whole body inverted grayscale b 800 diffusion-weighted images in a 74-year-old male with multiple myeloma. At diagnosis, (a) there is widespread diffuse marrow infiltration with superimposed focal lesions. After 3 cycles of chemotherapy treatment with VMP (bortezomib, melphalan and prednisone) almost complete normalization of marrow signal is noted (b).

the imaging modality of choice for the more accurate definition of a complete response.

Dynamic contrast-enhanced MRI in MM

Angiogenesis plays a major role in the development of MM from its precursor forms, MGUS and SMM. Studies have shown a statistically significant gradual increase of MVD along the disease spectrum of plasma cell dyscrasias from MGUS to SMM, newly diagnosed MM, and finally relapsed MM, in accordance with the so-called angiogenic switch theory.^{73,74} Complex interactions between plasma cells and the bone marrow microenvironment, mediated by angiogenic and anti-angiogenic molecules, are implicated in the promotion of bone marrow angiogenesis.⁷⁵ A direct link between enhanced angiogenesis and the development of myeloma-related bone disease has been suggested.⁷⁶ Moreover, increased angiogenesis is a marker of poor survival in MM.⁷³ Using conventional MRI sequences, Mouloupoulos et al found that a diffuse MRI pattern correlates with increased angiogenesis (as assessed by MVD in trephine biopsies) and inferior survival.⁶⁵ DCE-MRI provides a non-invasive method for assessing the angiogenetic state of the bone marrow in myeloma patients, at the time of diagnosis as well as following treatment. It has the advantage of evaluating a large volume of the bone marrow, overcoming the sampling bias of trephine biopsies.² For DCE-MRI in myeloma patients, a fat-suppressed T_1 weighted ultrafast gradient echo sequence of the thoracolumbar or lumbar spine (more rarely of the pelvis) is usually acquired. Semi-quantitative perfusion parameters can be extracted from TICs, by drawing ROIs. DCE-MRI studies in patients with plasma cell dyscrasias have revealed a gradual increase of microcirculation from MGUS to SMM and MM.^{33,77} This can be appreciated by visual assessment of the relevant TICs (Figure 10). In MM, the curve is typically characterized by an early, steep and high rise, which reflects the pronounced neovascularity of the infiltrated marrow, followed by an early and pronounced washout,

as a result of a small interstitial compartment.³⁵ TICs in patients with SMM vary, but in most cases, they display a relatively flat enhancement slope and a low maximum enhancement followed by a very slow washout.

In most studies of DCE-MRI in myeloma patients, quantitative analysis of the dynamic data using either the Brix or the Tofts pharmacokinetic model has been performed, with extraction of parameters reflecting blood volume and vascular permeability. A few studies have evaluated the prognostic value of DCE-MRI in MGUS, SMM and MM as well as its role in response assessment. A recent study using the Brix model examined the prognostic significance of the quantitative perfusion parameters amplitude A (reflecting blood volume) and exchange rate constant K_{ep} (reflecting vessel wall permeability), in patients with MGUS and SMM. Patients with SMM and an amplitude A above the cut-off point of 0.89 arbitrary units, had a 2 year progression rate into symptomatic disease of 80%. For patients with MGUS, no prognostic effect of either parameter could be found.² In an earlier study, the same group had also observed perfusion parameters indicating increased angiogenesis in occasional patients with MGUS or SMM.⁷⁷ These results are encouraging, but more studies are needed to validate the role of DCE-MRI in the identification of high-risk SMM patients. The prognostic role of DCE-MRI in MM has also been evaluated in a small number of studies. One previous study, in patients with progressive or relapsed MM, had shown that amplitude A derived from DCE-MRI of the lumbar spine could predict for event-free survival but not for overall survival (OS).⁷⁸ However, this has not been confirmed in patients with newly-diagnosed MM, where the same group found no correlation of DCE-MRI amplitude A with progression-free survival, although high median values of amplitude A did predict shorter OS.⁷⁹ With regard to post-therapeutic assessment, previous studies have shown that response in MM patients is accompanied by a reduction of MVD.^{80,81} DCE-MRI derived parameters, which have been shown to correlate well with MVD, could therefore, be used as non-invasive biomarkers of response in MM.⁸² In the largest prospective study investigating DCE-MRI in this setting, quantitative DCE-MRI was performed in 91 patients before systemic therapy and at a median time of 14 months after the first MRI. Responders exhibited significantly lower values for amplitude A and k_{ep} compared with nonresponders after therapy and higher median k_{ep} values at the second MRI were associated with shorter OS. Interestingly, patients not responding to therapy showed higher amplitude A values before the start of therapy compared with responders, suggesting prognostic implications for the baseline DCE-MRI exam.⁸³ One group has used a low temporal resolution, whole body DCE MRI protocol to assess treatment response in patients with MM semiquantitatively. Bone marrow maximum enhancement differed significantly between good and poor responders after completion of induction chemotherapy. Regarding focal lesions, the timing of maximum enhancement was significantly delayed in good responders compared with poor responders.⁴ It remains to be seen whether DCE-MRI can become a useful clinical tool in the management of patients with plasma cell neoplasms,

Figure 10. Fitted time-signal intensity curves from DCE-MRI exams of the lumbar spine in a patient with smoldering multiple myeloma (green curve) and a different patient with symptomatic multiple myeloma (red curve). The curve of the patient with symptomatic multiple myeloma was derived from a region of interest drawn on a focal lesion, and displays an early, very steep rise, with high maximum enhancement and pronounced contrast washout. DCE, dynamic contrast-enhanced.

especially in view of the growing use of DWI, WBDWI and IVIM, all techniques that do not require the administration of intravenous contrast material.

CONCLUSIONS

DWI, WBDWI and DCE-MRI are being increasingly used in patients with plasma cell neoplasms. There is a growing body of

evidence that the additional functional information related to tumor cellularity and angiogenesis that they provide, may assist diagnosis of marrow infiltration and could have prognostic value in both SMM and MM patients. Moreover, by providing reliable MRI quantitative biomarkers, these techniques may play a more important role in the future for the better definition of treatment response criteria in patients with MM.

REFERENCES

1. Padhani AR, Koh DM, Collins DJ. Whole-body diffusion-weighted MR imaging in cancer: current status and research directions. *Radiology* 2011; **261**: 700–18. doi: <https://doi.org/10.1148/radiol.11110474>
2. Hillengass J, Ritsch J, Merz M, Wagner B, Kunz C, Hielscher T, et al. Increased microcirculation detected by dynamic contrast-enhanced magnetic resonance imaging is of prognostic significance in asymptomatic myeloma. *Br J Haematol* 2016; **174**: 127–35. doi: <https://doi.org/10.1111/bjh.14038>
3. Koutoulidis V, Fontara S, Terpos E, Zagouri F, Matsaridis D, Christoulas D, et al. Quantitative diffusion-weighted imaging of the bone marrow: an adjunct tool for the diagnosis of a diffuse MR imaging pattern in patients with multiple myeloma. *Radiology* 2017; **282**: 484–93. doi: <https://doi.org/10.1148/radiol.2016160363>
4. Bourillon C, Rahmouni A, Lin C, Belhadj K, Beaussart P, Vignaud A, et al. Intravoxel incoherent motion diffusion-weighted imaging of multiple myeloma lesions: correlation with whole-body dynamic contrast agent-enhanced MR imaging. *Radiology* 2015; **277**: 773–83. doi: <https://doi.org/10.1148/radiol.2015141728>
5. Dimopoulos MA, Hillengass J, Usmani S, Zamagni E, Lentzsch S, Davies FE, et al. Role of magnetic resonance imaging in the management of patients with multiple myeloma: a consensus statement. *J Clin Oncol* 2015; **33**: 657–64. doi: <https://doi.org/10.1200/JCO.2014.57.9961>
6. Giles SL, Messiou C, Collins DJ, Morgan VA, Simpkin CJ, West S, et al. Whole-body diffusion-weighted MR imaging for assessment of treatment response in myeloma. *Radiology* 2014; **271**: 785–94. doi: <https://doi.org/10.1148/radiol.13131529>
7. Le Bihan D, Iima M. Diffusion magnetic resonance imaging: what water tells us about biological tissues. *PLoS Biol* 2015; **13**: e1002203. doi: <https://doi.org/10.1371/journal.pbio.1002203>
8. Koh DM, Collins DJ. Diffusion-weighted MRI in the body: applications and challenges in oncology. *AJR Am J Roentgenol* 2007; **188**: 1622–35. doi: <https://doi.org/10.2214/AJR.06.1403>
9. Khoo MM, Tyler PA, Saifuddin A, Padhani AR. Diffusion-weighted imaging (DWI) in musculoskeletal MRI: a critical review. *Skeletal Radiol* 2011; **40**: 665–81. doi: <https://doi.org/10.1007/s00256-011-1106-6>
10. Padhani AR, Liu G, Koh DM, Chenevert TL, Thoeny HC, Takahara T, et al. Diffusion-weighted magnetic resonance imaging as a cancer biomarker: consensus and recommendations. *Neoplasia* 2009; **11**: 102–25. doi: <https://doi.org/10.1593/neo.81328>
11. Guo AC, Cummings TJ, Dash RC, Provenzale JM. Lymphomas and high-grade astrocytomas: comparison of water diffusibility and histologic characteristics. *Radiology* 2002; **224**: 177–83. doi: <https://doi.org/10.1148/radiol.2241010637>
12. Le Bihan D, Breton E, Lallemand D, Aubin ML, Vignaud J, Laval-Jeantet M. Separation of diffusion and perfusion in intravoxel incoherent motion MR imaging. *Radiology* 1988; **168**: 497–505. doi: <https://doi.org/10.1148/radiology.168.2.3393671>
13. Teruel JR, Goa PE, Sjøbakk TE, Østlie A, Fjøsne HE, Bathen TF. A simplified approach to measure the effect of the microvasculature in diffusion-weighted MR imaging applied to breast tumors: preliminary results. *Radiology* 2016; **281**: 373–81. doi: <https://doi.org/10.1148/radiol.2016151630>
14. Lecouvet FE, El Mouedden J, Collette L, Coche E, Danse E, Jamar F, et al. Can whole-body magnetic resonance imaging with diffusion-weighted imaging replace Tc 99m bone scanning and computed tomography for single-step detection of metastases in patients with high-risk prostate cancer? *Eur Urol* 2012; **62**: 68–75. doi: <https://doi.org/10.1016/j.eururo.2012.02.020>
15. Horger M, Claussen C, Kramer U, Fenchel M, Lichy M, Kaufmann S. Very early indicators of response to systemic therapy in lymphoma patients based on alterations in water diffusivity—a preliminary experience in 20 patients undergoing whole-body diffusion-weighted imaging. *Eur J Radiol* 2014; **83**: 1655–64. doi: <https://doi.org/10.1016/j.ejrad.2014.05.027>
16. Giles SL, deSouza NM, Collins DJ, Morgan VA, West S, Davies FE, et al. Assessing myeloma bone disease with whole-body diffusion-weighted imaging: comparison with X-ray skeletal survey by region and relationship with laboratory estimates of disease burden. *Clin Radiol* 2015; **70**: 614–21. doi: <https://doi.org/10.1016/j.crad.2015.02.013>
17. Bray TJP, Singh S, Latifoltojar A, Rajesparan K, Rahman F, Narayanan P, et al. Diagnostic utility of whole body dixon MRI in multiple myeloma: a multi-reader study. *PLoS One* 2017; **12**: e0180562. doi: <https://doi.org/10.1371/journal.pone.0180562>
18. Latifoltojar A, Hall-Craggs M, Rabin N, Popat R, Bainbridge A, Dikaïos N, et al. Whole body magnetic resonance imaging in newly diagnosed multiple myeloma: early changes in lesional signal fat fraction predict disease response. *Br J Haematol* 2017; **176**: 222–33. doi: <https://doi.org/10.1111/bjh.14401>
19. Ricci C, Cova M, Kang YS, Yang A, Rahmouni A, Scott WW, et al. Normal age-related patterns of cellular and fatty bone marrow distribution in the axial skeleton: MR imaging study. *Radiology* 1990; **177**: 83–8. doi: <https://doi.org/10.1148/radiology.177.1.2399343>
20. Mouloupoulos LA, Koutoulidis V. *Bone marrow MRI: a pattern-based approach*. Milan: Springer; 2015.
21. Vande Berg BC, Malghe J, Lecouvet FE, Maldague B. Magnetic resonance imaging of the normal bone marrow. *Skeletal Radiol* 1998; **27**: 471–83. doi: <https://doi.org/10.1007/s002560050423>
22. Kricun ME. Red-yellow marrow conversion: its effect on the location of some solitary bone lesions. *Skeletal Radiol* 1985; **14**: 10–19. doi: <https://doi.org/10.1007/BF00361188>

23. Nonomura Y, Yasumoto M, Yoshimura R, Haraguchi K, Ito S, Akashi T, et al. Relationship between bone marrow cellularity and apparent diffusion coefficient. *J Magn Reson Imaging* 2001; **13**: 757–60. doi: <https://doi.org/10.1002/jmri.1105>
24. Messiou C, Collins DJ, Morgan VA, Desouza NM. Optimising diffusion weighted MRI for imaging metastatic and myeloma bone disease and assessing reproducibility. *Eur Radiol* 2011; **21**: 1713–8. doi: <https://doi.org/10.1007/s00330-011-2116-4>
25. Herrmann J, Krstin N, Schoennagel BP, Sornsakrin M, Derlin T, Busch JD, et al. Age-related distribution of vertebral bone-marrow diffusivity. *Eur J Radiol* 2012; **81**: 4046–9. doi: <https://doi.org/10.1016/j.ejrad.2012.03.033>
26. Griffith JF, Yeung DK, Antonio GE, Wong SY, Kwok TC, Woo J, et al. Vertebral marrow fat content and diffusion and perfusion indexes in women with varying bone density: MR evaluation. *Radiology* 2006; **241**: 831–8. doi: <https://doi.org/10.1148/radiol.2413051858>
27. Tang GY, Lv ZW, Tang RB, Liu Y, Peng YF, Li W, et al. Evaluation of MR spectroscopy and diffusion-weighted MRI in detecting bone marrow changes in postmenopausal women with osteoporosis. *Clin Radiol* 2010; **65**: 377–81. doi: <https://doi.org/10.1016/j.crad.2009.12.011>
28. Biffar A, Dietrich O, Sourbron S, Duerr HR, Reiser MF, Baur-Melnyk A. Diffusion and perfusion imaging of bone marrow. *Eur J Radiol* 2010; **76**: 323–8. doi: <https://doi.org/10.1016/j.ejrad.2010.03.011>
29. Herneth AM, Friedrich K, Weidekamm C, Schibany N, Krestan C, Czerny C, et al. Diffusion weighted imaging of bone marrow pathologies. *Eur J Radiol* 2005; **55**: 74–83. doi: <https://doi.org/10.1016/j.ejrad.2005.03.031>
30. Padhani AR, van Ree K, Collins DJ, D'Sa S, Makris A. Assessing the relation between bone marrow signal intensity and apparent diffusion coefficient in diffusion-weighted MRI. *AJR Am J Roentgenol* 2013; **200**: 163–70. doi: <https://doi.org/10.2214/AJR.11.8185>
31. Li SP, Padhani AR. Tumor response assessments with diffusion and perfusion MRI. *J Magn Reson Imaging* 2012; **35**: 745–63. doi: <https://doi.org/10.1002/jmri.22838>
32. Walker-Samuel S, Leach MO, Collins DJ. Evaluation of response to treatment using DCE-MRI: the relationship between initial area under the gadolinium curve (IAUGC) and quantitative pharmacokinetic analysis. *Phys Med Biol* 2006; **51**: 3593–602. doi: <https://doi.org/10.1088/0031-9155/51/14/021>
33. Dutoit JC, Vanderkerken MA, Verstraete KL. Value of whole body MRI and dynamic contrast enhanced MRI in the diagnosis, follow-up and evaluation of disease activity and extent in multiple myeloma. *Eur J Radiol* 2013; **82**: 1444–52. doi: <https://doi.org/10.1016/j.ejrad.2013.04.012>
34. Tofts PS, Kermode AG. Measurement of the blood-brain barrier permeability and leakage space using dynamic MR imaging. 1. Fundamental concepts. *Magn Reson Med* 1991; **17**: 357–67. doi: <https://doi.org/10.1002/mrm.1910170208>
35. Mouloupoulos LA, Maris TG, Papanikolaou N, Panagi G, Vlahos L, Dimopoulos MA. Detection of malignant bone marrow involvement with dynamic contrast-enhanced magnetic resonance imaging. *Ann Oncol* 2003; **14**: 152–8. doi: <https://doi.org/10.1093/annonc/mdg007>
36. Montazel JL, Divine M, Lepage E, Kobeiter H, Breil S, Rahmouni A. Normal spinal bone marrow in adults: dynamic gadolinium-enhanced MR imaging. *Radiology* 2003; **229**: 703–9. doi: <https://doi.org/10.1148/radiol.2293020747>
37. Griffith JF, Yeung DK, Antonio GE, Lee FK, Hong AW, Wong SY, et al. Vertebral bone mineral density, marrow perfusion, and fat content in healthy men and men with osteoporosis: dynamic contrast-enhanced MR imaging and MR spectroscopy. *Radiology* 2005; **236**: 945–51. doi: <https://doi.org/10.1148/radiol.2363041425>
38. Savvopoulou V, Maris TG, Koureas A, Gouliamos A, Mouloupoulos LA. Degenerative endplate changes of the lumbosacral spine: dynamic contrast-enhanced MRI profiles related to age, sex, and spinal level. *J Magn Reson Imaging* 2011; **33**: 382–9. doi: <https://doi.org/10.1002/jmri.22444>
39. Chen WT, Shih TT, Chen RC, Lo SY, Chou CT, Lee JM, et al. Vertebral bone marrow perfusion evaluated with dynamic contrast-enhanced MR imaging: significance of aging and sex. *Radiology* 2001; **220**: 213–8. doi: <https://doi.org/10.1148/radiology.220.1.r01j132213>
40. Breault SR, Heye T, Bashir MR, Dale BM, Merkle EM, Reiner CS, et al. Quantitative dynamic contrast-enhanced MRI of pelvic and lumbar bone marrow: effect of age and marrow fat content on pharmacokinetic parameter values. *AJR Am J Roentgenol* 2013; **200**: W297–W303. doi: <https://doi.org/10.2214/AJR.12.9080>
41. Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med* 2011; **364**: 1046–60. doi: <https://doi.org/10.1056/NEJMra1011442>
42. Dhodapkar MV, Sexton R, Waheed S, Usmani S, Papanikolaou X, Nair B, et al. Clinical, genomic, and imaging predictors of myeloma progression from asymptomatic monoclonal gammopathies (SWOG S0120). *Blood* 2014; **123**: 78–85. doi: <https://doi.org/10.1182/blood-2013-07-515239>
43. Rajkumar SV, Dimopoulos MA, Palumbo A, Blade J, Merlini G, Mateos MV, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol* 2014; **15**: e538–e548. doi: [https://doi.org/10.1016/S1470-2045\(14\)70442-5](https://doi.org/10.1016/S1470-2045(14)70442-5)
44. Dimopoulos M, Terpos E, Comenzo RL, Tosi P, Beksac M, Sezer O, et al. International myeloma working group consensus statement and guidelines regarding the current role of imaging techniques in the diagnosis and monitoring of multiple myeloma. *Leukemia* 2009; **23**: 1545–56. doi: <https://doi.org/10.1038/leu.2009.89>
45. Hillengass J, Mouloupoulos LA, Delorme S, Koutoulidis V, Mosebach J, Hielscher T, et al. Whole-body computed tomography versus conventional skeletal survey in patients with multiple myeloma: a study of the International Myeloma Working Group. *Blood Cancer J* 2017; **7**: e599. doi: <https://doi.org/10.1038/bcj.2017.78>
46. Libshitz HI, Malthouse SR, Cunningham D, MacVicar AD, Husband JE. Multiple myeloma: appearance at MR imaging. *Radiology* 1992; **182**: 833–7. doi: <https://doi.org/10.1148/radiology.182.3.1535904>
47. Mouloupoulos LA, Varma DG, Dimopoulos MA, Leeds NE, Kim EE, Johnston DA, et al. Multiple myeloma: spinal MR imaging in patients with untreated newly diagnosed disease. *Radiology* 1992; **185**: 833–40. doi: <https://doi.org/10.1148/radiology.185.3.1438772>
48. Stäbler A, Baur A, Bartl R, Munker R, Lamerz R, Reiser MF. Contrast enhancement and quantitative signal analysis in MR imaging of multiple myeloma: assessment of focal and diffuse growth patterns in marrow correlated with biopsies and survival rates. *AJR Am J Roentgenol* 1996; **167**: 1029–36. doi: <https://doi.org/10.2214/ajr.167.4.8819407>
49. Walker R, Barlogie B, Haessler J, Tricot G, Anaissie E, Shaughnessy JD, et al. Magnetic resonance imaging in multiple myeloma: diagnostic and clinical implications. *J Clin Oncol* 2007; **25**: 1121–8. doi: <https://doi.org/10.1200/JCO.2006.08.5803>
50. Hillengass J, Fechtner K, Weber MA, Bäuerle T, Ayyaz S, Heiss C, et al. Prognostic significance of focal lesions in whole-body

- magnetic resonance imaging in patients with asymptomatic multiple myeloma. *J Clin Oncol* 2010; **28**: 1606–10. doi: <https://doi.org/10.1200/JCO.2009.25.5356>
51. Kastritis E, Mouloupoulos LA, Terpos E, Koutoulidis V, Dimopoulos MA. The prognostic importance of the presence of more than one focal lesion in spine MRI of patients with asymptomatic (smoldering) multiple myeloma. *Leukemia* 2014; **28**: 2402–3. doi: <https://doi.org/10.1038/leu.2014.230>
 52. Mouloupoulos LA, Dimopoulos MA, Smith TL, Weber DM, Delasalle KB, Libshitz HI, et al. Prognostic significance of magnetic resonance imaging in patients with asymptomatic multiple myeloma. *J Clin Oncol* 1995; **13**: 251–6. doi: <https://doi.org/10.1200/JCO.1995.13.1.251>
 53. Dutoit JC, Claus E, Offner F, Noens L, Delanghe J, Verstraete KL. Combined evaluation of conventional MRI, dynamic contrast-enhanced MRI and diffusion weighted imaging for response evaluation of patients with multiple myeloma. *Eur J Radiol* 2016; **85**: 373–82. doi: <https://doi.org/10.1016/j.ejrad.2015.11.040>
 54. Messiou C, Giles S, Collins DJ, West S, Davies FE, Morgan GJ, et al. Assessing response of myeloma bone disease with diffusion-weighted MRI. *Br J Radiol* 2012; **85**: e1198–e1203. doi: <https://doi.org/10.1259/bjr/52759767>
 55. Pearce T, Philip S, Brown J, Koh DM, Burn PR. Bone metastases from prostate, breast and multiple myeloma: differences in lesion conspicuity at short-tau inversion recovery and diffusion-weighted MRI. *Br J Radiol* 2012; **85**: 1102–6. doi: <https://doi.org/10.1259/bjr/30649204>
 56. Taouli B, Beer AJ, Chenevert T, Collins D, Lehman C, Matos C, et al. Diffusion-weighted imaging outside the brain: Consensus statement from an ISMRM-sponsored workshop. *J Magn Reson Imaging* 2016; **44**: 521–40. doi: <https://doi.org/10.1002/jmri.25196>
 57. Pawlyn C, Fowkes L, Otero S, Jones JR, Boyd KD, Davies FE, et al. Whole-body diffusion-weighted MRI: a new gold standard for assessing disease burden in patients with multiple myeloma? *Leukemia* 2016; **30**: 1446–8. doi: <https://doi.org/10.1038/leu.2015.338>
 58. Padhani AR, Lecouvet FE, Tunariu N, Koh DM, De Keyser F, Collins DJ, et al. METastasis reporting and data system for prostate cancer: practical guidelines for acquisition, interpretation, and reporting of whole-body magnetic resonance imaging-based evaluations of multiorgan involvement in advanced prostate cancer. *Eur Urol* 2017; **71**: 81–92. doi: <https://doi.org/10.1016/j.eururo.2016.05.033>
 59. Usmani SZ, Mitchell A, Waheed S, Crowley J, Hoering A, Petty N, et al. Prognostic implications of serial 18-fluorodeoxyglucose emission tomography in multiple myeloma treated with total therapy 3. *Blood* 2013; **121**: 1819–23. doi: <https://doi.org/10.1182/blood-2012-08-451690>
 60. Baur A, Stähler A, Nagel D, Lamerz R, Bartl R, Hiller E, et al. Magnetic resonance imaging as a supplement for the clinical staging system of Durie and Salmon? *Cancer* 2002; **95**: 1334–45. doi: <https://doi.org/10.1002/cncr.10818>
 61. Lecouvet FE, Vande Berg BC, Michaux L, Malghem J, Maldague BE, Jamart J, et al. Stage III multiple myeloma: clinical and prognostic value of spinal bone marrow MR imaging. *Radiology* 1998; **209**: 653–60. doi: <https://doi.org/10.1148/radiology.209.3.9844655>
 62. Mai EK, Hielscher T, Kloth JK, Merz M, Shah S, Hillengass M, et al. Association between magnetic resonance imaging patterns and baseline disease features in multiple myeloma: analyzing surrogates of tumour mass and biology. *Eur Radiol* 2016; **26**: 3939–48. doi: <https://doi.org/10.1007/s00330-015-4195-0>
 63. Song MK, Chung JS, Lee JJ, Min CK, Ahn JS, Lee SM, et al. Magnetic resonance imaging pattern of bone marrow involvement as a new predictive parameter of disease progression in newly diagnosed patients with multiple myeloma eligible for autologous stem cell transplantation. *Br J Haematol* 2014; **165**: 777–85. doi: <https://doi.org/10.1111/bjh.12820>
 64. Filonzi G, Mancuso K, Zamagni E, Nanni C, Spinnato P, Cavo M, et al. A comparison of different staging systems for multiple myeloma: can the MRI pattern play a prognostic role? *AJR Am J Roentgenol* 2017; **209**: 152–8. doi: <https://doi.org/10.2214/AJR.16.17219>
 65. Mouloupoulos LA, Dimopoulos MA, Christoulas D, Kastritis E, Anagnostou D, Koureas A, et al. Diffuse MRI marrow pattern correlates with increased angiogenesis, advanced disease features and poor prognosis in newly diagnosed myeloma treated with novel agents. *Leukemia* 2010; **24**: 1206–12. doi: <https://doi.org/10.1038/leu.2010.70>
 66. Mouloupoulos LA, Dimopoulos MA, Kastritis E, Christoulas D, Gkotzamanidou M, Roussou M, et al. Diffuse pattern of bone marrow involvement on magnetic resonance imaging is associated with high risk cytogenetics and poor outcome in newly diagnosed, symptomatic patients with multiple myeloma: a single center experience on 228 patients. *Am J Hematol* 2012; **87**: 861–4. doi: <https://doi.org/10.1002/ajh.23258>
 67. Kumar S, Paiva B, Anderson KC, Durie B, Landgren O, Moreau P, et al. International Myeloma Working Group consensus criteria for response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol* 2016; **17**: e328–e346. doi: [https://doi.org/10.1016/S1470-2045\(16\)30206-6](https://doi.org/10.1016/S1470-2045(16)30206-6)
 68. Zamagni E, Nanni C, Mancuso K, Tacchetti P, Pezzi A, Pantani L, et al. PET/CT improves the definition of complete response and allows to detect otherwise unidentifiable skeletal progression in multiple myeloma. *Clin Cancer Res* 2015; **21**: 4384–90. doi: <https://doi.org/10.1158/1078-0432.CCR-15-0396>
 69. Mouloupoulos LA, Dimopoulos MA, Alexanian R, Leeds NE, Libshitz HI. Multiple myeloma: MR patterns of response to treatment. *Radiology* 1994; **193**: 441–6. doi: <https://doi.org/10.1148/radiology.193.2.7972760>
 70. Bannas P, Hentschel HB, Bley TA, Treszl A, Eulenburger C, Derlin T, et al. Diagnostic performance of whole-body MRI for the detection of persistent or relapsing disease in multiple myeloma after stem cell transplantation. *Eur Radiol* 2012; **22**: 2007–12. doi: <https://doi.org/10.1007/s00330-012-2445-y>
 71. Horger M, Weisel K, Horger W, Mroue A, Fenchel M, Lichy M. Whole-body diffusion-weighted MRI with apparent diffusion coefficient mapping for early response monitoring in multiple myeloma: preliminary results. *AJR Am J Roentgenol* 2011; **196**: W790–W795. doi: <https://doi.org/10.2214/AJR.10.5979>
 72. Latifoltojar A, Hall-Craggs M, Bainbridge A, Rabin N, Popat R, Rismani A, et al. Whole-body MRI quantitative biomarkers are associated significantly with treatment response in patients with newly diagnosed symptomatic multiple myeloma following bortezomib induction. *Eur Radiol* 2017; **27**: 5325–36. doi: <https://doi.org/10.1007/s00330-017-4907-8>
 73. Rajkumar SV, Mesa RA, Fonseca R, Schroeder G, Plevak MF, Dispenzieri A, et al. Bone marrow angiogenesis in 400 patients with monoclonal gammopathy of undetermined significance, multiple myeloma, and primary amyloidosis. *Clin Cancer Res* 2002; **8**: 2210–6.
 74. Vacca A, Ribatti D, Roncali L, Ranieri G, Serio G, Silvestris F, et al. Bone marrow angiogenesis and progression in multiple

- myeloma. *Br J Haematol* 1994; **87**: 503–8. doi: <https://doi.org/10.1111/j.1365-2141.1994.tb08304.x>
75. Jakob C, Sterz J, Zavrski I, Heider U, Kleeberg L, Fleissner C, et al. Angiogenesis in multiple myeloma. *Eur J Cancer* 2006; **42**: 1581–90. doi: <https://doi.org/10.1016/j.ejca.2006.02.017>
 76. Tanaka Y, Abe M, Hiasa M, Oda A, Amou H, Nakano A, et al. Myeloma cell-osteoclast interaction enhances angiogenesis together with bone resorption: a role for vascular endothelial cell growth factor and osteopontin. *Clin Cancer Res* 2007; **13**: 816–23. doi: <https://doi.org/10.1158/1078-0432.CCR-06-2258>
 77. Hillengass J, Zechmann C, Bäuerle T, Wagner-Gund B, Heiss C, Benner A, et al. Dynamic contrast-enhanced magnetic resonance imaging identifies a subgroup of patients with asymptomatic monoclonal plasma cell disease and pathologic microcirculation. *Clin Cancer Res* 2009; **15**: 3118–25. doi: <https://doi.org/10.1158/1078-0432.CCR-08-2310>
 78. Hillengass J, Wasser K, Delorme S, Kiessling F, Zechmann C, Benner A, et al. Lumbar bone marrow microcirculation measurements from dynamic contrast-enhanced magnetic resonance imaging is a predictor of event-free survival in progressive multiple myeloma. *Clin Cancer Res* 2007; **13**: 2 Pt 1: 475–81. doi: <https://doi.org/10.1158/1078-0432.CCR-06-0061>
 79. Merz M, Moehler TM, Ritsch J, Bäuerle T, Zechmann CM, Wagner B, et al. Prognostic significance of increased bone marrow microcirculation in newly diagnosed multiple myeloma: results of a prospective DCE-MRI study. *Eur Radiol* 2016; **26**: 1404–11. doi: <https://doi.org/10.1007/s00330-015-3928-4>
 80. Oh HS, Choi JH, Park CK, Jung CW, Lee SI, Park Q, et al. Comparison of microvessel density before and after peripheral blood stem cell transplantation in multiple myeloma patients and its clinical implications: multicenter trial. *Int J Hematol* 2002; **76**: 465–70.
 81. Sezer O, Niemöller K, Eucker J, Jakob C, Kaufmann O, Zavrski I, et al. Bone marrow microvessel density is a prognostic factor for survival in patients with multiple myeloma. *Ann Hematol* 2000; **79**: 574–7. doi: <https://doi.org/10.1007/s002770000236>
 82. Nosàs-Garcia S, Moehler T, Wasser K, Kiessling F, Bartl R, Zuna I, et al. Dynamic contrast-enhanced MRI for assessing the disease activity of multiple myeloma: a comparative study with histology and clinical markers. *J Magn Reson Imaging* 2005; **22**: 154–62. doi: <https://doi.org/10.1002/jmri.20349>
 83. Merz M, Ritsch J, Kunz C, Wagner B, Sauer S, Hose D, et al. Dynamic contrast-enhanced magnetic resonance imaging for assessment of antiangiogenic treatment effects in multiple myeloma. *Clin Cancer Res* 2015; **21**: 106–12. doi: <https://doi.org/10.1158/1078-0432.CCR-14-1029>