Published in final edited form as:

J Vet Behav. 2019 ; 29: 4–10. doi:10.1016/j.jveb.2018.08.002.

Handler training improves decontamination of working canines with oil-based exposure in field conditions using disposable kits.

Ellie B. Powell^a, Gary A. Apgar^a, Eileen K. Jenkins^b, Stephen Y. Liang^c, and Erin B. Perry^{1,a}

^aSouthern Illinois University, Department of Animal Science Food & Nutrition, Carbondale, Illinois 62901

^bFirst Year Graduate Veterinary Education Program Public Health Activity – Fort Bragg Fort Bragg, North Carolina 28310

^cDivisions of Emergency Medicine and Infectious Diseases Washington University School of Medicine St. Louis, Missouri 63110

Abstract

Exposure to contaminants in the field is a reality for deployed canines. To date, there is no data evaluating the benefits of training for handlers associated with canine decontamination efforts. The objective of our work was to investigate the impact of handler training on the reduction of oil-based contaminants in working canines. Canine teams ($n = 10$) were randomly assigned to either TRAINED or UNTRAINED groups. Each team (handler and dog) in the TRAINED group received 30-minutes of interactive training using an illustrated guide on proper utilization of equipment provided. Teams in the UNTRAINED group received the same equipment and illustrated guide but no interactive training. Decontamination efforts were measured using an oil-based pseudo-contaminant (GloGerm®, Moab, UT) topically applied to four anatomical locations: cranial neck, between the shoulder blades, left medial hindlimb and left hind paw with pre- and post-washing images collected from a fixed distance of 20 inches. Visual assessment of contaminant reduction was scored as follows: 0 = <24% contaminant reduction; 1 = 25–50% contaminant reduction; 2 = 51–75% contaminant reduction; and 3 = >76% contaminant reduction. No score discrepancies >1 were reported between reviewers. Trained handlers were more effective at contamination reduction ($P = .0093$) as compared to their untrained counterparts. These results indicate that handlers, when properly trained, can achieve reduction of oil-based contaminants with a disposable decontamination kit and a garden hose.

¹Corresponding author: erin.perry@siu.edu.

DISCLAIMER

The findings and conclusions in this manuscript represent the opinion of the authors and do not necessarily represent those of the U.S. Department of Homeland Security, Federal Emergency Management Agency, or the U.S. Department of Defense.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

decontamination; search-and-rescue; disaster response; working canines

INTRODUCTION

Large-scale disasters can result in the dispersal of potentially toxic chemicals and hazardous materials into the environment (Murphy et al., 2003; Soric et al., 2008). Floods, volcanic eruptions, earthquakes, landslides, hurricanes, tornadoes and man-made events (Binder et al., 1987; Otto et al., 2002) can release phenols, toxic petroleum products, asbestos-laden dust, heavy metals and a wide range of volatile organic solvents and other compounds that may pose harm to animal and human health (Murphy et al., 2003; Soric et al., 2008).

Although some efforts have been made to identify methods associated with decontamination following disasters, few data have identified effective techniques applicable to working canines deployed as part of a response team (Welte et al., 1991; Soric et al., 2008; Washington Department of Fish and Wildlife, 2009; Venable et al., 2017). Previous work has demonstrated current techniques are inadequate for complete removal of oil-based contaminants (Venable et al., 2017). In contrast, data from wildlife studies have reported complete contaminant removal, but suggest a protocol that is prohibitive due to the extensive amount of time required (Welte et al., 1991; Washington Department of Fish and Wildlife, 2009). These studies involve the use of a decontamination corridor (Figure 1) that is both labor intensive and unlikely to be available in the early hours of a disaster response.

Response time for effective decontamination is critical and should include efforts to reduce exposure, restrict absorption and enhance elimination (Jones et al., 2004). The handler may better achieve each of these mandates through rapid decontamination in the field. To date, no data is available regarding the efficacy on contaminant reduction associated with handler decontamination in the field. Therefore, the objectives of our study were to assess the contaminant reduction potential of handlers using disposable field kits for canine decontamination under field conditions and investigate the impact of training on handler decontamination effectiveness.

MATERIALS & METHODS

Animals

This research was approved by the Southern Illinois University Institutional Animal Care and Use Committee (Protocol 15–032). Canine teams ($n = 10$) from the National Disaster Search Dog Foundation (Santa Paula, California) were randomly assigned to TRAINED or UNTRAINED groups for participation in a two-day study. Breeds included German shepherds ($n = 2$), Labrador retrievers ($n = 5$), Belgian malinois ($n = 1$) and mixed breeds ($n = 2$). Each team (handler and dog) in the TRAINED group received 30-minutes of interactive instructions using an illustrated guide on proper utilization of equipment provided. Teams in the UNTRAINED group received the same equipment and illustrated

guide but no interactive instructions. No handler had any knowledge or use of the decontamination field kit prior to the study.

Field kit

Both groups received identical field decontamination kits (Figure 2A), designed by the researchers for the purpose of this study. Each field decontamination kit contained a rubber grooming brush (ZoomGroom®, Salisbury, UK), two disposable nylon leashes, disposable towels for drying, Johnson and Johnson's Head-To-Toe-Baby Wash (Proctor & Gamble, Cincinnati, OH), cotton balls, a package of baby wipes (Huggies®, Neenah, WI), disposable gloves and an illustrated guide (Figure 2B) with step-by-step procedures for decontamination. Instructions for use of the field decontamination kit and illustrated guide (Figure 3) were developed according to researchers' recommendation.

Procedure

An oil-based contaminant (GloGerm®, Moab, Utah) was topically applied to each canine at four anatomic sites: 1) cranial neck; 2) between the shoulders blades; 3) left medial hindlimb; and 4) left hind paw (Figure 4). Equal pressure was applied during application to maximize consistency of contaminant contact. Images were collected before (Figure 5A) and after decontamination efforts (Figure 5B) with a Nikon D3400 DSLR Camera with 18–55mm VR lens and a UV flashlight provided by GloGerm® at a fixed position of 20 inches. Care was taken to ensure that room lighting and camera distance remained the same for all images.

Photos were analyzed using two methods 1) assignment of visual score; 2) measurement of fluorescent reduction. Scores were assigned on visual inspection of each of the four anatomic sites, using two blinded reviewers with a >80% agreement as previously described with no score discrepancies of >1 between reviewers (Venable et al., 2017). Visual score assignments were allotted as follows: 0 = <24% contaminant reduction; 1 = 25–50% contaminant reduction; 2 = 51–75% contaminant reduction; and 3 = >76% contaminant reduction adapted from prior work (Lee et al., 2014). Additionally, contaminant reduction was measured quantitatively at each site using the software program, ImageJ (NIH, Bethesda, MD) and by calculating the percent change in fluorescence using the equation $\left(\frac{PRE - POST}{PRE} \times 100\right) = \Delta\%$.

Categorical data (visual scores) were analyzed as a Chi square using PROC FREQ (SAS, v 9.4). Numerical data (%) were compared using a one-way ANOVA. Following calculations of both data sets, PROC REG was used to identify relatedness. The P value for all analysis was set at 0.05 and effective decontamination was defined as a reduction in fluorescence by at least 50% or a contaminant reduction score of ≥ 2 (Venable et al., 2017).

Cross-contamination was assessed visually in the kennel area for five canines and their handlers. Canine kennel areas were inspected 24-hours following decontamination, while the handlers were inspected immediately following field decontamination of their canines. Private kennel space was maintained for each dog to simulate kennel stations at a typical base of operations during deployments.

RESULTS

Efficacy of decontamination

Training improved contaminant reduction compared to their untrained counterparts ($P = 0.0093$) as shown in Figure 6 as determined by visually assigned scores. Similarly, measured fluorescent reduction also indicated improved contaminant reduction associated with the TRAINED vs. UNTRAINED group ($P = 0.0042$) (Figure 7).

Anatomic site also impacted contaminant reduction ($P = 0.0055$) as in Figure 8. The shoulder blades were most effectively decontaminated (i.e., greatest amount of reduction) at 50.3%, followed by the left hind paw at 40%. Both the paw and shoulder blades were more effectively decontaminated when compared to the cranial neck (29.2% reduction) and the left medial hindlimb (20.4% reduction).

Cross-contamination

Cross-contamination was defined as the presence of contaminant on environmental surfaces and was assessed using an ultraviolet flashlight (The Bramton Company LLC, Dallas, TX). Of the five canine kennels surveyed, all showed evidence of residue transfer (Figure 9). Specific areas of contamination included 60% of the canines' water buckets and beds, 80% of the kennel floors and walls and 40% of the kennel doors handles. Handlers checked for cross-contamination had residue transfer to their boots, pants, belts, shirts, watches and hats, suggesting cross-contamination from canine to handler.

Comparison between categorical and measured fluorescent reduction data

A linear regression analysis was performed and indicated relatedness ($P = 0.0126$) between the visual scores and measured reduction. Upon further analysis, utilizing a Spearman's correlation, a weak relatedness ($r_s = 0.2476$) as determined between both sources of data. r_s is defined as:

$$r_s = 1 - \frac{6 \sum D^2}{n(n^2 - 1)}$$

where the Spearman rank correlation coefficient, D is the difference between each pair of ranks of corresponding values of the variables X and Y , and n is the number of pairs of values in the sample.

DISCUSSION

According to the Urban Search and Rescue Veterinary Group, canine decontamination is a recognized component of Federal Emergency Management Agency (FEMA) Urban Search-and-Rescue (US&R) canine care and part of the expected set-up to maintain a clean base of operations environment (FEMA, 2017). Although efforts towards improved decontamination of working canines have been made, lack of available data and standardized protocols have inhibited opportunities for replication. Consequently, evidence-based guidelines are needed to protect canines and handlers when disasters strike.

Gordon (FEMA, 2012; 2017) reported attempted decontamination by federal teams deployed to hurricanes in 2012 and 2017. Decontamination stations were utilized with 63% – 100% of canines decontaminated (FEMA 2012; 2017). However, procedures were inconsistent and lacked adequate detail. Various combinations of water temperatures (cold or warm), cleansers (soap or Simple Green®) and baby wipes were utilized to decontaminate the canines with some teams reporting the use of water alone.

Other reports have included greater detail but indicate a lack of evidence regarding cleanser selection. In 2010, following an earthquake in Haiti, twenty-two US&R canines were deployed. Twenty-one of the canines underwent decontamination on all search days, while the remaining canine only received decontamination on half of the search days (FEMA, 2012). However, no corridor was set up for decontamination. Eighteen canines were decontaminated with soap and water, two canines had a combination of soap, water, moist wipes and eye flush, while the remaining two canines were decontaminated with water alone (FEMA, 2012). Following a 2014 landslide in Oso, Washington, the Washington National Guard established two mass decontamination units, and Washington Task Force 1 established a decontamination corridor to decontaminate the twenty-five canines deployed (Gordon, 2015). Decontamination was provided for canines after every shift. Water and dish soap were utilized, but veterinary medical stations also provided additional water, wipes and towels to further decontaminate the canines as needed (Gordon, 2015). Veterinary assessments revealed that the dish soap utilized caused dermatological issues, leading to scratching and skin damage. Therefore, an oatmeal-based shampoo formulated for canines was acquired and used for the remaining duration of the deployment to mitigate skin drying (Gordon, 2015). This report seems to suggest that repetitive decontamination, as is often necessary under deployment conditions, may require investigation so as to provide evidence-based recommendations for cleansers that are effective and safe.

Decontamination protocols for oil-based contaminant exposure exist for sea otters and waterfowl (Welte et al., 1991; Washington Department of Fish and Wildlife, 2009). Both utilize Dawn® (Proctor & Gamble, Cincinnati, OH) due to degreasing properties. In sea otters, multiple applications of diluted 5% solution were applied at washing tables. (Washington Department of Fish and Wildlife, 2009). Each wash table required four to six people to decontaminate the sea otters, as the detergent was massaged into the oiled fur and then rinsed off under moderate pressure (30–40 psi) with a spray nozzle (Washington Department of Fish and Wildlife, 2009). This procedure was repeated until all of the oil was lifted and removed from the otter's fur, requiring 40–60 minutes to complete (Washington Department of Fish and Wildlife, 2009). Following the wash procedure, a final rinse lasted an additional 40–60 minutes. Although this method has been deemed effective for wildlife, the length of time required for implementation makes this approach impractical for working canines in the field.

In addition to sea otter decontamination, the Washington Department of Fish and Wildlife also has protocols in place for waterfowl exposed to oil-based contaminants. Warm water (104°F) and Dawn® solution (2–15%) are used in conjunction with ten-gallon tubs in order to wash oil from feathers. Once again, the cleaning effort takes approximately 60-minutes (Welte et al., 1991). Clearly, these methods, although potentially effective, are not plausible

for use in the field where warm water may be unavailable and lengthy decontamination procedures are prohibitive. Decontamination procedures must proceed efficiently and rapidly in order to accommodate the influx of teams returning to base.

Some decontamination efforts have utilized Dawn® and Simple Green® as cleansers despite the lack of data for safety and efficacy in the canine. The use of Simple Green® to decontaminate canines is alarming. Simple Green® is a product approved for use on inert surfaces, not on animals. Soaps and detergents, also known as surface active agents, work as emulsifiers to surround and lift organic matter from hair, which can then be rinsed away with water (Soric et al., 2008). Although that chemical characteristic provides a sound theory for the use of dish soap, when Dawn® was compared to baby shampoo in a recent study, baby shampoo (Johnson & Johnson Head-To-Toe Baby Wash, Proctor & Gamble, Cincinnati, OH) was found to be more effective at reducing oil-based contaminants than Dawn® (Venable et al., 2017). These pilot data are the first to suggest that dish soap may be ineffective at oil-based contaminant reduction using standard decontamination techniques in working canines.

In a study conducted by the Sea World Research Institute, a decontamination protocol for otters was developed, utilizing diluted Dawn® solution (1:16 in water). The researchers designed a protocol that required a 40-minute wash, followed by a 40-minute rinse once there was no indication of oil residue on the fur of the otters (Davis et al., 1988). However, gas chromatography performed on fur samples taken from the abdomen (unoled) or the sternum (oiled) of otters revealed changes in squalene concentrations after use of Dawn® (Davis et al., 1988). Squalene is important in maintaining the water-repellent and insulating qualities of otter fur. Squalene was not restored in the oiled and cleaned fur after 7 days (Davis et al., 1988) suggesting dermatological impacts associated with the use of dish soap.

The impact of decontamination on canine skin is dependent on the type and frequency of cleanser used, as well as the health and composition of the skin to which the cleanser is applied. The use of surfactants in cleansers facilitates removal of dirt and soil. Unfortunately, the interaction of surfactant with proteins and lipids in mammalian skin can be detrimental to dermatologic health (Ananthapadmanabhan, 2004). Anionic surfactants, which typically have an alkaline pH, are notably more irritating to skin than cationic surfactants, which typically have a neutral to slightly acidic pH (Ananthapadmanabhan, 2004). Differences in cleanser structural properties and skin pH influence the amount and type of damage inflicted by surfactants interacting with proteins and lipids within the stratum corneum (Beranda, 2002; Ananthapadmanabhan, 2004).

In order to maximize effectiveness and minimize irritation, the pH of the cleanser should resemble the pH of the skin. Most products currently used for canine decontamination are shampoos formulated for human skin and hair, or dishwashing soap formulated to protect human skin. The pH of canine skin is slightly acidic to neutral, varies by body region and is influenced by excitement, but is not influenced by male or female sex (Meyer, 1991). While human skin typically has a pH of 5.5 – 6.5 (Chan 2012), the mean pH of canine skin can range from 5.86 to 7.42 in the unexcited state and 7.15 to 8.03 in the excited state (Meyer, 1991). Thus, the use of cleansers formulated for human skin may not be the most effective or safe product for decontamination of canine skin.

Regardless of the cleansing product used, repeated washing of skin can have deleterious effects. With repeated use, surfactants can cause a decrease in skin hydration and viscoelasticity, cause denaturation of skin proteins, and can alter skin lipid biosynthetic pathways (Ananthapadmanabhan, 2004). Repeated washing with dish soap caused increased epidermal irritation in search & rescue canines bathed daily (Gordon, 2015). The serial use of Dawn dish soap for canine decontamination resulted in mild to moderate epidermal irritation starting at a mean of 4.9 days (Mabry, 2015). The serial use of DermaLyte soap, a product marketed specifically for dogs, also resulted in mild to moderate epidermal irritation starting at a mean of 5.8 days (Mabry, 2015). More research is needed to determine the most effective and least irritating products to use for serial decontamination of canine skin. The ideal cleanser will rapidly and effectively remove a wide variety of contaminating substances, will not enhance cutaneous absorption or penetration of the contaminant, is readily rinsed or removed after use, does not damage skin, and is affordable and readily available (Chan, 2012).

One objective of this study was to determine the effect of training on contaminant reduction, utilizing a field kit. Handlers receiving no training failed to remove soiled collars from the canine's neck, thus the collar acted as a vector for reintroduction of contaminants. This is similar to prior work demonstrating a failure to remove soiled collars from canines during decontamination procedures (Venable et al., 2017). In addition, training also had an impact on the amount of contaminant reduction on the canine's paw. The paws are likely to have the highest amount of contamination. This is especially concerning as floodwaters may contain potentially dangerous chemicals and pathogens. This is particularly worrisome as canines groom themselves in areas where accumulated contaminants are likely to occur: their paws, the area immediately around the sides of the mouth, along their limbs and the areas of short hair, such as the anogenital area (Beaver, 2009). This behavior provides a pathway for oral exposure (Fitzgerald et al., 2008) and requires further investigation in order to better understand oral ingestion of contaminants and gastrointestinal complications frequently observed in dogs during deployment (Otto, 2002; Murphy, 2003; Gordon 2010; 2015).

The benefits of training associated with decontamination have been studied extensively in hospital settings. Formal education programs significantly improve physician and nurse hand hygiene compliance. However, improvements are usually short-lived; without ongoing reinforcement, compliance tends to return to baseline over a period of time (Lam et al., 2004). Therefore, it seems reasonable to assume that improvements in canine decontamination efforts as a result of formal training are also perishable and will require regular reinforcement.

Another objective of this study was to examine the efficacy of a portable field decontamination kit. Based on the results, the kit provided an effective in-field decontamination option for handlers. This is particularly important when timely intervention is critical and a decontamination corridor is not available.

The observations of cross-contamination noted in this study are particularly worrisome. Researchers noted fluorescent residue transfer to the kennels and handlers. During a disaster, contaminated water, particularly in the setting of poor or compromised sanitation, has the

potential to harbor pathogens that could be detrimental to working dog and/or human health. Working dogs exposed to contaminated water can become infected with certain pathogens (e.g., *Leptospira*, *Giardia*), develop illness, and transmit disease to humans. They could also serve as unwitting carriers of enteric pathogens on their fur coat that pose a threat primarily to humans (e.g., *Escherichia coli*, *Vibrio*, enteroviruses) during large-scale disasters. Close contact between working dogs and humans combined with poor hand hygiene practices could lead to human oral ingestion of pathogens and subsequent infection.

CONCLUSION

Training was shown to have a positive effect on contaminant reduction in this study. Further study is recommended to analyze this effect across different breeds with varying coat types. Effects related to coat type should be examined with particular attention given to type of cleanser and repetitive cleansing. In addition, the impact of the cleanser on the skin needs to be evaluated more thoroughly. Currently there is insufficient evidence available to determine which cleansers are safest for canine skin. Improved training and further investigation into the safety and efficacy of existing cleansers are needed to protect the health of our canine partners in the field. Future work should also assess likely areas of contamination on the canine so as to maximize decontamination efforts and improve results.

ACKNOWLEDGEMENTS

The authors would like to give a special thanks to Sonja Heritage and the National Disaster Search Dog Foundation for their participation of this work and for the use of their canines and facility. The authors also thank Wayne Rasband for his assistance in utilizing the ImageJ program for fluorescence analysis and Dakota Discepolo for traveling and assisting in gathering the data for this study.

FINANCIAL DISCLOSURES

SYL was supported by the KM1 Comparative Effectiveness Research Career Development Award (grant number KM1CA156708-01); the Clinical and Translational Science Award (CTSA) program (grant number UL1RR024992) of the National Center for Advancing Translational Sciences (NCATS); and the Barnes-Jewish Patient Safety & Quality Career Development Program, which is funded by the Foundation for Barnes-Jewish Hospital.

LITERATURE CITED

- Ananthapadmanabhan KP, Moore DJ, Subramanyan K, Misra M, Meyer F. 2004 Cleansing without compromise: the impact of cleansers of the skin barrier and the technology of mild cleansing. *Dermatol. Ther* 17: 16–25. [PubMed: 14728695]
- Beaver BV 2009 Canine grooming behavior In: *Canine behavior insights and answers*. Saunders Elsevier, St. Louis, MO p. 285–293.
- Beranda L, González R, Torres-Alvarez B, Alvarez C, Ramírez V, 2002 Correlation between pH and irritant effect of cleansers marketed for dry skin. *Int. J. Dermatol* 41: 494–499. [PubMed: 12207765]
- Binder S, Sanderson LM, 1987 The role of the epidemiologist in natural disasters. *Ann. Emerg. Med* 16: 1081–1084. [PubMed: 3631672]
- Cibulsky SM, Kirk MA, Ignacio JS, Leary AD, Schwartz MD, 2014 Patient decontamination in a mass chemical exposure incident: national planning guidance for communities US Department of Homeland Security Washington, DC.
- Chan HP, Zhai H, Hui X, Maibach HI, 2012 Skin decontamination: principles and perspectives. *Toxicol. Ind. Health* 29(10): 955–968. [PubMed: 22851522]

- Davis RW, Williams TM, Thomas JA, A Kastelein R, Cornell LJ, . The effects of oilContamination and cleaning on sea otters (*Enhydra lutris*). II. Metabolism, thermoregulation and behavior. *Can J. Zool* 66: 2782–2790.
- Euripidou E, Murray V, 2004 Public health impacts of floods and chemical contamination. *J. Public Health* 26:376–383. doi:10.1093/PUBMED/FDH163.
- FEMA. 2012 Handler and canine deployment survey: Hurricane Sandy. Available at www.fema.gov. Accessed February 28, 2018.
- FEMA. 2017 Hurricane deployment triplet - canine health data report. Available at www.fema.gov. Accessed February 28, 2018.
- Fitzgerald SD, Rumbelha WK, Braselton WE, Downend AF, Otto CM, 2008 Pathology and toxicology findings for search-and-rescue dogs deployed to the September 11, 2001, terrorist attack sites: initial five-year surveillance. *J Vet Diagn Invest.* 20: 477–484. [PubMed: 18599853]
- Gordon LE, 2010 Injuries and illnesses among urban search-and-rescue dogs deployed to Haiti following the January 12, 2010, earthquake. *JAVMA.* 240: 396–403.
- Gordon LE, 2015 Injuries and illnesses among Federal Emergency Management Agency-certified search-and-recovery and search-and-rescue dogs deployed to Oso, Washington, following the March 22, 2014, State Route 530 landslide. *JAVMA.* 247: 901–908. [PubMed: 26421402]
- Irving BA, Y Weltman J, Brock DW, Davis CK, Gaesser GA, Weltman A, 2007 NIH imagej and slice-o-matic computed tomography imaging software to quantify soft tissue. *Obesity* 15(2): 370–376. doi:10.1038/oby.2007.573 [PubMed: 17299110]
- Jensen EC 2013 Quantitative analysis of histological staining and fluorescence using Image. *J. Anat. Rec.* 296: 378–381. doi:10.1002/ar.22641
- Jones KE, Dashfield K, Downend AB, Otto CM, . 2004 Search-and-rescue dogs: an overview for veterinarians. *JAVMA.* 225: 854–860. [PubMed: 15485043]
- Lam BCC, Lee J, . Lau YL, . 2004 Hand hygiene practices in a neonatal intensive care unit: a multimodal intervention and impact on nosocomial infection. *Pediatrics* 114 (5): 565–571.
- Lee RHL, Lee PH, 2014 To evaluate the effects of simplified hand washing improvement program in schoolchildren with mild intellectual disability: a pilot study. *Res. Dev. Disabil* 35: 3017–3018.
- Lillquise DR, McCabe ML, Church KH, . 2005 A comparison of traditional handwashing training with active handwashing training in the food handler industry. *J. Environ. Health* 67 (6): 13–16. [PubMed: 15690900]
- Ling ML, How KN, . 2012 Impact of a hospital-wide hand hygiene promotion strategy on healthcare-associated infections. *Antimicrob Resist Infect Control.* 1:13. [PubMed: 22958911]
- Mabry CJ 2015 Comparison of dermatological complications associated with repeated use of a high anionic vs. a low anionic soap to decontaminate canines. Master's thesis, Texas A&M University. Available electronically from <http://hdl.handle.net/1969.1/154965>.
- Meyer W, Neurand K, 1991 Comparison of skin pH in domesticated and laboratory animals. *Arch. Dermatol. Res.* n 283: 16–18.
- Murphy LA, Gwaltney-Brant SM, Albretsen JC, Wismer TA, 2003 Toxicologic agents of concern for search-and-rescue dogs responding to urban disasters. *JAVMA.* 222: 296–302. [PubMed: 12564590]
- Otto CM, Franz MA, Kellogg B, Lewis R, Murphy L, Lauber G, 2002 Field treatment of search dogs: lessons learned from the World Trade Center. *J Vet Emerg Crit Care.* 12: 33–42.
- Soric S, Belanger MP, Wittnich C, 2008 A method for decontamination of animals involved in floodwater disasters. *JAVMA.* 232: 364–370. [PubMed: 18241100]
- Venable E, Discepolo D, Powell E, Liang SY, 2017 An evaluation of current working canine decontamination procedures and methods for improvement. *J. Vet. Behav.: Clin. Appl. Res* 21: 53–58.
- Washington Department of Fish and Wildlife. 2009 Washington oiled sea otter response handbook. p. 9–17. Available at: <http://wdfw.wa.gov/publications/00302/wdfw00302.pdf> Accessed September 12, 2016.
- Welte S, Frink L., 1991 Rescue and rehabilitation of oiled birds. *Waterfowl Management Handbook.* Wilmington, DE p. 1–8.

Young S, Balluz L, aMalilay J. 2004 Natural and technologic hazardous material releases during and after natural disasters: a review. *Sci. Total Environ* 322: 3–20. [PubMed: 15081734]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

HIGHLIGHTS

- 1.** Training on decontamination of working canines improves the efficacy of contaminant reduction
- 2.** Even when decontaminated, dogs can serve as vectors for cross-contamination
- 3.** Disposable field kits can serve to decontaminate dogs when full-scale facilities are unavailable

¹Zones: Hot = contaminant present; Warm = decontamination procedure; Cold = no contaminant present and medical checks performed.

Figure 1:
Illustration of typical decontamination corridor setup in the triage zone¹.

Figure 2A.
Disposable field kit designed to decontaminate canines in the field.

Figure 2B.

Illustrated guide with instructions on proper decontamination procedures.

Figure 3.

Instructions for use of the field decontamination kit, based on researcher's recommendations.

Figure 4.
Anatomic sites for contaminant application: (1) cranial neck, (2) between the shoulder blades, (3) left medial hindlimb and (4) left hind paw.

Figure 5.

A & B. Images representing pre-and post-washing levels of fluorescent residue. Image depicts a score of 3 (> 75% reduction in contamination).

¹Training: Trained handlers = 30-minute formal presentation on field kit and use. Untrained handlers = no formal presentation on field kit and use.

²Frequency of score was utilized to determine percent occurrence.

³Unsuccessful = Contaminant reduction score of 0-1.

⁴Successful = Contaminant reduction score of 2-3.

Figure 6:

Effect of training¹ on frequency of contaminant reduction score² visually assigned across all anatomic sites ($P = 0.0093$).

¹Training: Trained handlers = 30-minute formal presentation on field kit and use. Untrained handlers = no formal presentation on field kit and use.

Figure 7:

Effect of training¹ on contaminant reduction measured across all anatomic sites (P=0.0042).

Figure 8:

Effect of anatomical site on measured contaminant reduction following field decontamination ($P = 0.0055$).

Figure 9:
Evidence of cross-contamination of an untrained handler's canine's kennel, following field decontamination.