Published in final edited form as:

Tetrahedron Lett. 1997 October 13; 38(41): 7171–7174. doi:10.1016/S0040-4039(97)01765-6.

ESTER DERIVED TITANIUM ENOLATE ALDOL REACTION: HIGHLY DIASTEREOSELECTIVE SYNTHESIS OF SYN- AND ANTI-ALDOLS

Arun K. Ghosh, Steve Fidanze, Masanobu Onishi, and Khaja Azhar Hussain

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607.

Abstract

Aldol reactions of bidentate aldehydes and *cis*-1-arylsulfonamido-2-indanyl ester derived titanium enolates proceed with excellent syn-diastereoselectivities and good to excellent isolated yields.

Asymmetric aldol reactions are often utilized in the synthesis of complex organic molecules of biological importance.¹ Over the years, numerous studies led to the development of a number of effective methodologies for syn² and anti-aldol³ reactions. Recently, we reported⁴ *cis*-1-arylsulfonamido-2-indanyl ester derived titanium enolate anti-aldol reactions with high diastereofacial selectivity. In our continuing effort to understand the origin of anti selectivity, we have now established that the choice of *p*-toluenesulfonamido group, the presence of indanyl ring and the choice of metal all are critical to observed anti-aldol diastereoselectivity. Furthermore, based on the possible transition state assembly, we speculated that the incorporation of a chelating substituent on the aldehyde side chain would alter the stereochemical outcome from an anti-aldol to a syn-aldol product. Herein, we report that indeed, the reactions of a number of ester derived titanium enolates with three representative bidentate oxyaldehydes proceeded with excellent syn diastereoselectivity (up to 99% de) with good to excellent isolated yields. The current methodology is convenient and has practical synthetic potential since either the syn or anti-aldol product can be prepared from the same chiral template in a stereopredictable fashion utilizing inexpensive and versatile titanium reagents.⁵

Enantiomerically pure 1*S*, 2*R*-sulfonamide **1**⁴ was converted to propionate ester **2a** with propionyl chloride and pyridine in CH₂Cl₂ at 0°C for 1 h (84% yield). The acylation of **1** with hydrocinnamic acid and 4-methylvaleric acid with DCC and DMAP afforded the respective esters **2b** and **2c** in 85% and 74% yield. Various titanium enolates of **2a-c** were generated by reaction of the respective ester with TiCl₄ (1.2 equiv) in CH₂Cl₂ at 0–23°C for 15 min followed by addition of *N*-ethyl-diisopropylamine (4 equiv) at 23°C and stirring of the resulting brown solution for 2 h. The titanium enolate was then added to the representative aldehyde (2 equiv) precomplexed with TiCl₄ (2.2 equiv) at –78°C and the mixture was stirred at –78°C for 2 h before quenching with aqueous NH₄Cl. As shown in Table 1, aldol reactions of **2a-c** derived titanium enolates with various homologous oxyaldehydes proceeded with excellent syn-diastereoselectivity whereas reactions with a representative aliphatic aldehyde such as isovaleraldehyde, proceeded with excellent anti-

diastereoselectivity. The syn-anti mixture ratio was determined by ^1H NMR (400 MHz) as well as by reverse phase HPLC before and after chromatography. Aldol reactions of benzyloxyacetaldehyde⁶ with propionate derivative **2a**, hydrocinnamate derivative **2b** and 4-methylvalerate derivative **2c** have afforded virtually a single syn-isomer in excellent isolated yield. Similarly, reactions of these esters with benzyloxypropionaldehyde provided excellent syn-diastereoselectivity, but with relatively lower isolated yields. The aldol reaction of **2b** with benzyloxybutyraldehyde⁶ also afforded a syn-aldol product with slightly lower selectivity compared to other oxyaldehydes. In contrast, reactions of isovaleraldehyde with **2b** and **2c** have resulted in anti-aldol reaction with excellent anti-diastereoselectivity and isolated yields.

The assignment of relative and absolute stereochemistry of various syn aldolates **3** (entry 1 to 7) was firmly established after removal of the chiral template. The comparison of ^1H -NMR and ^{13}C -NMR spectra as well as the optical rotation of the resulting acids **5** were compared against the authentic optically pure samples prepared utilizing the boron enolate^{2a} aldol reaction.⁷ The removal of the chiral template was effected by exposure to lithium hydroperoxide in THF at 23°C for 2–3 h affording the corresponding β -hydroxy acid in 84–95% yield. The chiral template **1** was fully recovered without loss of optical purity ($\alpha_{\text{D}}^{23} +34.2$, c 1.8, CHCl_3). Alternatively, the hydrolysis of aldol adduct can be carried out with aqueous lithium hydroxide in THF at 23°C. However, a much longer reaction time is required for complete hydrolysis (for **3**, 18–24 h).

To rationalize the anti-aldol stereoselectivities with monodentate aldehydes, we previously⁴ postulated a Zimmerman-Traxler⁸ type transition state model **A** in which the metalocycle is assumed to adopt a chair-like conformation with a possible π -stacking interaction between the aromatic rings. This model is the basis of our further speculation that the addition of a chelating substituent on the aldehyde side chain would result in a transition state assembly such as **B**.⁹ The observed syn-stereoselectivity of bidentate aldehydes is consistent with this postulated model. The present syn-stereoselectivity can also be explained by an acyclic transition state similar to that proposed by Gennari *et al.*¹⁰ The enhanced selectivity for benzyloxyacetaldehyde and benzyloxy-propionaldehyde compared to benzyloxybutyraldehyde is most likely due to effective metal chelation through five and six membered transition states rather than a less favorable seven membered ring system. The involvement of crucial metal chelation was further evidenced by the fact that the reaction of enolate of **2a** with *tert*-butyldiphenylsilyloxyacetaldehyde afforded a 70:30 mixture of syn/anti aldol products due to steric bulk of the surrounding ether oxygen which hinders effective chelation. The possible aromatic π -stacking interaction in the transition state model is supported by the fact that incorporation of methylsulfonamide in place of tosylsulfonamide in **2a** resulted in a mixture (70:30) of anti/syn diastereomers with isovaleraldehyde (through **B**). Furthermore, removal of the indane aromatic ring in **2a** also resulted in a mixture (50:50) of syn/anti diastereomers.¹¹ Further evidence for such a π —stacking interaction is the subject of ongoing investigation.

In summary, an ester derived titanium enolate based highly selective (96–98% de) syn-aldol process has been developed. The reaction of the same titanium enolate with monodentate aldehyde however provides anti-aldol product with excellent diastereoselectivity. Thus, with

proper choice of chiral template and aldehyde one can prepare either syn- or anti-aldol product in a stereopredictable fashion. Synthetic applications and mechanistic investigations are currently in progress.

Acknowledgment:

Financial support of this work by the National Institute of Health (GM 53386) is gratefully acknowledged. The authors thank Professors Thomas Hoye and George Gould for helpful discussions. M. O. is a visiting scientist from Nihon Nohyaku Co. Ltd., Japan.

References and Notes:

- (a). Kim B-M; Williams SF; Masamune S *Comprehensive Organic Synthesis*; Trost BM; Fleming I; Eds.; Pergamon Press; Oxford, 1991; Vol 2 (Heathcock CH, Ed.), Chapter 1.7, 239;(b)Franklin AS; Paterson I; *Contemp. Org. Synth* 1994, 317 and references cited therein.
- (a). Evans DAAldrich. *Acta* 1982 15, 23;(b)Roder H; Helmchen G; Peters E-M; von Schmering H-G *Angew. Chem. Int. Ed. Engl* 1984, 23, 898;(c)Masamune S; Choy W; Peterson JS; Sita LR *ibid.* 1985, 24, 1;(d)Paterson I; Lister MA; McClure CK *Tetrahedron Lett* 1986, 27, 4787;(e)Corey EJ; Imwinkelreid R; Pikul S; Xiang YB J. *Am. Chem. Soc* 1989, 111, 5493;(f)Oppolzer W; Blagg J; Rodriguez I; Walther EJ *ibid* 1990, 112, 2767;(f)Ghosh AK; Duong TT; McKee SP J. *Chem. Soc., Chem. Commun* 1992, 1673; [PubMed: 30364406] (g)Yan T-H; Tan C-W; Lee H-C; Lo H-C J. *Am. Chem. Soc* 1993, 115, 2613;(h)Abiko A; Liu J-F; Masamune SJ. *Org. Chem* 1996, 61, 2590 and references cited therein. [PubMed: 11667081]
- (a). Meyers AI; Yamamoto Y *Tetrahedron* 1984, 40, 2309;(b)Helmchen G; Leikauf U; Taufer-Knopfel I *Angew. Chem., Int. Ed. Eng* 1985, 24, 874;(c)Gennari C; Bernardi A; Colombo L; Scolastico CJ. *Am. Chem. Soc* 1985, 107, 5812;(d)Oppolzer W; Marco-Contelles J *Helv. Chim. Acta* 1986, 69, 1699;(e)Masamune S; Sato T; Kim BM; Wollman TA J. *Am. Chem. Soc* 1986, 108, 8279;(f)Danda H; Hansen MM; Heathcock CH J. *Org. Chem* 1990, 55, 173;(g)Corey EJ; Kim SS J. *Am. Chem. Soc* 1990, 112, 4976;(h)Corey EJ; Kim SS *Tetrahedron Lett* 1990, 31, 3715;(i)Meyers AG; Widdowson KL J. *Am. Chem. Soc* 1990, 112, 9672;(j)Duthaler RO; Herold P; Helfer S-W; Riediker M *Helv. Chim. Acta* 1990, 73, 659;(k)Walker MA; Heathcock CH J. *Org. Chem* 1991, 56, 5747;(l)Gennari C; Moresca D; Vieth S; Vulpetti A *Angew. Chem., Int. Ed. Eng* 1993, 32, 1618;(m)Paterson I; Wren SPJ. *C. S. Chem. Comm* 1993, 1790;(n)Abiko A; Liu J-F; Masamune SJ *Am. Chem. Soc* 1997, 119, 2586 and references cited therein.
- Ghosh AK; Onishi MJ *Am. Chem. Soc* 1996, 118, 2527.
- Reetz MT *Stereoselectivities in Lewis Acid Promoted Reactions*, Schinzer D, Ed., Kluwer Academic Publishers, 1989, Chapter 6, 107.
- Hon Y-S; Lin S-W; Lu L; Chen Y-J *Tetrahedron* 1995, 51, 5019.
- Observed optical rotation of the corresponding β -hydroxy acid in CHCl_3 : 5a, $[\alpha]_{\text{D}}^{23} +12.97$ (c, 3.7); 5b, $[\alpha]_{\text{D}}^{23} +9.44$ (c, 1.8); 5c, $[\alpha]_{\text{D}}^{23} -9.09$ (c, 1.1); 5d, $[\alpha]_{\text{D}}^{23} -8.98$ (c, 3.2); 5e, $[\alpha]_{\text{D}}^{23} +11.9$ (c, 1.1); 5f, $[\alpha]_{\text{D}}^{23} -4.14$ (c, 4.3); 5g, $[\alpha]_{\text{D}}^{23} -3.33$ (c, 0.9); 6i, $[\alpha]_{\text{D}}^{23} -18.2$ (c, 2.1); 6j, $[\alpha]_{\text{D}}^{23} -26.7$ (c, 1.6); 6k, $[\alpha]_{\text{D}}^{23} -28.4$ (c, 2.4).
- Zimmerman HE; Traxler MD J. *Am. Chem. Soc* 1957, 79, 1920.
- In this model, Ti presumably assumes a six-coordinate ion-pair with chloride. For related example, see, Gennari C; Bernardi A; Scolastico C; Potenza D *Tetrahedron Lett* 1985, 26, 4129.
- Mukaiyama aldol reaction of N-methylephedrine derived silyl ketene acetal and benzyloxypropionaldehyde has also provided a syn-selective aldol reaction (selectivity 78:11:11), see Gennari C; Cozzi GJ *Org. Chem* 1988, 53, 4015.
- For the preparation of the corresponding cis-2-amino-cyclopentan-1-ol, see, Ghosh AK; Cho H; Onishi M *Tetrahedron:Asymm.* 1997, 8, 821.

Figure 1.

Scheme I:

(a) CH₃CH₂COCl, pyridine, CH₂Cl₂, 0°C, 1 h for **2a**; RCO₂H, DCC, DMAP, CH₂Cl₂, 23°C, 18–24 h for **2b-c**; (b) TiCl₄, *i*Pr₂NEt, 23°C then R₁CHO and TiCl₄, CH₂Cl₂, –78°C, 2 h; (c) LiOH, 30% H₂O₂, THF-H₂O, 23°C, 2–3 h.

Table 1.Aldol reaction of various esters **2a-c** with representative aldehydes

Entry	Ester	Aldehyde	Compd ^a	% Yield ^b	Syn:Anti (3/4) ^c
1.	2a	BnOCH ₂ CHO	3a	84	98 : 2
2.	2a	BnO(CH ₂) ₂ CHO	3b	51	98 : 2
3.	2b	BnOCH ₂ CHO	3c	84	99 : 1
4.	2b	BnO(CH ₂) ₂ CHO	3d	51	99 : 1
5.	2b	BnO(CH ₂) ₃ CHO	3e	55	94 : 6
6.	2c	BnOCH ₂ CHO	3f	83	99 : 1
7.	2c	BnO(CH ₂) ₂ CHO	3g	56	99 : 1
8.	2a	<i>i</i> BuCHO	4i	92	1 : 99
9.	2b	<i>i</i> BuCHO	4j	91	1 : 99
10.	2c	<i>i</i> BuCHO	4k	83	1 : 99

^a Only isolated product; **3a** (R=Me, R₁=CH₂OBn); **3b** (R=Me, R₁=(CH₂)₂OBn); **3c** (R=Bn, R₁=CH₂OBn); **3d** (R=Bn, R₁=(CH₂)₂OBn); **3e** (R=Bn, R₁=(CH₂)₃OBn); **3f** (R=*i*Bu, R₁=CH₂OBn); **3g** (R=*i*Bu, R₁=(CH₂)₂OBn); **4i** (R=Me, R₁=*i*Bu); **4j** (R=Bn, R₁=*i*Bu); **4k** (R=*i*Bu, R₁=*i*BU);

^b Isolated yield after chromatography.

^c Ratios determined by ¹H-NMR and HPLC analysis before and after chromatography. Reaction time = 1.5–2 h.