has already started to address this issue, taking the steps to better serve the Hispanic population by not only making available educational materials in Spanish but also, and as important, exploring the needs of this community through a recent national survey. We look forward to their findings and plans to improve the health of all patients with CF. These efforts in aggregate to scientific investigations will permit the clearer identification of factors amenable to intervention and achieve a goal of eliminating problematic differentials in outcomes. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

MyMy C. Buu, M.D.
Carlos E. Milla, M.D.
Center for Excellence in Pulmonary Biology
Stanford University School of Medicine
Stanford, California

References

1. Cystic Fibrosis Foundation Patient Registry. 2016 Annual Data Report. Bethesda, MD: Cystic Fibrosis Foundation; 2017.
2. The Canadian Cystic Fibrosis Registry 2013 Annual Report. Cystic Fibrosis Canada; 2015 Jan [accessed 2018 Sept 7]. Available from: <https://www.cysticfibrosis.ca/uploads/cf%20care/Canadian-CF-Registry-2013-FINAL.pdf>.
3. Barr HL, Britton J, Smyth AR, Fogarty AW. Association between socioeconomic status, sex, and age at death from cystic fibrosis in England and Wales (1959 to 2008): cross sectional study. *BMJ* 2011; 343:d4662.
4. Colby SL, Ortman JM. Projections of the size and composition of the US population: 2014 to 2060: current population reports, P25-1143. Washington, DC: US Census Bureau; 2014.
5. Kharrazi M, Yang J, Bishop T, Lessing S, Young S, Graham S, *et al.*; California Cystic Fibrosis Newborn Screening Consortium. Newborn screening for cystic fibrosis in California. *Pediatrics* 2015;136: 1062–1072.
6. World Health Organization. The molecular genetic epidemiology of cystic fibrosis: report of a Joint Meeting of WHO/ECFTN/ICF(M)/AECFS. Geneva: World Health Organization; 2004.
7. US Census Bureau. 2010 Census of Population and Housing, Summary Population and Housing Characteristics, CPH-1-1, United States. Washington, DC: US Government Printing Office; 2013.
8. Buu M, Milla C, Chan J, Wise P, Cornfield D. Hospitalization and death rates of Hispanic cystic fibrosis pediatric patients in California. *Pediatr Pulmonol* 2010;S33:390–391.
9. Buu MC, Sanders LM, Mayo JA, Milla CE, Wise PH. Assessing differences in mortality rates and risk factors between Hispanic and non-Hispanic patients with cystic fibrosis in California. *Chest* 2016; 149:380–389.
10. Rho J, Ahn C, Gao A, Keller A, Jain R. Evaluating outcomes disparities in the Hispanic cystic fibrosis population: a need for a national analysis. *Chest* 2016;150:751–753.
11. Rho J, Ahn C, Gao A, Sawicki GS, Keller A, Jain R. Disparities in mortality of Hispanic patients with cystic fibrosis in the United States: a national and regional cohort study. *Am J Respir Crit Care Med* 2018;198:1055–1063.
12. McKone EF, Emerson SS, Edwards KL, Aitken ML. Effect of genotype on phenotype and mortality in cystic fibrosis: a retrospective cohort study. *Lancet* 2003;361:1671–1676.
13. Drumm ML, Konstan MW, Schluchter MD, Handler A, Pace R, Zou F, *et al.*; Gene Modifier Study Group. Genetic modifiers of lung disease in cystic fibrosis. *N Engl J Med* 2005;353:1443–1453.
14. Dorfman R, Sandford A, Taylor C, Huang B, Frangolias D, Wang Y, *et al.* Complex two-gene modulation of lung disease severity in children with cystic fibrosis. *J Clin Invest* 2008;118:1040–1049.
15. Buu MC, Sanders LM, Mayo JA, Milla CE, Wise PH. Assessing differences in mortality rates and risk factors between Hispanic and non-Hispanic patients with cystic fibrosis in California. *Chest* 2016; 149:380–389.
16. Pino-Yanes M, Thakur N, Gignoux CR, Galanter JM, Roth LA, Eng C, *et al.* Genetic ancestry influences asthma susceptibility and lung function among Latinos. *J Allergy Clin Immunol* 2015;135: 228–235.
17. Seldin MF, Pasaniuc B, Price AL. New approaches to disease mapping in admixed populations. *Nat Rev Genet* 2011;12:523–528.
18. Sandefur GD, Campbell ME, Eggerling-Boeck J. Racial and ethnic identification, official classifications, and health disparities. In: Anderson NB, Bulatao RA, Cohen B, editors; National Research Council (US) Panel on Race, Ethnicity, and Health in Later Life. Critical perspectives on racial and ethnic differences in health in late life. Washington, DC: National Academies Press; 2004.

Copyright © 2018 by the American Thoracic Society

Respiratory Syncytial Virus–induced Acute Disease Severity and Long-Term Wheezing Uncovering the Unexpected

Despite over 60 years of research efforts, there are fundamental aspects of the pathogenesis of respiratory syncytial virus (RSV) infection, especially in children, that remain to be elucidated. For years, most research on RSV pathogenesis was performed on animal models, more specifically mice (1, 2). Those studies provided a detailed understanding of the pathogenesis of RSV infection, and began to uncover how the virus and its proteins

interacting with the different components of the immune system influence both the clinical manifestations of the acute disease and long-term airway hyperreactivity (3–5), a phenotype similar to the persistent wheezing observed after RSV infection in young children (6). As more advanced technologies made it possible to study RSV immunopathogenesis in infants and young children, it became evident that one could not directly translate the observations derived from mouse models to the immune responses to RSV in young children.

Understanding how the interactions between virus replication and immune response determine disease severity has attracted major attention, especially in the current era of active development

O.R. and A.M. are supported in part by NIH grants AI112524 and AI131386. Originally Published in Press as DOI: 10.1164/rccm.201805-0908ED on May 30, 2018

of new interventions against RSV, including antiviral agents, monoclonal antibodies, and vaccines (7). For years, the leading hypothesis was that an exaggerated inflammatory response in children hospitalized with severe RSV infection was the major determinant of disease severity.

More recently, however, research by different investigators began to unravel a more complex situation. A number of studies in children showed an association between RSV viral loads and disease severity (although not consistently), as demonstrated by significant correlations between initial viral loads and duration of hospitalization or the need for intensive care (8, 9). At the same time, several observations began to support an alternative view, as studies showed that children with severe RSV disease displayed what appeared to be an inadequate immune response. Investigators found that children with bronchiolitis who required hospitalization had reduced IFN- γ , IL-4, IL-17, and IP-10 concentrations in nasal wash samples compared with those with milder disease who were discharged home and managed as outpatients (10). Likewise, blood transcriptome analyses showed that among young children hospitalized with RSV infection, those with more severe disease had suboptimal expression of IFN-related genes but overexpression of neutrophil- and inflammation-related genes (11). In addition, children who died of severe RSV infection had a near absence of CD8 T cell and natural killer cell infiltrates in the lungs (12). In a recent study, Piedra and colleagues began to integrate these apparently discrepant observations and found that RSV gene copy numbers were higher in children with less severe RSV disease, and significantly correlated with the concentrations of the aforementioned cytokines and chemokines (13). They proposed a distinct model in which higher RSV loads early during the infection would induce a robust innate immune response, which would protect against disease progression.

In this context, the study by Thwaites and colleagues (pp. 1074–1084) in this issue of the *Journal* provides additional and relevant insights into this new paradigm (14). The investigators longitudinally analyzed viral loads and immune responses in the upper respiratory tract of infants with and without additional comorbidities who were hospitalized with acute bronchiolitis caused by RSV and other viruses. They took advantage of a new technique (nasosorption) that, with some limitations, allows noninvasive and repeated sampling of the upper respiratory tract in young infants. Using this methodology, they measured viral loads, cytokine and chemokine concentrations, and gene expression patterns sequentially. Their results indicate that infants with severe disease display lower RSV viral loads as well as lower concentrations of IFN- γ and CCL5/RANTES compared with infants with moderate disease. A gene-expression analysis in 43 infants with bronchiolitis (20 of whom had RSV) also showed decreased gene expression of type I IFN and increased expression of MUC5AC and IL-17A in severe cases. The longitudinal sampling and innovative time-weighted analyses of the virologic and immunologic variables provided a better understanding of the pathogenesis of RSV infection in young children, and how the interaction between viral replication and the immune response contributes to disease severity. These findings confirm and extend previous studies based on one-time sample analyses, and add to the increasing evidence suggesting that severe RSV disease is associated with an inadequate immune response and lower viral loads.

In the same issue of the *Journal*, a study conducted by Turi and colleagues (pp. 1064–1073) provides novel information regarding the nasal immune response pathways that are activated or suppressed upon RSV or rhinovirus infection and their association with subsequent development of childhood wheeze (15). The investigators leveraged the analyses of respiratory samples obtained as part of a birth cohort study (INSPIRE [Infant Susceptibility to Pulmonary Infections and Asthma following RSV Exposure]) in which 193 healthy, term infants were followed during the first 2 years of life to capture their first RSV (or rhinovirus) infection and its association with recurrent wheezing. By applying a 53 multianalyte panel in nasal samples, the authors identified two distinct immune-response phenotypes, or clusters, present at the time of a first RSV or rhinovirus infection that were associated with subsequent wheezing. They validated their findings by using a number of elegant analytical strategies, such as bootstrap analyses, to compensate for the small number of infants in the RSV and rhinovirus clusters. The authors found that cluster 2, which was dominated by type 2 and type 17 cytokines, but also IFN- α and IFN- γ , was independently associated with an increased risk of wheezing during the first 2 years of life in infants with RSV infection.

These studies demonstrate the need to improve our understanding of the complex pathogenesis of acute RSV infection and its association with long-term sequelae without preconceived assumptions. As illustrated by the works of Thwaites and colleagues (14) and Turi and colleagues (15), the implementation of high-resolution assays and sophisticated statistical tools that can be applied in the clinical context has accelerated our ability to conduct in-depth studies of RSV pathogenesis in young children with limited sample sizes. Although their direct clinical applicability is still limited, these studies represent an important foundation for future work. It is now time to design larger, prospective cohort studies to validate these findings and link these observations by carefully examining whether (and if so, how) the virologic and immunologic factors that determine acute disease severity influence the development of post-RSV wheezing. The clinical development of new prophylactic and therapeutic interventions for RSV infection will provide a unique opportunity to incorporate detailed virologic and immunologic analyses to determine how modifications of the host/virus variables associated with an intervention correlate with both acute disease severity and post-RSV wheezing. Such an approach should help us achieve the ultimate goal of diminishing the significant acute and long-term morbidity, as well as mortality, that is associated with RSV infection in young children. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Octavio Ramilo, M.D.
 Asuncion Mejias, M.D., Ph.D.
Division of Pediatric Infectious Diseases
 and
Center for Vaccines and Immunity
Nationwide Children's Hospital
Columbus, Ohio
 and
The Ohio State University
Columbus, Ohio

ORCID IDs: 0000-0002-0565-5239 (O.R.); 0000-0002-5983-8006 (A.M.).

References

1. Graham BS, Perkins MD, Wright PF, Karzon DT. Primary respiratory syncytial virus infection in mice. *J Med Virol* 1988;26:153–162.
2. Openshaw PJ, Pemberton RM, Ball LA, Wertz GW, Askonas BA. Helper T cell recognition of respiratory syncytial virus in mice. *J Gen Virol* 1988;69:305–312.
3. Estripeaut D, Torres JP, Somers CS, Tagliabue C, Khokhar S, Bhoj VG, et al. Respiratory syncytial virus persistence in the lungs correlates with airway hyperreactivity in the mouse model. *J Infect Dis* 2008;198:1435–1443.
4. Jafri HS, Chavez-Bueno S, Mejias A, Gomez AM, Rios AM, Nassi SS, et al. Respiratory syncytial virus induces pneumonia, cytokine response, airway obstruction, and chronic inflammatory infiltrates associated with long-term airway hyperresponsiveness in mice. *J Infect Dis* 2004;189:1856–1865.
5. Mejias A, Chávez-Bueno S, Rios AM, Saavedra-Lozano J, Fonseca Aten M, Hatfield J, et al. Anti-respiratory syncytial virus (RSV) neutralizing antibody decreases lung inflammation, airway obstruction, and airway hyperresponsiveness in a murine RSV model. *Antimicrob Agents Chemother* 2004;48:1811–1822.
6. Stein RT, Sherrill D, Morgan WJ, Holberg CJ, Halonen M, Taussig LM, et al. Respiratory syncytial virus in early life and risk of wheeze and allergy by age 13 years. *Lancet* 1999;354:541–545.
7. Mazur NI, Martínón-Torres F, Baraldi E, Fauroux B, Greenough A, Heikkinen T, et al. Respiratory Syncytial Virus Network (ReSViNET). Lower respiratory tract infection caused by respiratory syncytial virus: current management and new therapeutics. *Lancet Respir Med* 2015;3:888–900.
8. Hasegawa K, Jartti T, Mansbach JM, Laham FR, Jewell AM, Espinola JA, et al. Respiratory syncytial virus genomic load and disease severity among children hospitalized with bronchiolitis: multicenter cohort studies in the United States and Finland. *J Infect Dis* 2015;211:1550–1559.
9. El Saleeby CM, Bush AJ, Harrison LM, Aitken JA, Devincenzo JP. Respiratory syncytial virus load, viral dynamics, and disease severity in previously healthy naturally infected children. *J Infect Dis* 2011;204:996–1002.
10. Nicholson EG, Schlegel C, Garofalo RP, Mehta R, Scheffler M, Mei M, et al. Robust cytokine and chemokine response in nasopharyngeal secretions: association with decreased severity in children with physician diagnosed bronchiolitis. *J Infect Dis* 2016;214:649–655.
11. Mejias A, Dimo B, Suarez NM, Garcia C, Suarez-Arrabal MC, Jartti T, et al. Whole blood gene expression profiles to assess pathogenesis and disease severity in infants with respiratory syncytial virus infection. *PLoS Med* 2013;10:e1001549.
12. Welliver TP, Garofalo RP, Hosakote Y, Hintz KH, Avendano L, Sanchez K, et al. Severe human lower respiratory tract illness caused by respiratory syncytial virus and influenza virus is characterized by the absence of pulmonary cytotoxic lymphocyte responses. *J Infect Dis* 2007;195:1126–1136.
13. Piedra FA, Mei M, Avadhanula V, Mehta R, Aideyan L, Garofalo RP, et al. The interdependencies of viral load, the innate immune response, and clinical outcome in children presenting to the emergency department with respiratory syncytial virus-associated bronchiolitis. *PLoS One* 2017;12:e0172953.
14. Thwaites RS, Coates M, Ito K, Ghazaly M, Feather C, Abdulla F, et al. Reduced nasal viral load and IFN responses in infants with respiratory syncytial virus bronchiolitis and respiratory failure. *Am J Respir Crit Care Med* 2018;198:1074–1084.
15. Turi KN, Shankar J, Anderson LJ, Rajan D, Gaston K, Gebretsadik T, et al. Infant viral respiratory infection nasal immune-response patterns and their association with subsequent childhood recurrent wheeze. *Am J Respir Crit Care Med* 2018;198:1064–1073.

Copyright © 2018 by the American Thoracic Society

Nothing but a Number? Age and Precision Treatment in Pulmonary Arterial Hypertension

Pulmonary arterial hypertension (PAH) is an incurable disease associated with significant morbidity and mortality. Research during the last several decades has led to the development and approval of numerous PAH-specific therapies. Although more pharmacologic treatment options are available for patients with PAH, with progress comes increased cost in a disease for which substantial treatment burden and healthcare use already exist (1).

For newly diagnosed patients with PAH not at high risk, guidelines recommend initial combination therapy with an oral ERA (endothelin receptor antagonist) and a PDE5i (phosphodiesterase-5 inhibitor) (2). This recommendation was based on data from a large randomized trial that demonstrated improved time to clinical worsening with ERA and PDE5i compared with either therapy alone (3). Despite a variable definition of clinical worsening across trials, a recent meta-analysis similarly showed improved outcomes with combination therapy compared with monotherapy (4). Up-front triple therapy with ERA, PDE5i, and prostacyclin analog therapy has been proposed, and a large, multicenter randomized trial is currently ongoing

to assess its efficacy (NCT02558231). Although a multidrug approach might be effective by combining therapies with synergistic mechanisms, it is also possible that this strategy shows benefit in that it simply ensures that a greater proportion of individuals receive the drug most effective for them.

Despite these advances, the choice of initial PAH therapy remains based on clinical risk assessment and patient and provider preference. It is well established that certain PAH populations have worse prognosis than others; namely, men (5) and those with PAH as a complication of connective tissue disease (6). However, there are few data examining differential responses to therapy, particularly with respect to key demographic characteristics such as age, sex, and race and ethnicity. One study by Gabler and colleagues (7) found that women with PAH had greater increases in 6-minute-walk distance than men when treated with an ERA, and that white individuals may have responded better to ERA treatment than black individuals. In another study, Mathai and colleagues (8) showed that compared with women, men with PAH were more likely to have clinically meaningful responses in functional and health-related quality-of-life measures when treated with tadalafil.

In this issue of the *Journal*, Heresi and colleagues (pp. 1090–1093) retrospectively compared differences in survival in patients with

Originally Published in Press as DOI: 10.1164/rccm.201806-1075ED on July 9, 2018