Images in Cardiovascular Medicine

Pivotal Role of Intraoperative Transesophageal Echocardiography for Detecting Iatrogenic Aortic Regurgitation due to Cardiac Catheterization

Kantha R. Kolla , MBBS¹, Carolyn M. Larsen , MD¹, Roger L. Click, MD, PhD², Hari P. Chaliki , MD¹, and Simon Maltais, MD, PhD³

¹Department of Cardiovascular Diseases, Mayo Clinic, Scottsdale, AZ, USA

²Department of Cardiovascular Medicine, Mayo Clinic, Rochester, MN, USA

³Department of Cardiovascular Surgery, Mayo Clinic, Rochester, MN, USA

OPEN ACCESS

Received: Apr 18, 2018

Revised: Aug 21, 2018

Accepted: Sep 4, 2018

Correspondence to

Hari P. Chaliki, MD

Division of Cardiovascular Diseases, Mayo Clinic, 13400 E Shea Blvd, Scottsdale, AZ 85259, USA.

E-mail: chaliki.hari@mayo.edu

Copyright © 2018. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Kantha R. Kolla

<https://orcid.org/0000-0003-1959-928X>

Carolyn M. Larsen

<https://orcid.org/0000-0001-6530-3545>

Hari P. Chaliki

<https://orcid.org/0000-0002-9470-9605>

Conflict of Interest

The authors have no financial conflicts of interest.

A 67-year-old man with a history of fatigue and dyspnea on exertion was diagnosed with atrial fibrillation and severe mitral regurgitation. Preoperative transthoracic echocardiography (TTE) showed a normal aortic valve without aortic regurgitation (**Figure 1** and **Supplementary Video 1**) and bileaflet mitral valve prolapse with severe mitral regurgitation. As part of the preoperative evaluation, cardiac catheterization was performed, which showed normal coronary arteries. The patient was referred for mitral valve repair.

Intraoperative transesophageal echocardiography (TEE) before sternotomy showed a moderate, unexpected eccentric jet of aortic regurgitation (**Figure 2** and **Supplementary Video 2**). The preoperative left ventriculogram showed the multipurpose guiding catheter intermittently entering the aortic sinus (**Figure 3** and **Supplementary Video 3**). Because aortic regurgitation was not present before catheterization, the catheter likely caused trauma to the aortic leaflet, resulting in aortic regurgitation. Two areas with defects were found during surgery: an apparent cleft in the right coronary cusp and a tear in the central right coronary cusp (**Figure 4A**). Aortic valve repair consisted of commissuroplasty between the right and noncoronary cusps, repair of the tear in the right coronary cusp, and resuspension of all 3 commissures (**Figure 4B**). Mitral valve repair, ligation of the left atrial appendage, and surgical CryoMaze of the left atrium were performed as planned. Follow-up TTE showed insignificant aortic regurgitation, trivial mitral regurgitation, and normal left ventricular ejection fraction.

Figure 1. Transthoracic echocardiogram, showing a normal aortic valve without aortic regurgitation. AO = aorta; LV = left ventricle.

Author Contributions

Conceptualization: Chaliki HP, Maltais S;
Data curation: Kolla KR, Larsen CM, Click RL;
Writing - original draft: Kolla KR, Chaliki HP;
Writing - review & editing: Kolla KR, Click RL,
Chaliki HP, Maltais S.

Figure 2. Intraoperative transesophageal echocardiogram before surgery, showing aortic regurgitation. Ao = aorta; LA = left atrium; LV = left ventricle.

Figure 3. Cineangiogram during contrast injection of the left ventriculogram. (A) Left ventricular catheter (arrow) just below the right coronary cusp. (B) Left ventricular catheter (arrow) just above the right coronary cusp.

Figure 4. Intraoperative image. (A) Tear in the right coronary cusp (arrows). (B) Aortic valve repair (arrows).

Iatrogenic aortic regurgitation due to cardiac catheterization is a rare complication most often associated with aortic dissection.¹⁾ Few cases have been reported of iatrogenic aortic regurgitation due to aortic leaflet trauma during cardiac catheterization.²⁻⁴⁾ This case highlights the importance of intraoperative TEE in identifying unexpected findings such as aortic regurgitation secondary to iatrogenic aortic valve cusp trauma.

SUPPLEMENTARY MATERIALS

Supplementary Video 1

Preoperative transthoracic echocardiogram showing the presence of mitral regurgitation and absence of aortic regurgitation.

[Click here to view](#)

Supplementary Video 2

Intraoperative transesophageal echocardiogram showing an eccentric jet of aortic regurgitation.

[Click here to view](#)

Supplementary Video 3

Left ventricular cineangiogram showing the multipurpose guiding catheter briefly entering the aortic sinus from the left ventricle during the contrast injection.

[Click here to view](#)

REFERENCES

1. Wyss CA, Steffel J, Lüscher TF. Isolated acute iatrogenic aortic dissection during percutaneous coronary intervention without involvement of the coronary arteries. *J Invasive Cardiol* 2008;20:380-2.
[PUBMED](#)
2. Denyer MH, Elliott CM, Robicsek F. Pericardial patch repair of aortic cusp perforation caused by cardiac catheterization. *J Card Surg* 1988;3:155-7.
[PUBMED](#) | [CROSSREF](#)
3. Fundarò P, Di Mattia DG, Cialfi A, Santoli C. Repair of aortic cusp traumatic laceration caused by cardiac catheterization. *J Heart Valve Dis* 1996;5:281-2.
[PUBMED](#)
4. Rolf T, Tozzi P, Roumy A, Iglesias JF, von Segesser LK. Aortic valve lesion after coronary angiography. *Interact Cardiovasc Thorac Surg* 2011;12:205-6.
[PUBMED](#) | [CROSSREF](#)