Published in final edited form as:

Int J Hyg Environ Health. 2019 January ; 222(1): 101–110. doi:10.1016/j.ijheh.2018.08.008.

Dibutyl-Phthalate Exposure from Mesalamine Medications and Serum Thyroid Hormones in Men

Feiby L. Nassan^{a,b}, Tim I.M. Korevaar^a, Brent A. Coull^c, Niels E. Skakkebaek^e, Stephen A. Krawetz^f, Molly Estill^f, Elizabeth J. Hait^g, Joshua R. Korzenik^h, Jennifer B. Ford^a, Ralph A. De Poortereⁱ, Maarten A. Broerenⁱ, Alan C. Moss^j, Thomas R. Zoeller^k, and Russ Hauser^{a,d,l}

^aDepartment of Environmental Health Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA. ^bDepartment of Nutrition Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA. ^cDepartment of Biostatistics Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA. ^dDepartment of Epidemiology Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA. ^eDepartment of Growth and Reproduction and EDMaRC, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark. ^fDepartment of Obstetrics & Gynecology, Center for Molecular Medicine & Genetics, Wayne State University, Detroit, Michigan, USA. ^gDivision of Gastroenterology, Children's Hospital Boston, Harvard Medical School, Boston, Massachusetts, USA. ^hDivision of Gastroenterology, Hepatology and Endoscopy, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA. ⁱLaboratory of Clinical Chemistry and Haematology, Máxima Medical Centre, Veldhoven, De Run, 4600 The Netherlands ^jCenter for Inflammatory Bowel Disease, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA. ^kDepartment of Biology, University of Massachusetts, Amherst, Massachusetts, USA. ^lVincent Obstetrics and Gynecology, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA.

Abstract

Background: Dibutyl phthalate (DBP) is an endocrine disruptor and used in some medication coatings, such as mesalamine for treatment inflammatory bowel disease (IBD).

Objectives: To determine whether high-DBP from some mesalamine medications alters thyroid function.

Methods: Seventy men with IBD, without thyroid disease or any radiation history participated in a crossover-crossback prospective study and provided up to 6 serum samples (2:baseline, 2:crossover, 2:crossback). Men on non-DBP mesalamine (background exposure) at baseline

Address correspondence to: Feiby L. Nassan, Sc.D., Harvard T.H. Chan School of Public Health, Building II 3rd Floor, 655 Huntington Avenue, Boston, MA 02115, Tel: (857) 244-3312, fen769@mail.harvard.edu.

Author contributions

F.L.N performed the statistical analysis and wrote the manuscript. F.L.N, B.A.C and R.H wrote and received a pilot grant for this work from NIEHS center at the Harvard T. H. Chan School of Public Health. T.I.M.K, B.A.C, N.E.S, S.A.K, M.E, E.J.H, J.R.K, J.B.F, R.A.DP, M.A.B, A.C.M., T.R.Z, and R.H reviewed and edited the manuscript; F.L.N and R.H had primary responsibility for final content. All authors read and approved the final manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

crossed-over to DBP-mesalamine (high exposure) then crossed-back to non-DBP mesalamine (B₁HB₂-arm) and vice versa for men on DBP-mesalamine at baseline (H₁BH₂-arm). Serum concentrations of total triiodothyronine (T3), total thyroxine (T4), free triiodothyronine (FT3), free thyroxine (FT4), thyroid-stimulating hormone (TSH) and thyroid peroxidase antibody (TPOAb), and thyroglobulin antibody (TgAb).

Results: After crossover in B₁HB₂-arm (26 men, 134 samples), T3 decreased 10% (95% confidence interval (CI): -14%, -5%), T3/T4 ratio decreased 8% (CI: -12%, -3%), TPOAb, and TgAb concentrations decreased, 11% (-20%, -2%) and 15% (-23%, -5%), respectively; after crossback, they increased. When men in the H₁BH₂-arm (44 men, 193 samples) crossed-over, T3 decreased 7% (CI: -11%, -2%) and T3/T4 ratio decreased 6% (CI: -9%, -2%). After crossback, only TgAb increased and FT4 decreased.

Conclusions: High-DBP novel exposure or removal from chronic high-DBP exposure could alter elements of the thyroid system, and most probably alters the peripheral T4 conversion to T3 and thyroid autoimmunity, consistent with thyroid disruption. After exposure removal, these trends were mostly reversed.

Keywords

Phthalate; endocrine disruptor; thyroid; hormones; men

Introduction

Thyroid hormone regulates metabolism in practically all tissues of the human body. Differences in thyroid function, even within the clinical reference range, have been associated with a higher risk of adverse clinical outcomes including cardiovascular disease, dementia and diabetes¹⁻³. Although recent experimental⁴⁻⁶ studies suggest that phthalate exposure can disrupt the hypothalamic-pituitary-thyroid axis, the limited human studies are inconsistent⁷⁻¹². Most of the human studies were population-based on background exposure to phthalates among pregnant women and children⁹⁻¹⁶. The effects of exposure to high phthalate concentrations, specifically on thyroid function, has not been studied.

Ortho-phthalates (hereafter referred to as phthalates) are a commonly used high-production family of chemicals to which the general population are ubiquitously exposed¹⁷. Phthalates are present in personal care products¹⁸, medical equipment, foods, water and house dust¹⁷. Dibutyl phthalate (DBP)¹⁷ is also used for enteric coating in some specific mesalamine formulations¹⁹⁻²³ despite the recent US Food and Drug Administration (FDA) recommendation against the use of phthalates in drug delivery vehicles²⁴. Mesalamine or 5-aminosalicylic acid (5-ASA) medications are prescribed for inflammatory bowel diseases (IBD), including ulcerative colitis (UC) and Crohn's disease (CD). Mesalamine is the active ingredient in Asacol[®] and Asacol[®]HD, with DBP acting as an excipient in their enteric coating²⁵. Other mesalamine formulations such as Pentasa[®], Lialda[®], Apriso[®], and Delzicol[®] do not contain DBP^{23,26}. Asacol[®], widely used to treat IBD in adults and children, was a first line of therapy for patients with UC and often used in pregnant women with IBD^{26,27}.

Research including ours has shown that mesalamine medications with DBP-containing coatings contribute to high-DBP exposure as measured by concentrations of urinary monobutyl phthalate (MBP), the primary DBP metabolite^{28,29}. Specifically, in individuals taking mesalamine medications that contain DBP, their MBP urinary concentrations were approximately 1,000 times higher than the median levels for men in the US general population (National Health and Nutrition Examination Survey (NHANES))³⁰. Therefore, patients with IBD taking DBP-containing mesalamine have chronic high exposure to DBP because the medication is taken daily.

We recently found that sperm motility decreased in newly exposed men to high-DBP from mesalamine and continued to decrease even after a four month withdrawal³¹ and disrupted their reproductive hormones and perhaps the hypothalamic-pituitary-testis axis¹⁸. However, to our knowledge, there are few studies that investigated the association between DBP and thyroid hormones, despite the observation that other phthalates appeared to affect the hypothalamic-pituitary-thyroid axis^{7,8}. Specifically, there are no studies on effects of high-DBP exposures from medications on hypothalamic-pituitary-thyroid axis. Therefore, as an extension of our previous work focusing on semen quality³¹ and reproductive hormones¹⁸, we took advantage of the marked difference in DBP-exposure from different mesalamine formulations using a crossover-crossback prospective cohort to assess the effects of high-DBP exposure from mesalamine medications on thyroid function and autoimmunity in adult men. We hypothesized that exposure to high-DBP is associated with a decrease in thyroid hormone concentrations and an increase in thyroid autoimmunity.

Methods

Participants

For this study, men aged 18 to 55 years old, with mild IBD, and on mesalamine for at least three months at enrollment, without history of steroid medication in the last three months, vasectomy, diabetes mellitus, hepatic, or renal diseases were eligible to participate. As previously described^{18,31}, 73 men were enrolled from gastroenterology clinics at Beth Israel Deaconess Medical Center (BIDMC), Brigham and Women's Hospital (BWH) and Massachusetts General Hospital (MGH) in Boston into the Mesalamine And Reproductive health Study (MARS) (2010–2016). We excluded 3 men who had history of thyroid disease or radiation therapy leaving 70 men in our current analysis. The institutional review boards of Harvard T.H. Chan School of Public Health, BIDMC, BWH and MGH approved MARS. All men signed informed consents before participation.

Study design

Men participated in up to six visits (2 baseline, 2 crossover, and 2 crossback). B₁H_B₂-arm (Background₁-High-Background₂) represents men who started on non-DBP mesalamine (background-DBP exposure from other sources) then crossed-over to high-DBP mesalamine (high-DBP) then crossed-back to non-DBP mesalamine (background-DBP) and vice versa in H₁BH₂-arm (High₁-Background-High₂) men who started on high-DBP mesalamine (Figure 1). Because the major aim of this study was to investigate semen parameters³¹, four month crossover and crossback periods were chosen *a priori* taking into account the

spermatogenesis cycle³². The duration between the two visits in the baseline, crossover and crossback phase were two weeks. At time 0 (T0), men reported their medications, demographics, lifestyle and health information and our research nurse measured their height and weight. At each visit, men reported their mesalamine medications and gave blood samples. Twelve out of the 44 men in the H₁BH₂-arm participated in a short protocol requiring only up to four visits that only required crossover as previously described and justified³¹. These 12 men participated in 29 baseline and crossover visits and were also included in the analysis.

Exposure assessment

We defined DBP exposure by medication type, i.e., high-DBP mesalamine or non-DBP mesalamine, in each visit as prescribed and self-reported which has been shown to be a reliable method for medication compliance^{18,31,33}. We *a priori* chose to define high-DBP exposure by the mesalamine medication type^{18,31} rather than measuring urinary MBP concentrations that, due to the short DBP half-life³⁴, only represent exposure over the past several hours.

Thyroid function assessment

At each visit, nurses collected non-fasting blood samples between 7 am and 2 pm. All samples were stored at -80° C and shipped in a single batch for analysis to the laboratory of clinical chemistry and haematology at the Máxima Medical Centre in Veldhoven, The Netherlands. We measured TSH, FT4, total thyroxine (T4), total triiodothyronine (T3), free triiodothyronine (FT3), thyroperoxidase antibodies (TPOAbs) and thyroglobulin antibodies (TgAbs) using a electrochemiluminescence immunoassay (Cobas e601, Roche Diagnostics, Mannheim Germany) in serum. The clinical reference values were 0.4 – 4.0 mU/L for TSH, 0.9 – 2.8 nmol/L for T3, 3.5 – 6.5 pmol/L for FT3, 58 – 161 nmol/L for T4, and 10 –24 pmol/L for FT4. Inter and intra-assay coefficients of variation were 2.1%, 3.5%, 3.8%, 3.8%, and 7.7% for TSH, FT4, T4, FT3 and T3, respectively. TPOAb and TgAb concentrations were considered positive if >35 IU/ml or >115 IU/ml (manufacturer cut-offs), respectively, and the coefficients of variation were 12.4% and 7.1% for TPOAbs at 33 or 100 IU/l, respectively, 10.9% and 8.6% for TgAbs at 76 and 218 IU/l, respectively. In addition, we calculated the ratio (%) of serum total T3 to serum total T4 (T3/T4) from the respective hormone concentrations (T₃ (nmol/L)/T₄ (nmol/L) * 100) as an index of the action of thyroid hormones in peripheral tissues³⁵ and employed previously by Johns et al.³⁶. Serum total T₄ is maintained by a balance between T₄ secretion from the thyroid gland and its clearance from blood (half-life of 7–10 days), whereas serum total T₃ is largely controlled by peripheral deiodination in tissues as it becomes activated³⁷. Others have shown that the T₃/T₄ ratio is a good measure of thyroid state in health and disease³⁸. Therefore, when the ratio decreases, this suggests that deiodination decreased and vice versa.

Statistical Analysis

We defined the exposure as an indicator variable based on DBP-exposure (high versus background) at each phase (baseline, crossover and crossback) for the two study arms (H₁BH₂ and B₁HB₂). We modeled the thyroid function measurements as natural log-transformed continuous outcomes to satisfy model assumptions.

We performed descriptive statistics and tested for differences between men in the two arms using Fisher's exact test for categorical variables and Kruskal-Wallis test for continuous variables. We selected the covariates based on directed acyclic graph (Supplementary Fig 1) and statistical considerations (>10% change in the effect estimate). The final model included age (years, continuous), body mass index (BMI) (Kg/m², continuous), sample drawing time (hours, continuous, time varying), race (Caucasian or not), current smoking (binary), and duration on high-DBP mesalamine at T0 (years, continuous). Initially, we also considered adjustment for IBD severity, IBD diagnosis (UC/CD), duration since diagnosis, and education, but these variables were not confounders and thus not retained in the final models.

We estimated the crossover (crossover versus baseline), and crossback (crossback versus crossover) percent changes in the thyroid hormones within each arm separately. We also examined the cross-sectional differences at T0 between arms. We used linear mixed effect models (LMEM) with a random intercept to account for within-person correlation among longitudinal measures of a given hormone outcome arising from man-to-man heterogeneity across the study participants. We tested for effect modification by duration under high-DBP at T0 by adding an interaction term and we used the same alpha level as for the other analyses (0.05).

We applied fixed effect models (FEM) as a secondary analysis to estimate the terms as ordinary fixed regression coefficients, rather than assuming a random distribution for the person-specific intercepts. These models isolate the purely longitudinal within-person effect of exposure³⁷. As a sensitivity analysis, we assessed LMEM model sensitivity to the covariance structure implied by the random intercept model, using robust empirical standard errors³⁹. In addition, we conducted the analysis after further adjusting for the antibodies concentrations, one at a time. We also conducted the analysis after excluding men with high observations of the antibodies, suspecting that they may have undiagnosed thyroid diseases. We considered two-sided alpha <0.05 as statistically significant. We conducted all statistical analyses using SAS version 9.4 (SAS Institute Inc., Cary, NC).

LMEM equation:

$$Y_{it} = \beta_0 + \beta_1 \text{med}_{it} + \beta_2 Z_{it} + \gamma W_i + u_i + \varepsilon_{it}$$

Where Y_{it} denotes the outcome for subject i at time t (baseline, crossover, or crossback), med_{it} is an indicator denoting DBP/non-DBP mesalamine medication, Z_{it} is a vector of time-varying variables e.g., sample drawing time, W_i is a vector of non-time varying variables e.g., race, β_0 and β_1 are scalar parameters, β_2 and γ are vectors of coefficients, and u_i is random subject-specific errors and ε_{it} is within-subject errors.

FEM equation:

$$Y_{it} = \beta_0 + \beta_1 \text{med}_{it} + \beta_2 Z_{it} + u_i + \varepsilon_{it}$$

Where u_i is a fixed regression coefficient reflecting the constant effect for all measurements on a given subject. Note: u_i controls for all fixed differences in outcome across subject, measured or unmeasured.

Results

Seventy men (mean age 34.7 years) with no thyroid disease or any radiation history were enrolled and provided 327 blood samples (average: five samples/man) (Supplementary Fig. 2). Among the 58 men who participated in the full-protocol, 49 men (85%) crossed-over and 45 (78%) crossed-back; 89% and 77%, respectively, in the B₁HB₂-arm, with 81% and 78%, respectively, in the H₁BH₂-arm. Men in the B₁HB₂-arm (26 men, 134 samples) were on non-DBP mesalamine medication at baseline for a median of 1 year. Men in the H₁BH₂-arm (44 men, 193 serum samples) were on their high-DBP mesalamine medication at baseline (H₁) for a median duration of 3 years. Men's characteristics did not differ significantly between the two arms except for smoking where 85% of the men in the B₁HB₂ arm and 75% of the men of H₁BH₂ arm never smoked (Table 1). Nine men were lost to follow-up, three in B₁HB₂-arm and six in H₁BH₂-arm (p-value= 0.64). All thyroid function distributions are shown in Table 2.

When the B₁HB₂-arm (26 men, 134 samples) (Table 3A and Figure 2) crossed-over, concentrations of T3 decreased 10% (95% confidence interval (CI): -14%, -5%), T3/T4 ratio decreased 8% (CI: -12%, -3%), TPOAb, and TgAb decreased 11% (CI: -20%, -2%) and 15% (CI: -23%, -5%), respectively. After crossback, T3 increased 8% (CI: 2%, 14%), T3/T4 ratio increased 7% (CI: 3%, 13%), and FT4 decreased 5% (CI: -8%, -2%). Both TPOAb, and TgAb increased 16% (CI: 4%, 29%) and 39% (CI: 24%, 57%) respectively. There was no significant change in T4, TSH, or FT3.

When the men in the H₁BH₂-arm (44 men, 193 samples) (Table 3B and Figure 2) crossed-over, T3 decreased by 7% (CI: -11%, -2%) and T3/T4 ratio decreased by 6% (CI: -9%, -2%) while the changes in other thyroid parameters were not statistically significant. After crossback, FT4 decreased 5% (CI: -7%, -2%), TSH decreased 13.2% (CI: -25%, 0%) whereas TgAb increased 37% (CI: 24%, 52%).

In the cross-sectional adjusted analysis at T0, on average men in B₁HB₂-arm had higher FT3, FT4, T3, and T4 and lower TSH concentrations compared to men in H₁BH₂-arm, but the difference was not statistically significant (Supplementary Table 1). The duration of high-DBP mesalamine intake at T0 modified the changes in T3, T3/T4 ratio and TgAb (p, interaction= 0.007, 0.006, 0.004) respectively (data not shown). For example, among men with a history of no high-DBP mesalamine at T0 (i.e., B₁HB₂-arm) or men with shorter duration of high-DBP mesalamine at T0 in the H₁BH₂-arm, high-DBP exposure was associated with lower T3, T3/T4 ratio and TgAb. This association was the opposite among the men in the H₁BH₂-arm with longer duration of high-DBP mesalamine at T0. The results were consistent in all sensitivity analyses (Supplementary Table 2 to 5) using the fixed effect models, robust empirical standard errors, after further adjusting for the antibodies concentrations one at a time and after excluding men with observation of high antibodies leaving 305 samples from 66 men (excluding two men from each arm).

Discussion

In the current study, men who were newly-exposed to high-DBP mesalamine in the B₁HB₂-arm exhibited a decrease in serum total T3, serum T3/T4 ratio, and both TPOAb and TgAb concentrations relative to T0. When these men crossed-back to non-DBP mesalamine, all of these measures returned toward their T0 concentrations while FT4 decreased. Because 80% of serum T3 is derived from peripheral T4 conversion to T3, these findings are consistent in part with the interpretation that DBP interferes with this conversion. Thus, DBP would decrease serum T3 and the T3/T4 ratio and would tend to increase serum T4 as a result, though increased T4 was not observed. Reducing DBP exposure among the newly-exposed men would be expected to reverse these trends. It is not immediately clear why antibodies for TPO and Tg decreased.

The decrease in T3 but not T4 in men newly exposed to high-DBP suggests that new high-DBP exposure could affect type 1 and/or type 2 deiodinase that is responsible for deiodination of T4 to T3. This may occur by reducing Dio1/Dio2 expression or by directly inhibiting Dio1/Dio2 activity. Another explanation could be that decrease in the T3/T4 ratio were mediated by changes in homeostasis of growth hormone and consequently insulin-like growth factor 1 (IGF-1). A decreased IGF-1 could lead to decrease in T4 to T3 conversion as hypothesized and observed before ^{40,41}. Although we cannot confirm from this in our study, Boas et al., observed a negative association between concentrations of urinary MBP and serum IGF-1 and T3 in Danish children ¹¹. Alternatively, DBP may alter thyroid function in subtle ways that we don't currently understand.

Paradoxically, in men chronically exposed to high-DBP from mesalamine medications, removal of high-DBP mesalamine medication was also associated with a decrease in T3 and a decrease in the T3/T4 ratio while a subsequent restart of the high-DBP mesalamine was associated with a decrease in FT4 and an increase in TgAb concentrations. One potential interpretation is that long-term exposure to DBP could cause 'adaptation' in the thyroid system that may cause these paradoxical findings after removal of high-DBP exposure. This is supported by the observation of significant interactions of medication type (DBP exposure) and the duration on high-DBP mesalamine use at T0 on the response of T3, T3/T4 ratio and TgAb. For example, among men with history of no (B₁HB₂-arm) or shorter duration of high-DBP mesalamine at T0, high-DBP exposure was associated with decreased T3, T3/T4 ratio and TgAb concentrations. However, in the longer term exposed men in the H₁BH₂-arm, subsequent high-DBP exposure was associated with increased T3, T3/T4 ratio and TgAb concentrations. Overall, these results in both arms suggest that new high-DBP exposure or removal from chronic high-DBP exposure alters the thyroid system and thyroid autoimmunity, consistent with thyroid disruption. The seemingly similar changes in both arms may suggest that high-DBP exposure changes may have met the threshold for "change" in either direction.

In addition, in cross-sectional analysis, men exposed to high-DBP concentrations (H₁BH₂-arm) at T0 had lower T4, T3 and, FT4 concentrations and higher TSH concentrations compared to men not exposed to high-DBP concentrations (B₁HB₂-arm). It is important to note that the cross-sectional results may be due to chance sampling

variability due to non-randomization at T0. Finally, although unlikely, we cannot exclude the possibility of a non-causal explanation for the similar trends in the two arms such as an unmeasured time-varying confounding or changes in health behaviors due to participation in the study.

Only a few studies have investigated the association of DBP exposure with thyroid hormones in adult men. These studies were limited by their cross-sectional study design^{7,8,42} or by restricting to infertile/subfertile men⁸, or Dirtu et al⁴³ who investigated the association in obese and control lean group. None of these studies measured thyroid antibodies. Consistent with our results for the newly exposed men, Park et al.,⁴² observed that urinary MnBP concentrations were significantly associated with lower T3 in adult Korean men. Janjua et al.,³⁴ conducted the only human study with some similarity to our design, examining the association between dermal DBP-exposure and short term effects on thyroid hormones, concluding that their results were most likely due to chance. The dermal DBP-exposure was much lower compared to our study, and one week as washout period was likely too short compared to the four months in the current study. Clearly, more research is needed in the adult men.

Our study had some limitations. First, we did not randomize men to mesalamine formulations at T0. However, this should not influence the within person comparison of interest. Second, although our sample size was not very large due to the innovative study and length of participation, the unique design of MARS and the use of the men as their own control avoided the purely cross-sectional analysis that could be confounded by inter-individual variables. Third, we did not have data on the iodine status of our participants, which is an essential trace element for thyroid function⁴⁴. Although some studies observed correlations between urinary iodine and phthalate exposure, it is unclear whether an individual's phthalate exposure directly influences iodine status (note: low iodine = elevated T₃). In a study conducted among a representative sample of U.S. adults including men, iodine excretion had a negligible impact on the significant correlations observed between phthalate metabolites and thyroid hormone levels⁴⁵. In addition, iodine status is unlikely to be associated with the prescribed mesalamine medications and, most importantly, is expected to be constant within the same man. Therefore, iodine status is unlikely to confound our longitudinal results. Also, because mesalamine medications (with or without DBP coating) prescribed for treating IBD have the same active ingredient (mesalamine), confounding by indication was unlikely. Finally, there may be a concern of generalizing results from men with IBD to men in the general population due to the possible relation between IBD and thyroid disease⁴⁶. Results are still important at least for IBD patients as a growing disease especially in the developed world.

Our study had several strengths. First, the innovative prospective design is rarely conducted in environmental epidemiology studies. The design enabled us to compare the hormone concentrations within the same men during periods of high-DBP and background-DBP exposure while accounting by design for confounding by measured and unmeasured non-time-varying variables^{18,31}. Second, having two arms with different sequences of exposure and non-exposure allowed us to determine the reversibility of the changes after exposure removal and observe if the exposure history may change the hormonal response. Third, we

had comprehensive measures for both free and total thyroid hormones, TSH and, in addition, thyroid antibodies. Finally, this is the first study to examine the thyroid hormone response to such high-DBP exposure (1,000 times higher than background) compared to environmental background exposure, and to examine the reversibility of the association when the source of high-DBP exposure was removed.

DBP and other phthalates are still used in the enteric coatings of some licensed and over-the-counter medications including specific mesalamine formulations in the US^{23,28,29}, Canada¹⁹, United Kingdom⁴⁷, Germany²⁰ and China⁴⁸. Mesalamine, the active ingredient in Asacol[®], Asacol[®]HD, Pentasa[®], Lialda[®], Apriso[®], and Delzicol[®], is commonly prescribed for IBD patients. The enteric coating of Asacol[®], and Asacol[®]HD contains DBP as an excipient^{49,50}, whereas it is not used in other mesalamine formulations²³. Although manufacturers are required to disclose the inactive ingredients in the drug product labeling, they are not required to do so when the composition has a patented delivery mechanism (U.S. Pharmacopeial Convention 2009). The FDA has information on the type and amount of phthalates used in specific approved drug products, however this information is generally not publically available²³.

There are no guidelines for use of mesalamine or any other medications that contain phthalate coatings in men. Recently, there was a consensus that women with IBD who are pregnant or trying to get pregnant should avoid high-DBP mesalamine¹⁹. Our results are of a concern because thyroid function is critical for body functions, such as energy level, heart rate and fertility⁵¹ and, on a population level, the observed changes would be important. Undiagnosed thyroid disease may put patients at risk for certain serious conditions, such as cardiovascular diseases, osteoporosis and infertility^{51,52}. Therefore, our results support the need to reconsider the use of DBP-containing mesalamine and other DBP-containing medications for men, as well as facilitated education and access to information for physicians and patients.

Conclusion

Some medications are considered a source of high-DBP exposure including mesalamine. These medications are prescribed with little attention given to the potential health effects from such high overlooked exposure source. Our study is the first to examine the association between high-DBP exposure from mesalamine and thyroid hormones using the crossover-crossback design. Our results suggest that high-DBP exposure for four months or removal from chronic high-DBP exposure for four months could alter elements of the thyroid system most probably due to alteration of the peripheral T4 conversion to T3 and thyroid autoimmunity, consistent with thyroid disruption. After four months of exposure removal, these trends were mostly reversed. More research is needed to determine the underlying mode of action of high-DBP and the effect of exposure duration on thyroid functions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors gratefully thank the study participants, all members of the MARS team especially Ramace Dadd, Pat Morey, and Myra Keller and the clinical staff.

Support:

a pilot grant for this work from NIEHS center at the Harvard T. H. Chan School of Public Health and NIH: R01ES017285 & P30 ES000002.

Disclosure Statement

J.R.K., receives research support from Abbvie, Transparency Life Sciences and Takeda, consults for Abbvie, and is a founder of a company called ColonyConcepts. A.C.M. has no conflict currently and had previously received prior research grants from Shire, Salix and Aptalis, manufacturers of mesalamine medications. F.L.N., T.I.M.K., B.A.C., N.E.S., S.A.K., M.E., E.J.H., J.B.F., R.A.DP., M.A.B., T.R.Z., and R.H. have nothing to declare.

References

1. Chaker L, Ligthart S, Korevaar TI, et al. Thyroid function and risk of type 2 diabetes: a population-based prospective cohort study. *BMC medicine*. 2016;14(1):150. [PubMed: 27686165]
2. Chaker L, Wolters FJ, Bos D, et al. Thyroid function and the risk of dementia: The Rotterdam Study. *Neurology*. 2016;87(16):1688–1695. [PubMed: 27638924]
3. Chaker L, Heeringa J, Dehghan A, et al. Normal Thyroid Function and the Risk of Atrial Fibrillation: the Rotterdam Study. *The Journal of clinical endocrinology and metabolism*. 2015;100(10):3718–3724. [PubMed: 26262438]
4. Boas M, Feldt-Rasmussen U, Main KM. Thyroid effects of endocrine disrupting chemicals. *Molecular and cellular endocrinology*. 2012;355(2):240–248. [PubMed: 21939731]
5. Liu C, Zhao L, Wei L, Li L. DEHP reduces thyroid hormones via interacting with hormone synthesis-related proteins, deiodinases, transthyretin, receptors, and hepatic enzymes in rats. *Environmental science and pollution research international*. 2015;22(16):12711–12719. [PubMed: 25913319]
6. Zhai W, Huang Z, Chen L, Feng C, Li B, Li T. Thyroid endocrine disruption in zebrafish larvae after exposure to mono-(2-ethylhexyl) phthalate (MEHP). *PloS one*. 2014;9(3):e92465. [PubMed: 24658602]
7. Meeker JD, Ferguson KK. Relationship between urinary phthalate and bisphenol A concentrations and serum thyroid measures in U.S. adults and adolescents from the National Health and Nutrition Examination Survey (NHANES) 2007–2008. *Environmental health perspectives*. 2011;119(10):1396–1402. [PubMed: 21749963]
8. Meeker JD, Calafat AM, Hauser R. Di(2-ethylhexyl) Phthalate Metabolites May Alter Thyroid Hormone Levels in Men. *Environmental health perspectives*. 2007;115(7):1029–1034. [PubMed: 17637918]
9. Johns LE, Ferguson KK, McElrath TF, Mukherjee B, Meeker JD. Associations between Repeated Measures of Maternal Urinary Phthalate Metabolites and Thyroid Hormone Parameters during Pregnancy. *Environmental health perspectives*. 2016.
10. Huang PC, Kuo PL, Guo YL, Liao PC, Lee CC. Associations between urinary phthalate monoesters and thyroid hormones in pregnant women. *Human reproduction (Oxford, England)*. 2007;22(10):2715–2722.
11. Boas M, Frederiksen H, Feldt-Rasmussen U, et al. Childhood exposure to phthalates: associations with thyroid function, insulin-like growth factor I, and growth. *Environmental health perspectives*. 2010;118(10):1458–1464. [PubMed: 20621847]
12. Jung H, Hong Y, Lee D, Pang K, Kim Y. The association between some endocrine disruptors in human plasma and the occurrence of congenital hypothyroidism. *Environmental toxicology and pharmacology*. 2013;35(2):278–283. [PubMed: 23376177]

13. Brucker-Davis F, Ferrari P, Boda-Buccino M, et al. Cord Blood Thyroid Tests in Boys Born With and Without Cryptorchidism: Correlations with Birth Parameters and In Utero Xenobiotics Exposure. *Thyroid*. 2011;21(10):1133–1141. [PubMed: 21875366]
14. Tsai HJ, Wu CF, Tsai YC, et al. Intake of Phthalate-tainted Foods and Serum Thyroid Hormones in Taiwanese Children and Adolescents. *Scientific reports*. 2016;6:30589. [PubMed: 27470018]
15. Huang PC, Tsai CH, Liang WY, Li SS, Huang HB, Kuo PL. Early Phthalates Exposure in Pregnant Women Is Associated with Alteration of Thyroid Hormones. *PloS one*. 2016;11(7):e0159398. [PubMed: 27455052]
16. Huang HB, Chuang CJ, Su PH, et al. Prenatal and Childhood Exposure to Phthalate Diesters and Thyroid Function in a 9-Year Follow-up Birth Cohort Study: Taiwan Maternal and Infant Cohort Study. *Epidemiology*. 2017;28 Suppl 1:S10–s18. [PubMed: 29028671]
17. CDC. (Centers for Disease Control and Prevention). National Report on Human Exposure to Environmental Chemicals. 2018; <https://www.cdc.gov/exposurereport/>. Accessed July 2018.
18. Nassan FL, Coull BA, Skakkebaek NE, et al. A crossover-crossback prospective study of dibutylphthalate exposure from mesalamine medications and serum reproductive hormones in men. *Environmental research*. 2018;160:121–131. [PubMed: 28978458]
19. Nguyen GC, Seow CH, Maxwell C, et al. The Toronto Consensus Statements for the Management of Inflammatory Bowel Disease in Pregnancy. *Gastroenterology*. 2016;150(3):734–757.e731. [PubMed: 26688268]
20. Wittassek M, Koch HM, Angerer J, Bruning T. Assessing exposure to phthalates - the human biomonitoring approach. *Molecular nutrition & food research*. 2011;55(1):7–31. [PubMed: 20564479]
21. Hernandez-Diaz S, Mitchell AA, Kelley KE, Calafat AM, Hauser R. Medications as a potential source of exposure to phthalates in the U.S. population. *Environmental health perspectives*. 2009;117(2):185–189. [PubMed: 19270786]
22. Seckin E, Fromme H, Volkel W. Determination of total and free mono-n-butyl phthalate in human urine samples after medication of a di-n-butyl phthalate containing capsule. *Toxicology letters*. 2009;188(1):33–37. [PubMed: 19433267]
23. Kelley KE, Hernandez-Diaz S, Chaplin EL, Hauser R, Mitchell AA. Identification of phthalates in medications and dietary supplement formulations in the United States and Canada. *Environmental health perspectives*. 2012;120(3):379–384. [PubMed: 22169271]
24. FDA. (U.S. Food and Drug Administration) Limiting the use of certain phthalates as excipients in CDER-regulated products. 2012; <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm330792.htm>. Accessed March 2016.
25. FDA. Summary Review for Regulatory Action, Division Director Review NDA 204412, Division of Gastroenterology and Inborn Errors Products, CENTER FOR DRUG EVALUATION AND RESEARCH 2012; http://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/204412Orig1s000SumR.pdf Accessed January, 2016.
26. Gallinger ZR, Nguyen GC. Presence of phthalates in gastrointestinal medications: is there a hidden danger? *World J Gastroenterol*. 2013;19(41):7042–7047. [PubMed: 24222946]
27. Khan N, Abbas AM, Koleva YN, Bazzano LA. Long-term mesalamine maintenance in ulcerative colitis: which is more important? Adherence or daily dose. *Inflamm Bowel Dis*. 2013;19(6):1123–1129. [PubMed: 23514878]
28. Hauser R, Duty S, Godfrey-Bailey L, Calafat AM. Medications as a source of human exposure to phthalates. *Environmental health perspectives*. 2004;112(6):751–753. [PubMed: 15121520]
29. Hait EJ, Calafat AM, Hauser R. Urinary phthalate metabolite concentrations among men with inflammatory bowel disease on mesalamine therapy. *Endocr Disruptors (Austin)*. 2014;1(1).
30. CDC. Centers for Disease Control and Prevention, Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables, February 2015 2015; http://www.cdc.gov/biomonitoring/pdf/FourthReport_UpdatedTables_Feb2015.pdf Accessed February 2016.
31. Nassan FL, Coull BA, Skakkebaek NE, et al. A crossover-crossback prospective study of dibutylphthalate exposure from mesalamine medications and semen quality in men with inflammatory bowel disease. *Environment international*. 2016;95:120–130. [PubMed: 27575365]

32. Heller CG, Clermont Y. Spermatogenesis in man: an estimate of its duration. *Science (New York, NY)*. 1963;140(3563):184–186.
33. Gifford AE, Berg AH, Lahiff C, Cheifetz AS, Horowitz G, Moss AC. A random urine test can identify patients at risk of mesalamine non-adherence: a prospective study. *Am J Gastroenterol*. 2013;108(2):249–255. [PubMed: 23295279]
34. Janjua NR, Mortensen GK, Andersson AM, Kongshoj B, Skakkebaek NE, Wulf HC. Systemic uptake of diethyl phthalate, dibutyl phthalate, and butyl paraben following whole-body topical application and reproductive and thyroid hormone levels in humans. *Environmental science & technology*. 2007;41(15):5564–5570. [PubMed: 17822133]
35. Dietrich JW, Landgrafe G, Fotiadou EH. TSH and Thyrotropic Agonists: Key Actors in Thyroid Homeostasis. *Journal of thyroid research*. 2012;2012:351864. [PubMed: 23365787]
36. Johns LE, Ferguson KK, McElrath TF, Mukherjee B, Meeker JD. Associations between Repeated Measures of Maternal Urinary Phthalate Metabolites and Thyroid Hormone Parameters during Pregnancy. *Environmental health perspectives*. 2016;124(11):1808–1815. [PubMed: 27152641]
37. Jameson JL. *Endocrinology : adult & pediatric*. 7th edition. ed. Philadelphia, PA: Elsevier Saunders; 2016.
38. Mortoglou A, Candiloros H. The serum triiodothyronine to thyroxine (T3/T4) ratio in various thyroid disorders and after Levothyroxine replacement therapy. *Hormones (Athens, Greece)*. 2004;3(2):120–126.
39. Fitzmaurice GM, Laird NM, Ware JH. *Applied Longitudinal Analysis*. 2nd ed 2011.
40. Huang HB, Pan WH, Chang JW, et al. Does exposure to phthalates influence thyroid function and growth hormone homeostasis? The Taiwan Environmental Survey for Toxicants (TEST) 2013. *Environmental research*. 2017;153:63–72. [PubMed: 27907809]
41. Jorgensen JO, Pedersen SA, Laurberg P, Weeke J, Skakkebaek NE, Christiansen JS. Effects of growth hormone therapy on thyroid function of growth hormone-deficient adults with and without concomitant thyroxine-substituted central hypothyroidism. *The Journal of clinical endocrinology and metabolism*. 1989;69(6):1127–1132. [PubMed: 2685007]
42. Park C, Choi W, Hwang M, et al. Associations between urinary phthalate metabolites and bisphenol A levels, and serum thyroid hormones among the Korean adult population - Korean National Environmental Health Survey (KoNEHS) 2012–2014. *The Science of the total environment*. 2017;584–585:950–957.
43. Dirtu AC, Geens T, Dirinck E, et al. Phthalate metabolites in obese individuals undergoing weight loss: Urinary levels and estimation of the phthalates daily intake. *Environment international*. 2013;59:344–353. [PubMed: 23892227]
44. Zimmermann MB, Kohrle J. The impact of iron and selenium deficiencies on iodine and thyroid metabolism: biochemistry and relevance to public health. *Thyroid*. 2002;12(10):867–878. [PubMed: 12487769]
45. Mendez W, Jr., Eftim SE. Biomarkers of perchlorate exposure are correlated with circulating thyroid hormone levels in the 2007–2008 NHANES. *Environmental research*. 2012;118:137–144. [PubMed: 22749113]
46. Shizuma T Concomitant Thyroid Disorders and Inflammatory Bowel Disease: A Literature Review. *BioMed research international*. 2016;2016:5187061. [PubMed: 27042663]
47. Jamieson L, McCully W. Review: UK medicines likely to be affected by the proposed European Medicines Agency's guidelines on phthalates. *BMC pharmacology & toxicology*. 2015;16:17. [PubMed: 26070463]
48. Jia LL, Lou XY, Guo Y, Leung KS, Zeng EY. Occurrence of phthalate esters in over-the-counter medicines from China and its implications for human exposure. *Environment international*. 2016.
49. FDA. ASACOL® (mesalamine) delayed-release tablets, for oral use - HIGHLIGHTS OF PRESCRIBING INFORMATION. 2015; <http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm215476.htm>, http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/019651s025lbl.pdf Accessed April, 2017.
50. FDA. Asacol® HD (mesalamine) delayed-release tablet for oral administration-HIGHLIGHTS OF PRESCRIBING INFORMATION. 2010; http://www.accessdata.fda.gov/drugsatfda_docs/label/2010/021830s005lbl.pdf Accessed April, 2017.

51. (ATA) ATA. Prevalence and Impact of Thyroid Disease. 2017; <http://www.thyroid.org/media-main/about-hypothyroidism/>. Accessed November, 2017.
52. Patel N, Kashanian JA. Thyroid Dysfunction and Male Reproductive Physiology. *Seminars in reproductive medicine*. 2016;34(6):356–360. [PubMed: 27741548]

Figure 1. Mesalamine And Reproductive health Study (MARS) Design.

B₁HB₂-arm: B₁ represents background-exposure at baseline, H represents high-DBP exposure after crossover and B₂ represents background-exposure after crossback; included 26 men taking mesalamine that did not contain DBP at baseline. H₁BH₂-arm: H₁ represents high-DBP exposure at baseline, B represents background-exposure after crossover and H₂ represents high-DBP exposure after crossback; included 44 men taking mesalamine containing DBP at baseline (24 men for less than 3 years and 20 men with 3 years or more).

Abbreviations: MARS, Mesalamine And Reproductive health Study; DBP, dibutyl phthalate.

Figure 2. Adjusted geometric means (95%CI) of thyroid hormone parameters at baseline, crossover and crossback for the 2 arms (B₁HB₂- arm, and H₁BH₂-arm).

^a Adjusted for race (Caucasian or not), age (continuous), BMI (continuous), hour of sample draw (continuous), current smoking, and duration on DBP-containing mesalamine medication at baseline (in years) in the mixed effect model.

^b Geometric means are presented for TSH, TPOAb, and TgAb.

^c B₁HB₂-arm: B₁ represents background low-DBP exposure at baseline, H represents high-DBP exposure after crossover and B₂ represents background low-exposure after crossback.

^d H₁BH₂-arm: H₁ represents high-DBP at baseline, B represents background-DBP exposure after crossover and H₂ represents high-DBP after crossback.

* Indicates statistical significance of P-value < 0.05

Abbreviations: MARS, Mesalamine And Reproductive health Study; N, number of men; B₁HB₂, Background₁-High-Background₂ DBP exposure; H₁BH₂, High₁-Background-

High₂ DBP exposure; 95% CI, 95% Confidence Interval; DBP, dibutyl phthalate; T3, Total Triiodothyronine; T4, Total thyroxine; FT3, Free Triiodothyronine; FT4.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1:

Demographics of 70 men contributing 327 blood samples in the MARS Study by arm

	B₁HB₂-arm (26 men, 134 visits)	H₁BH₂-arm (44 men, 193 visits)	Total (70 men, 327 visits)	
	N (%) / Mean (SD), [Range]	N (%) / Mean (SD), [Range]	N (%) / Mean (SD), [Range]	P-value^b
Baseline characteristics (at Visit 1), men (N)^a				
Age (Years)	33.9(8.97), [20.2, 54.6]	35.1 (9.69), [20.8, 55.7]	34.7 (9.38), [20.2, 55.7]	0.78
Race				0.23
Caucasian	25 (96)	35 (80)	60 (86)	–
Black/African American	1(4)	2(5)	3(4)	–
Asian	0	4(9)	4(6)	–
Other	0	3(7)	3(4)	–
BMI (Kg/m2)	25.3 (3.56), [19.4, 32.7]	26.3 (6.53), [19.5, 54.0]	26.0(5.61), [19.4, 54.0]	0.95
BMI-categories				0.99
Normal weight (18.5 BMI<25)	14(54)	23 (52)	37(53)	–
Overweight (25 BMI<30)	9(35)	15 (34)	24 (34)	–
Obese (BMI 30)	3(12)	6(14)	9(13)	–
Education				0.69
Below college	3(14)	4(10)	7(12)	–
College graduate	10(48)	16(41)	26 (43)	–
Graduate degree	8(38)	19(49)	27 (45)	–
Smoking status				0.007
Never smoker	22 (85)	33 (75)	55 (79)	–
Former smoker	1(4)	11(25)	12(17)	–
Current smoker	3(11)	0	3(4)	–
Current Marijuana smoke	6(46)	6(27)	12 (34)	0.29
Warm season at baseline^c	5(19)	17(39)	22(31)	0.11
IBD diagnosis				0.80
Ulcerative colitis	16(62)	29 (66)	45 (64)	–
Crohn's disease	10(38)	15(34)	25 (36)	–
IBD score^d	1.65(1.38), [0,4]	1.20(1.34), [0, 5]	1.37(1.36), [0,5]	0.15
Duration since IBD diagnosis (years)	9.00 (7.84), [0.28, 29.5]	10.7(9.24), [0.7, 39.0]	10.1 (8.73), [0.28, 39.0]	
Alcohol^e				0.18
None or< 1 days/week	7(32)	17(45)	24 (40)	–
1 or 2 days/week	13(59)	13(34)	26 (43)	–
2 days/week	2(9)	8(21)	10(17)	–
Time-varying characteristics, visits^a				
Warm season of blood draw^c	60 (45)	86 (45)	146 (45)	0.99
Hour of the blood draw	9.68(1.62), [7.25, 13.92]	9.68(1.64), [7.17, 13.5]	9.68(1.63), [7.17, 13.9]	0.97

	B₁HB₂–arm (26 men, 134 visits)	H₁BH₂–arm (44 men, 193 visits)	Total (70 men, 327 visits)	
	N (%) / Mean (SD), [Range]	N (%) / Mean (SD), [Range]	N (%) / Mean (SD), [Range]	P-value^b
Time of blood draw				0.99
7 am to 10 am	74 (55)	106(55)	180(55)	–
>10 am to 2 pm	60 (45)	87 (45)	147 (45)	–

Notes: B₁HB₂–arm, B₁ represents background-exposure at baseline, H represents high-DBP exposure after crossover and B₂ represents background-exposure after crossback; included 26 men taking mesalamine that did not contain DBP at baseline; 16 men (62%) on Lialda[®], 8 men (31%) on Pentasa[®], one man (4%) on Apriso[®] and one man (4%) was not on mesalamine medication at the time of recruitment.

H₁BH₂–arm, H₁ represents high-DBP exposure at baseline, B represents background-exposure after crossover and H₂ represents high-DBP exposure after crossback; included 44 men taking mesalamine-containing DBP at baseline; 23 men (53%) on Asacol[®] and 21 men (48%) on Asacol[®]HD.

^aN (%) for categorical/binary variables and mean (SD), [Range] for continuous variables

^bP-values are based on Fisher exact test for categorical variables and Kruskal Wallis test for continuous variables.

^cSeason of sample collection, Warm: April Through September

^dIBD score: included bowel frequency and urgency, presence of blood in the stool and general wellbeing. Mild IBD score: 5 or less on the simple clinical colitis activity index for UC and 4 or less on the Harvey-Bradshaw index for CD.

Abbreviations: MARS, Mesalamine And Reproductive health Study; BMI, body mass index; Kg, Kilogram; m, meter; SD, standard deviation; IBD, Inflammatory Bowel Disease; UC, ulcerative colitis; CD, Crohn's disease; N, number of men; DBP, dibutyl phthalate; B₁HB₂, Background₁–High-Background₂ DBP exposure; H₁BH₂, High₁–Background-High₂ DBP exposure.

Table 2.

Hormone concentrations for 70 men (327 blood samples) in the MARS study

Hormone Parameters	Arithmetic Mean	SD	Geometric means	Minimum	5th Perc	25th Perc	50th Perc	75th Perc	95th Perc	Maximum	QR	Clinical normal range
TSH (mU/L)	2.61	1.41	2.36	0.04	1.03	1.70	2.21	3.24	5.33	9.60	1.54	0.4–4.0
FT4 (pmol/l)	16.7	2.07	16.8	10.8	13.1	15.5	16.6	18.1	20.0	23.8	2.64	10.0–24.0
FT3 (pmol/l)	5.75	0.68	5.75	3.95	4.67	5.35	5.70	6.13	6.91	8.53	0.78	3.5–6.5
T4 (nmol/l)	105	16.2	105	62.8	79.4	94.5	104	116	132.8	162	21.6	58–161
T3 (nmol/l)	2.01	0.35	2.05	1.22	1.47	1.75	1.99	2.26	2.6	3.18	0.51	0.9–2.8
T3/T4 Ratio, %	1.92	0.28	1.95	1.37	1.56	1.73	1.87	2.06	2.44	2.90	0.33	–
TPOAb (IU/ml)	21.3	26.3	17.5	6.00	9.00	12.2	15.3	19.7	65.0	215.6	7.48	35
TgAb (IU/ml)	25.8	49.7	18.2	10.0	10.2	13.1	16.8	22.2	36.0	506.3	9.18	115

Abbreviations: MARS, Mesalamine And Reproductive health Study; SD, standard deviation; Perc, percentile; QR, quartile range; T3, total triiodothyronine; T4, total thyroxine; FT3, free triiodothyronine; FT4, free thyroxine; TSH, thyroid-stimulating hormone; TPOAb, thyroid peroxidase antibody; TgAb, thyroglobulin antibody.

Table 3A.

Adjusted geometric means and percent changes (95%CI) of crossover and crossback on hormone concentrations among men starting on mesalamine medication not containing dibutyl phthalate (B₁HB₂–arm)

Hormone serum concentrations	Fitted Geometric Means (95%CI)			Comparisons			
	Baseline (B ₁)	Crossover (H)	Crossback (B ₂)	Crossover effect (95%CI) H-B ₁	P value	Crossback effect (95%CI) B ₂ -H	P value
TSH(mU/L)	2.30 (1.84, 2.88)	2.30 (1.83, 2.88)	2.06 (1.63, 2.61)	-0.07 (-14.3, 16.6)	0.99	-10.3 (-23.8, 5.68)	0.19
FT4(pmol/l)	17.2 (16.4, 18.1)	17.3 (16.5, 18.2)	16.4 (15.6, 17.2)	0.62 (-2.45, 3.78)	0.70	-5.40 (-8.45, -2.25)	0.001
FT3(pmol/l)	5.84 (5.6, 6.09)	5.68 (5.45, 5.93)	5.77 (5.52, 6.03)	-2.67 (-5.66, 0.41)	0.09	1.50 (-1.79, 4.91)	0.37
T4(nmol/l)	109 (103, 116)	106 (100, 113)	107 (100, 114)	-2.33 (-6.23, 1.75)	0.26	0.40 (-3.85, 4.84)	0.86
T3(nmol/l)	2.08 (1.96, 2.22)	1.88 (1.77, 2.01)	2.03 (1.90, 2.16)	-9.69 (-14.2, -5.00)	<.0001	7.56 (1.93, 13.5)	0.008
T3/T4 Ratio	1.91 (1.82, 2.00)	1.77 (1.68, 1.86)	1.90 (1.81, 2)	-7.52 (-11.5, -3.36)	0.001	7.44 (2.54, 12.6)	0.003
TPOAb(IU/ml)	19.5 (15.1, 25.2)	17.3 (13.4, 22.4)	20.1 (15.5, 26.1)	-11.2 (-19.6, -2.00)	0.02	16.0 (4.42, 28.8)	0.006
TgAb(IU/ml)	19.1 (14.9, 24.5)	16.3 (12.8, 20.9)	22.8 (17.7, 29.3)	-14.6 (-23.4, -4.61)	0.005	39.4 (24.1, 56.6)	<.0001

Table 3B.

Adjusted geometric means and percent changes (95%CI) of crossover and crossback on hormone concentrations among men starting on mesalamine medication containing dibutyl phthalate (H₁BH₂-arm).

Hormone serum concentrations	Fitted Geometric Means (95%CI)			Comparisons			
	Baseline (H ₁)	Crossover (B)	Crossback (H ₂)	Crossover Effect (95%CI) B-H ₁	P value	Crossback Effect (95%CI) H ₂ -B	P value
TSH(mU/L)	2.39 (2.03, 2.82)	2.36 (1.97, 2.83)	2.05 (1.7, 2.47)	-1.31 (-13.7, 12.9)	0.85	-13.2 (-24.8, 0.26)	0.05
FT4(pmol/l)	16.6 (16.0, 17.2)	16.8 (16.1, 17.4)	16.0 (15.4, 16.6)	0.86 (-1.83, 3.63)	0.53	-4.53 (-7.25, -1.74)	0.002
FT3(pmol/l)	5.73 (5.55, 5.91)	5.63 (5.44, 5.83)	5.69 (5.49, 5.89)	-1.67 (-4.30, 1.03)	0.22	1.01 (-1.89, 3.99)	0.50
T4(nmol/l)	102 (97.7, 107)	101 (96.5, 106)	102 (97.0, 107)	-0.76 (-4.23, 2.84)	0.67	0.64 (-3.13, 4.55)	0.74
T3(nmol/l)	2.02 (1.93, 2.12)	1.89 (1.79, 1.99)	1.95 (1.84, 2.05)	-6.51 (-10.5, -2.30)	0.003	2.89 (-1.87, 7.88)	0.24
T3/T4 Ratio	1.98 (1.91, 2.05)	1.86 (1.79, 1.94)	1.91 (1.83, 1.99)	-5.90 (-9.40, -2.26)	0.002	2.44 (-1.70, 6.76)	0.25
TPOAb(IU/ml)	16.1 (13.4, 19.4)	15.5 (12.8, 18.9)	16.9 (13.9, 20.5)	-3.42 (-11.5, 5.37)	0.43	8.75 (-0.80, 19.2)	0.07
TgAb(IU/ml)	17.3 (14.5, 20.7)	16.6 (13.8, 20.1)	22.8 (18.9, 27.6)	-4.06 (-12.9, 5.68)	0.40	37.2 (23.9, 52.0)	<.0001

Notes:

^aAdjusted for race (Caucasian or not), age (continuous), BMI (continuous), hour of sample draw (continuous), current smoking, and duration on DBP-containing mesalamine medication at baseline (in years) in the mixed effect model.

^bPercentage changes are presented by exponentiating the beta coefficient. Negative sign means % decrease for the corresponding outcome compared to the measure in the previous period.

^c**B₁HB₂-arm:** B₁ represents background low-DBP exposure at baseline, H represents high-DBP exposure after crossover and B₂ represents background low-exposure after crossback.

^d**H₁BH₂-arm:** H₁ represents high-DBP at baseline, B represents background-DBP exposure after crossover and H₂ represents high-DBP after crossback.

Abbreviations: MARS, Mesalamine And Reproductive health Study; N, number of men; B₁HB₂, Background₁-High-Background₂ DBP exposure; H₁BH₂, High₁-Background-High₂ DBP exposure; 95% CI, 95% Confidence Interval; DBP, dibutyl phthalate; T3, Total Triiodothyronine; T4, Total thyroxine; FT3, Free Triiodothyronine; FT4, Free thyroxine; TSH, Thyroid-stimulating hormone; TPOAb, Thyroid peroxidase antibody; TgAb, Thyroglobulin antibody.