Received:
10 October 2017

Revised:
22 February 2018

Accepted:
19 March 2018

<https://doi.org/10.1259/bjr.20170770>

Cite this article as:

Coupal TM, Munk PL, Ouellette HA, Al-Shikarchy H, Mallinson PI, Choudur H. Popping the cap: the constellation of MRI findings in patellofemoral syndrome. *Br J Radiol* 2018; **91**: 20170770.

PICTORIAL REVIEW

Popping the cap: the constellation of MRI findings in patellofemoral syndrome

¹TYLER M COUPAL, MD, ^{1,2}PETER L MUNK, MD, ¹HUGUE A OUELLETTE, MD, ¹HISHAM AL-SHIKARCHY, MD, ¹PAUL I MALLINSON, MBChB and HEMA CHOUDUR, MD

¹Department of Radiology, Vancouver General Hospital, Vancouver, BC, Canada

²Department of Radiology, Hamilton General Hospital, Hamilton, ON, Canada

Address correspondence to: Dr Tyler M Coupal
E-mail: tyler.coupal@medportal.ca

ABSTRACT

Patellofemoral syndrome (PFS) is a common etiology of anterior knee pain, particularly among young female athletes. Despite recent advancements in the resolution of MRI, there still remains a paucity of literature that has investigated the MRI findings associated with PFS. This pictorial essay will describe our institution's experience with cases of PFS and review what we believe to be a constellation of findings associated with this diagnostic entity. This review will also describe common pitfalls encountered during the diagnosis of PFS.

INTRODUCTION

Anterior knee pain is a commonly encountered clinical scenario for radiologists and clinicians alike, with an estimated incidence of 22/1000 patients per year.¹ Very often, this pain is localised to the peri- or retropatellar region and will be exacerbated by activities that repetitively increase patellofemoral joint compressive forces, such as running, squatting or cycling. This entity is clinically known as patellofemoral syndrome (PFS) and is very common among young female athletes. In fact, it has been estimated that PFS comprises up to 40% of all sports medicine consultations.¹ PFS is an important diagnostic entity, due to the fact that it can lead to chronic pain, limitation of activity and can predispose patients to the development of early osteoarthritis.² Unfortunately, while there has been widespread consensus that the development of PFS is multifactorial in etiology, there has yet to be clarity on its exact pathogenesis. As such, it is often considered to be a diagnosis of exclusion in the setting of chronic, activity exacerbated peri- or retropatellar knee pain.¹

Many clinicians and radiologists first hypothesised that chondromalacia patella was the predominant driving force behind the development of PFS. However, initial studies which correlated arthroscopic and MRI findings were highly discordant.³ Over recent years, however, there have been dramatic improvements in the resolution of MRI, which now allows radiologists to detect increasingly subtle findings. Despite this, there still remains a paucity of literature that has investigated the MRI findings associated with

PFS.^{4,5} This pictorial essay will describe our institution's experience with cases of PFS and review what we believe to be the constellation of MRI findings associated with this diagnostic entity. This review will also describe common pitfalls encountered during the diagnosis of PFS.

THE CONSTELLATION OF MRI FINDINGS IN PATELLOFEMORAL SYNDROME

Patellar malalignment

Patellar malalignment (Figure 1) is most commonly defined as an abnormal position of the patella with respect to the femoral trochlear groove. Predisposing factors for patellar malalignment include: trochlear dysplasia, medial patellofemoral ligamentous laxity, shortness of the lateral retinaculum, patella alta and increased tibial tubercle-trochlear groove (TT-TG) distances. These morphological abnormalities manifest as the clinical entity described as "patellar instability", which is most common in young females (second decade of life) with an estimated prevalence of 6–77 per 100,000 patients.¹

TT-TG distance is measured by superimposing axial images at the level of the femoral condyles and the tibial tuberosity (Figure 2), with TT-TG distances greater than 20 mm considered to be abnormal and an indicator of patellar instability.

Trochlear dysplasia is defined as an abnormally shallow trochlea which has lost its normal concavity and quantitative methods of measuring trochlear dysplasia on MRI have

Figure 1. Axial PD-T2 fat-saturated MRI from a 22-year-old female with PFS demonstrate patellar tendon lateralisation (white arrow) with thickening and reduced signal within the medial (a, black arrow) and lateral (b, black arrow) patellofemoral ligaments due to the associated increased strain. Moreover, there is focal synovial thickening (b, white arrow head) posterior to the lateral edge of the patella, in keeping with synovitis, and mild inflammation and oedema within Hoffa's fat pad from chronic impingement (b). PFS, patellofemoral syndrome.

in fact been validated.⁶ When measuring 3.0 cm above the femorotibial joint space, a trochlear groove depth of 3 mm or less has been shown to have a sensitivity of 100% and specificity of 96%.⁶

Patella alta is defined as an abnormally high patella and can be evaluated on MRI using various quantitative methods. The Insall-Salvati ratio remains the most commonly used index, which compares the patellar tendon length to the patellar height. It has been suggested that an Insall-Salvati ratio greater than 1.5 on MRI should be considered the upper limit of normal (Table 1).⁷ The modified Insall-Salvati ratio is often used as a confirmatory index, given that variability in the shape of the patellar lower pole can affect measurements with the traditional Insall-Salvati index. The modified Insall-Salvati ratio is measured by dividing the length of the patellar tendon by the length of the articular surface

Figure 2. Axial PD-T2 MRI of the knee at 30 degrees flexion acquired at the level of the trochlea (a) and tibial tuberosity (b) demonstrating calculation of the TT-TG distance. A line is drawn across the posterior aspects of the femoral condyles. Line A is drawn perpendicular to this, traversing the deepest aspect of the trochlear groove. Line B is drawn bisecting the tibial tuberosity and TT-TG distance is calculated as the distance between lines A and B. TT-TG, tibial tubercle-trochlear groove.

Table 1. Outline of normal and pathological values of four commonly utilised measurement indices for the assessment of patellar height.

Measurement index	Normal mean value	Patella alta
Insall-Salvati	1.0	>1.5
Modified Insall-Salvati	1.25	>2.0
Blackburn-Peel	0.6	>1.0
Caton-Deschamps	1.0	>1.3

of the patella, with values greater than 2.0 being defined as patella alta (Table 1). Both of these measurement techniques, however, can prove problematic in patients with Osgood Schlatter disease and when the patellar tendon insertion site is difficult to identify. In such cases, either the Blackburn-Peel or Caton-Deschamps indices may be of utility. The Blackburn-Peel index is measured by taking the shortest distance between the inferior margin of the patellar articular surface and a line extended parallel to the proximal tibial articular surface divided by the total length of the patellar articular surface. The Caton-Deschamps technique takes a ratio of the inferior patellar articular surface length divided by the distance to the anterior-most tibial plateau. While the Blackburn-Peel and Caton-Deschamps indices can certainly prove useful in some cases, it should be noted that they are greatly distorted in the setting of osteoarthritic or post-traumatic changes.

Patellar retinacular strain

Lateralisation of the patella (Figure 1) has been shown to induce significant strain on the patellar retinaculum (Figure 3).⁸ As expected, this increased strain manifests in PFS as a thickened retinaculum, which is best appreciated on axial PD-T2 sequences (Figure 3).

Hoffa's fat pad oedema

Inflammatory changes within the fat pads of the knee are well delineated on MRI, with the most recognised entity known as Hoffa's disease, which has distinct radiological appearances depending on its chronicity. The MRI signal characteristics associated with acute injury of Hoffa's fat pad are hyperintensity on T_2 weighted images and hypointensity on T_1 weighted images due to oedema and hemorrhage (Figure 4). Moreover, there is associated fat pad enlargement with adjacent mass effect. If contrast is administered, the fat pad will demonstrate enhancement. In the chronic stages of Hoffa's disease, the fat pad becomes enlarged, which may further predispose patients to impingement. There is often hemosiderin deposition and formation of fibrous bands which manifest as hypointensity on T_1 and T_2 weighted images.

Apart from Hoffa's disease, signal abnormalities in the fat pad can also be secondary to focal fat pad impingement, which is a well described cause of anterior knee pain.⁹ A common area of focal impingement of Hoffa's fat pad is between the lateral trochlear facet and lateral patellofemoral ligament, which can result in increased T_2 and proton density signal at the superolateral aspect of the fat pad.⁹ This differs from Hoffa's disease in that there is no

Figure 3. Axial PD-T2 fat-saturated MRI from a 27-year-old female with PFS demonstrates lateralisation of the patellar tendon (white arrow) with a thickened medial patellar retinaculum (black arrow) due to the resultant increased strain. Note is also made of Hoffa's fat pad oedema in this patient. PFS, patellofemoral syndrome.

predisposing injury that results in swelling or enlargement of the fat pad. This focal fat pad impingement has been associated with patella alta, which acts as a predisposing factor.⁹

Quadriceps fat pad oedema

In the same fashion, inflammation and oedema can affect the quadriceps fat pad (Figure 5), which is also known as the supra-patellar fat pad. This will share the same MRI signal characteristic and imaging findings as Hoffa's disease (Figure 4), only instead involving the quadriceps fat pad (Figure 5).

Chondromalacia patella

Chondromalacia patella is defined as softening or breakdown of the patellar cartilage (Figure 6). Well established factors that predispose one to developing chondromalacia patella include: prior knee trauma, structural abnormalities of the patellofemoral joint, or vascular insufficiency of the subchondral bone.¹⁰ While arthroscopy is still considered the gold standard for the diagnosis of chondromalacia patella, MRI has proven itself to be an accurate, non-invasive and highly reproducible tool for the assessment of patellar cartilage.¹⁰ Moreover, MRI provides visualisation of the patellofemoral joint in its entirety, which aids in the diagnosis of other contributory factors.

The modified Outerbridge grading system developed for arthroscopy is one of the systems used to grade chondromalacia patella on MRI, classifying it into four grades.¹¹ Grade I is focal areas of hyperintensity with normal cartilaginous contour. Grade II refers to fraying or swelling of the articular cartilage that extends to

Figure 4. Sagittal PD-T2 fat-saturated MRI (a) demonstrates patellar malalignment/lateralisation (white arrow) with resultant high signal intensity within Hoffa's fat pad secondary to inflammatory oedema (black arrow). This fluid also extends into the deep infrapatellar bursa, indicative of inflammatory infrapatellar bursitis. Axial PD-T2 fat-saturated MRI (b) from another patient with PFS also shows patellar tendon lateralisation (white arrow) with marked high signal and inflammatory changes with oedema and small fluid collections within Hoffa's fat pad (black arrow) from chronic impingement. Another case (c) shows extensive oedema within the superior Hoffa's fat pad (black arrow) from malalignment of the patella and patellar tendon. Note is also made of patellar tendinosis (white arrow). PFS, patellofemoral syndrome.

the surface. Grade III is partial thickness cartilage loss with focal ulceration and Grade IV refers to full thickness cartilage loss with underlying bony changes.¹¹ Moreover, it is not uncommon to see bone marrow oedema as a part of chondromalacia patella, although this is more often seen as a later stage feature in a non-traumatic setting, as discussed subsequently.

Patellar bone marrow oedema

Much like assessment of the patellar cartilage, MRI is highly effective in demonstrating bone marrow oedema within the patella (Figures 6b and 7). This will manifest as increased T_2 signal within the bone and often presents with concomitant chondromalacia patella in non-traumatic cases such as PFS.

Focal synovitis

In addition to chondromalacia patella and underlying bone marrow oedema of the patella, our experience has shown focal synovitis with synovial thickening to be a common associated finding in cases of PFS (Figure 1b).

Figure 5. Sagittal PD-T2 fat-saturated MRI demonstrates inflammatory high signal intensity and oedema of the quadriceps fat pad (black arrow) in a young female patient with PFS. Note is made of patella alta (white arrow), which is likely causing chronic impingement. In many cases of PFS, chronic impingement secondary to patellar malalignment will also be seen. PFS, patellofemoral syndrome.

Deep infrapatellar bursitis

The deep infrapatellar bursa is located inferior to the infrapatellar fat pad in the anterior inferior knee joint between the patellar tendon near its insertion on the tibial tubercle, the tibia and Hoffa's

Figure 6. Sagittal PD-T2 fat-saturated MRI (a) shows a high riding patella with resultant moderate-grade loss of the articular cartilage along the superior patella. Note the underlying patellar bone marrow oedema (white arrow) and high signal intensity of the quadriceps fat pad (black arrow) due to impingement and subsequent inflammation, thickening and oedema. Axial PD-T2 fat-saturated MRI (b) shows lateral subluxation of the patella with trochlear dysplasia (white arrow). There is resultant loss of the articular cartilage of the lateral patellar facet with underlying patellar bone marrow oedema (black arrow), highly suggestive of high-grade chondromalacia.

Figure 7. Axial PD-T2 fat-saturated MRI (a) from a young female with PFS demonstrates patellar subluxation with loss of articular cartilage (black arrow) and focal subarticular bone marrow oedema (white arrow). Sagittal PD-T2 fat-saturated MRI (b) from another patient with PFS shows patella alta with diminution of the patellar articular cartilage and underlying patellar bone marrow oedema (white arrow). PFS, patellofemoral syndrome.

fat pad above. While the deep infrapatellar bursa has been shown to be present on MRI more frequently than previously thought, an enlarged bursa with high T_2 inflammatory signal change should raise the suspicion for bursitis (Figures 4a and 8).

Infrapatellar plica oedema

The ligamentum mucosum is an embryonic synovial membrane which acts to compartmentally separate the knee. While the

Figure 8. Sagittal PD-T2 fat-saturated MRI from a young female with PFS demonstrates patella alta with subsequent high intensity inflammatory changes within the deep infrapatellar bursa, most compatible with bursitis (black arrow). PFS, patellofemoral syndrome.

Figure 9. Sagittal PD-T2 fat-saturated MRI demonstrates high signal intensity from inflammatory changes surrounding the infrapatellar plica (black arrow), indicative of infrapatellar plica syndrome.

majority of these will disappear with age, it is estimated that between 65–90% of knees will retain ligamentum mucosum into adulthood.¹² These remnants are often asymptomatic and will simply appear isointense to the infrapatellar fat pad on T_2 . It has been shown, however, that in patients undergoing repetitive flexion and extension activities, the ligamentum mucosum can become inflamed and thickened – at which point it is called an infrapatellar plica (Figure 9). In our experience, oedema and thickening of an infrapatellar plica is commonly seen in patients with PFS.

COMMON PITFALLS IN THE EVALUATION OF PATELLOFEMORAL SYNDROME

Mistaking Hoffa's fat pad and quadriceps fat pad oedema for volume averaging effects

Very often, oedema is confined to a small region of the fat pads and a study may appear grossly normal, as demonstrated in the sagittal PD-T2 fat-saturated images from a female presenting with anterior knee pain (Figure 10a). However, by scanning the extreme parasagittal planes, small areas of high T_2 intensity within Hoffa's fat pad are often discovered (Figure 10b). Correlation with axial sequences at the same level can confirm that this is in fact Hoffa's fat pad oedema (Figure 10c).

Figure 10. Sagittal PD-T2 MRI (a) from a young female with non-traumatic anterior knee pain show that initial interpretation can appear normal given that just a small amount of oedema is present. By scanning the extreme parasagittal planes (b), subtle oedema can often be appreciated in Hoffa's fat pad (white arrow). This can then be confirmed by correlating with axial imaging at the same level (c, white arrow). Note is also made of trace condylar bone marrow oedema in this case, which is commonly present in cases of PFS secondary to repetitive impaction of the inferior patella. PFS, patellofemoral syndrome.

A small amount of fluid in the infrapatellar bursa is not always to be considered pathological

In our experience, trace amounts of deep infrapatellar bursal fluid will be present in many normal studies. As such, trace

Figure 11. In a case from a young female with PFS, mild synovitis and patellar retinacular strain are not appreciated on the sagittal images (a) but are in fact present on axial T2-PD fat-saturated sequences (b, white arrow). PFS, patellofemoral syndrome.

infrapatellar oedema should be used only as a confirmatory finding when associated with more specific features of PFS, such as: patellar malalignment, impingement and pain of the overlying region on clinical exam.

Missed synovitis and patellar retinacular strain on sagittal reformats alone

We have found that synovitis and patellar retinacular strain are often not well appreciated on sagittal sequences alone and, in many cases, are only visible on axial T2-PD fat-saturated sequences (Figure 11). Search patterns should utilise all reformats to ensure the accurate diagnosis of PFS and its related pathological findings.

IMAGE-GUIDED INTERVENTION FOR PATELLOFEMORAL SYNDROME

By better recognising this constellation of MRI findings found in PFS at our institution, we have been able to expedite diagnosis and ensure that appropriate and targeted therapy is promptly

offered to patients. While many treatments such as physical rehabilitation and orthotics can be offered in cases of PFS, it has been found that these treatments are often inadequate and many patients suffer from recurrent pain.¹¹ By recognising the specific abnormalities associated with PFS, a number of image-guided percutaneous treatment options can be offered. In our practice, ultrasound guided injection of local anaesthetic and corticosteroid into Hoffa's fat pad or the infrapatellar bursa in particular have been highly efficacious in providing sustained pain relief, whereas other groups have described the use of alcohol ablation techniques as a viable treatment option.¹³

CONCLUSIONS

This pictorial essay reviews a number of findings our institution has found to be associated with PFS. By recognising the constellation of MRI findings found in cases of PFS, it can help radiologists expedite diagnosis and ensure that appropriate and targeted therapy is promptly offered to patients. Moreover, it is our hope that this review will initiate systematic reviews of these findings and their association with PFS in the forthcoming literature.

REFERENCES

1. Boling M, Padua D, Marshall S, Guskiewicz K, Pyne S, Beutler A. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. *Scand J Med Sci Sports* 2010; **20**: 725–30. doi: <https://doi.org/10.1111/j.1600-0838.2009.00996.x>
2. van der Heijden RA, Lankhorst NE, van Linschoten R, Bierma-Zeinstra SM, van Middelkoop M. Exercise for treating patellofemoral pain syndrome. *Cochrane Database Syst Rev* 2015; **1**: CD010387. doi: <https://doi.org/10.1002/14651858.CD010387.pub2>
3. Abernethy PJ, Townsend PR, Rose RM, Radin EL. Is chondromalacia patellae a separate clinical entity? *J Bone Joint Surg Br* 1978; **60-B**(2): 205–10. doi: <https://doi.org/10.1302/0301-620X.60B2.659466>
4. Kang S, Park J, Kang SB, Chang CB. MRI findings of young male soldiers with atraumatic anterior knee pain. *Scand J Med Sci Sports* 2016; **26**: 572–8. doi: <https://doi.org/10.1111/sms.12486>
5. van der Heijden RA, de Kanter JL, Bierma-Zeinstra SM, Verhaar JA, van Veldhoven PL, Krestin GP, et al. Structural abnormalities on magnetic resonance imaging in patients with patellofemoral pain: a cross-sectional case-control study. *Am J Sports Med* 2016; **44**: 2339–46. doi: <https://doi.org/10.1177/0363546516646107>
6. Pfirrmann CW, Zanetti M, Romero J, Hodler J. Femoral trochlear dysplasia: MR findings. *Radiology* 2000; **216**: 858–64. doi: <https://doi.org/10.1148/radiology.216.3.r00se38858>
7. Shabshin N, Schweitzer ME, Morrison WB, Parker L. MRI criteria for patella alta and baja. *Skeletal Radiol* 2004; **33**: 445–50. doi: <https://doi.org/10.1007/s00256-004-0794-6>
8. Arendt EA, Dejour D. Patella instability: building bridges across the ocean a historic review. *Knee Surg Sports Traumatol Arthrosc* 2013; **21**: 279–93. doi: <https://doi.org/10.1007/s00167-012-2274-1>
9. Draghi F, Ferrozzi G, Urciuoli L, Bortolotto C, Bianchi S. Hoffa's fat pad abnormalities, knee pain and magnetic resonance imaging in daily practice. *Insights Imaging* 2016; **7**: 373–83. doi: <https://doi.org/10.1007/s13244-016-0483-8>
10. Tuna BK, Semiz-Oysu A, Pekar B, Bukte Y, Hayirlioglu A. The association of patellofemoral joint morphology with chondromalacia patella: a quantitative MRI analysis. *Clin Imaging* 2014; **38**: 495–8. doi: <https://doi.org/10.1016/j.clinimag.2014.01.012>
11. Cole B. J, Malek M. eds. *Articular cartilage lesions*. New York, US: Springer; 2004.
12. Patel SJ, Kaplan PA, Dussault RG, Kahler DM. Anatomy and clinical significance of the horizontal cleft in the infrapatellar fat pad of the knee: MR imaging. *AJR Am J Roentgenol* 1998; **170**: 1551–5. doi: <https://doi.org/10.2214/ajr.170.6.9609172>
13. House CV, Connell DA. Therapeutic ablation of the infrapatellar fat pad under ultrasound guidance: a pilot study. *Clin Radiol* 2007; **62**: 1198–201. doi: <https://doi.org/10.1016/j.crad.2007.07.005>