Was There an Early Habitability Window for Earth's Moon?

Dirk Schulze-Makuch^{1,2} and Ian A. Crawford³

OUR MOON IS uninhabitable and lifeless today. It has no significant atmosphere, no liquid water on its surface, no magnetosphere to protect its surface from solar wind and cosmic radiation, no polymeric chemistry, and it is subject to large diurnal temperature variations (*e.g.*, Vaniman *et al.*, 1991; Schulze-Makuch and Irwin, 2008). Thus, associating our Moon with habitability seems outrageous, and certainly it would have been just a decade ago. However, results from recent space missions, as well as sensitive analyses of lunar rock and soil samples, have indicated that the Moon is not as dry as previously thought (*e.g.*, Anand, 2010; Hauri *et al.*, 2017). In addition to the probable occurrence of water ice in permanently shadowed polar craters (*e.g.*, Feldman *et al.*, 1998; Baker *et al.*, 2005; Lawrence, 2017), spectroscopic studies also indicate the presence of hydrated surface materials at high, but not permanently shadowed, latitudes (Clark, 2009; Pieters *et al.*, 2009; Li and Milliken, 2017), with evidence for temporal variations over the course of a lunar day (Sunshine *et al.*, 2009). In addition, recent studies of the products of lunar volcanism indicate that the lunar interior also contains more water than was once appreciated and that the lunar mantle may even be as comparably water-rich as Earth's upper mantle (see Hauri *et al.*, 2017, for a review).

The existence of indigenous sources of water implies that the Moon may not always have been as dead and dry as it is today. Insofar as water is required for habitability (*e.g.*, Kasting *et al.*, 1993; although it is not the sole criterion, see Schulze-Makuch *et al.*, 2011), we can speculatively identify two possible windows for lunar habitability. These may have occurred immediately following the accretion of the Moon and some hundreds of millions of years later following outgassing associated with lunar volcanic activity.

Current understanding is that the Moon originated from a gigantic impact 4.5 billion years ago (*e.g.*, Stevenson and Halliday, 2014). The extent to which volatiles were preserved in the Moon-forming debris disk produced by this impact is model-dependent, but impediments to the diffusion of water molecules in a silicate-dominated vapor are expected to result in some water retention in the disk and therefore in the newly formed Moon (Nakajima and Stevenson, 2014; Hauri *et al.*, 2017). The evidence for water concentrations of several hundred parts per million in the

mantle source regions of lunar basalts (Hauri *et al.*, 2017; Lin *et al.*, 2017) indicates either that volatiles were indeed preserved during the formation of the Moon or that they were added shortly afterwards by impacting asteroids (*e.g.*, Barnes *et al.*, 2016).

Following accretion, the Moon is expected to have been largely molten, with its silicate components existing in the form of a lunar magma ocean (LMO). Such magma oceans are expected to outgas volatiles, leading to the formation of significant transient atmospheres (Elkins-Tanton, 2008). Indeed, Lin *et al.* (2017) have invoked degassing from the LMO to reconcile the relatively low abundance of water in the post-LMO lunar mantle (at most a few hundred parts per million) with their predictions of much higher values (possibly >1000 ppm) prior to LMO crystallization. On the other hand, some authors have argued that the LMO would have been initially dry following accretion, with the current mantle volatile budget having been added by a subsequent "late veneer" of asteroidal volatiles (*e.g.*, Barnes *et al.*, 2016; Hauri *et al.*, 2017). However, in either case, it appears that significant quantities of water were present in the final stages of LMO evolution. Here, we merely note that outgassing 500 ppm water during the LMO phase (which would be required to bring the higher original values predicted by Lin *et al.* [2017] into agreement with current estimates) could in principle result in a surface water layer of an order of 1 km thickness. Of course, this would be a very optimistic estimate for the depth of any early lunar oceans—water would only be stable at the surface if protected by a sufficiently dense atmosphere, and significant losses would be expected owing to impact erosion (*e.g.*, Melosh and Vickery, 1989)—but it illustrates how much water might potentially have been available.

Needham and Kring (2017) have suggested a second phase of outgassing, and associated peak in lunar atmospheric pressure, as a result of mare basalt eruptions ~3.5 billion years ago. Gases derived from lava outpourings may have built up an atmosphere of about 10 mbar, which is above the triple-point pressure of water and about 1.5 times the present atmospheric pressure on Mars (and about 3 times as massive as the current martian atmosphere, given the difference in surface gravities). For comparison with the discussion above, Needham and Kring's estimated outgassing of water (~10¹⁴ kg) would equate to a global layer having an average depth of ~3 mm.

¹Center for Astronomy and Astrophysics, Technical University Berlin, Berlin, Germany.

²School of the Environment, Washington State University, Pullman, Washington, USA.

³Department of Earth and Planetary Sciences, Birkbeck College, University of London, London, UK.

The duration of hypothetical lunar atmospheres was first studied by Vondrak (1974) and reviewed by Stern (1999). For the dense atmospheres considered here (with total masses $>10^8$ kg), the mass loss rate from a collisionally dominated atmosphere is expected to be $\sim 10 \text{ kg s}^{-1}$, independent of atmospheric mass. The loss rate would be higher if impact erosion (Melosh and Vickery, 1989) or solar-wind stripping were important, but note that the early Moon may have been protected by a magnetic field (e.g., Hood, 2011), reducing the impact of the latter process. Based on these loss rates, Needham and Kring (2017) estimated a lifetime of ~ 70 million years for the transient atmosphere generated by lunar volcanic activity, and the duration of a hypothetical denser and earlier atmosphere immediately following LMO crystallization could have been even longer. In principle, liquid water could have existed on the lunar surface during these times, and perhaps even more likely in protected subsurface environments such as interstitial pore spaces within the impact-generated mega-regolith.

It is instructive to put this time frame into perspective from a natural history point of view: Lazcano and Miller (1994) speculated that the time it took from the formation of suitable building blocks of life to the first cyanobacteria was no more than 10 million years. If they are correct, the transition from a nonliving to a living system might have taken place in considerably less of a time span, perhaps as little as a few thousand years. On the other hand, Orgel (1998) has argued that we simply do not understand how life, particularly its replication system, originated on primitive Earth; thus it is not possible to come up with a reliable time estimate. If abiogenesis is able to occur on short timescales, then an origin of life on the Moon cannot be excluded. Moreover, there is an alternative scenario for introducing life to the Moon: the early history of the Solar System was dominated by giant impacts and the transfer of meteorites between planets (e.g., Mileikowsky *et al.*, 2000). During this time, and indeed subsequently, it is expected that meteorites blasted off the surface of Earth will have landed on the Moon (Armstrong *et al.*, 2002; Gutiérrez, 2002; Schulze-Makuch, 2013), and some of them are expected to have survived the impact (e.g., Crawford *et al.*, 2008). As life appears to have been present on Earth by 3.8–3.5 billion years ago (e.g., Schopf, 1993; Schidlowski, 2001; Brasier *et al.*, 2015; Nutman *et al.*, 2016; Hassenkam *et al.*, 2017; Schopf *et al.*, 2018), and possibly by 4.1 or even 4.28 billion years ago (e.g., Bell *et al.*, 2015; Dodd *et al.*, 2017), it is possible that Earth life could have inoculated transiently habitable lunar environments. We note that the chances of survival of microorganisms within terrestrial meteorites impacting the Moon would be increased by the presence of even a tenuous lunar atmosphere because this would reduce the impact velocity.

Of course, habitability requires much more than just the presence of a significant atmosphere and liquid water. Other constraints that need to be satisfied have been elaborated elsewhere (e.g., Schulze-Makuch *et al.*, 2011; Cockell *et al.*, 2016). We do not know whether there were any intrinsic organic compounds on the Moon at that time, but even if not, these would likely have been delivered from Earth, carbonaceous asteroids, and perhaps other sources, via meteorite impacts (e.g., Pierazzo and Chyba, 1999; Crawford *et al.*, 2008; Burchell *et al.*, 2010; Matthewman *et al.*, 2015; Svetsov and Shuvalov, 2015). Thus, sources of organics on the lunar surface may have been available. Moreover, as the early Moon appears to have had a magnetic field (e.g.,

Hood, 2011), its surface may have been partially protected from solar and cosmic radiation.

If these early habitable environments ever existed, would there be any evidence remaining? Clearly, we do not see the familiar water-modified topography on the Moon that we see on Mars (e.g., fluvial channels or crater rims that are altered by liquid water), and it is questionable whether any topographical evidence of early surface water would be preserved on the Moon after approximately 4 billion years of pounding by solar wind, cosmic radiation, and micrometeorites. On the other hand, there is some (albeit limited) evidence for oxidation and/or hydrothermal activity in lunar rocks (e.g., Shearer *et al.*, 2014; Joy *et al.*, 2015). One especially promising future line of inquiry, especially for evidence of near-surface water during the high point of mare volcanism, would be to search for evidence of hydration in paleoregolith layers trapped between lava flows dating from this time (see Crawford *et al.*, 2010, for a discussion of lunar paleoregolith deposits). It is also interesting to note that Cannon *et al.* (2017) have proposed an early episode of hydration, resulting from magma ocean degassing, as the source of primordial clay mineral formation on Mars, and if such a process operated on early Mars, then it seems likely to have operated on the early Moon as well. Although it seems certain that if liquid water ever existed on the early Moon it would have been much less prevalent than on early Mars, evidence for it may yet turn up in studies of lunar samples.

Thus, if liquid water and a significant atmosphere were present on the early Moon for millions of years, it can be assumed that the lunar surface was at least transiently habitable and probably also had an inventory of the building blocks required for life. Whether life ever arose on the Moon, or was transported to it from elsewhere, is of course highly speculative and can only be addressed by an aggressive future program of lunar exploration (e.g., Crawford and Joy, 2014). An important aspect of such an exploration program would be obtaining samples from paleoregolith deposits dating from time of peak mare volcanism (i.e., at ~ 3.5 Ga) to determine if hydrated conditions (or other evidence for habitable conditions, including possible biomarkers) existed at that time.

In addition, experiments could be conducted in lunar environment simulation chambers in laboratories on Earth to observe whether microorganisms can maintain viability under the environmental conditions predicted to have existed on the early Moon. Indeed, the surface conditions predicted by Needham and Kring (2017) are not very different from those routinely produced in Mars simulation chambers (Galletta *et al.*, 2006; Jensen *et al.*, 2008; de Vera *et al.*, 2013). Such facilities are available at many research institutions around the world and could easily be modified to simulate early lunar environmental conditions. Moon-analog experiments are already done on the International Space Station (ISS) as part of the BIOMEX project (Baqué *et al.*, 2017), and methods to analyze these samples for microbial viability are being developed (Liu *et al.*, 2018). Thus, we recommend utilizing both Moon simulation chambers on our planet and on the ISS to test whether there might have been an early habitability window on the Moon.

Acknowledgment

Dirk Schulze-Makuch acknowledges support from ERC Advanced Grant “HOME” (No. 339 231).

References

- Anand, M. (2010) Lunar water: a brief review. *Earth Moon Planets* 107:65–73.
- Armstrong, J., Wells, L., and Gonzalez, G. (2002) Rummaging through Earth's attic for remains of ancient life. *Icarus* 160: 183–196.
- Baker, V.R., Dohm, J.M., Fairén, A.G., Ferre, P.A., Ferris, J.C., Miyamoto, H., and Schulze-Makuch, D. (2005) Extraterrestrial hydrogeology. *Hydrogeol J* 13:51–68.
- Baqué, M., Böttger, U., Leya, T., and de Vera, J.-P.P. (2017) BIOMEX on EXPOSE-R2: first results on the preservation of Raman biosignatures after space exposure. In *19th EGU General Assembly*, European Geosciences Union, Munich, Germany, p 3697.
- Barnes, J.J., Tartèse, R., Anand, M., McCubbin, F.M., Neal, C.R., and Franchi, I.A. (2016) Early degassing of lunar urKREEP by crust-breaching impact(s). *Earth Planet Sci Lett* 447:84–94.
- Bell, E.A., Boehnke, P., Harrison, T.M., and Mao, W.L. (2015) Potentially biogenic carbon preserved in a 4.1 billion-year-old zircon. *Proc Natl Acad Sci USA* 112:14518–14521.
- Brasier, M.D., Antcliffe, J., Saunders, M., and Wacey, D. (2015) Changing the picture of Earth's earliest fossils (3.5–1.9 Ga) with new approaches and new discoveries. *Proc Natl Acad Sci USA* 112:4859–4864.
- Burchell, M.J., Parnell, J., Bowden, S.A., and Crawford, I.A. (2010) Hypervelocity impact experiments in the laboratory relating to lunar astrobiology. *Earth Moon Planets* 107:55–64.
- Cannon, K.M., Parman, S.W., and Mustard, J.F. (2017) Primordial clays on Mars formed beneath a steam or supercritical atmosphere. *Nature* 552:88–91.
- Clark, R.N. (2009) Detection of adsorbed water and hydroxyl on the Moon. *Science* 326:562–564.
- Cockell, C.S., Bush, T., Bryce, C., Direito, S., Fox-Powell, M., Harrison, J.P., Lammer, H., Landenmark, H., Martin-Torres, J., Nicholson, N., Noack, L., O'Malley-James, J., Payler, S.J., Rushby, A., Samuels, T., Schwendner, P., Wadsworth, J., and Zorzano, M.P. (2016) Habitability: a review. *Astrobiology* 16: 89–117.
- Crawford, I.A. and Joy, K.H. (2014) Lunar exploration: opening a window into the history and evolution of the inner Solar System. *Philos Trans A Math Phys Eng Sci* 372, doi:10.1098/rsta.2013.0315.
- Crawford, I.A., Baldwin, E.C., Taylor, E.A., Bailey, J.A., and Tsembeles, K. (2008) On the survivability and detectability of terrestrial meteorites on the Moon. *Astrobiology* 8:242–252.
- Crawford, I.A., Fagents, S.A., Joy, K.H., and Rumpf, M.E. (2010) Lunar palaeoregolith deposits as recorders of the galactic environment of the Solar System and implications for astrobiology. *Earth Moon Planets* 107:75–85.
- de Vera, J.-P., Schulze-Makuch, D., Khan, A., Lorek, A., Koncz, A., Ott, S., Möhlmann, D., and Spohn, T. (2013) Adaptation of the lichen *Pleopsidium chlorophanum* to martian surface conditions within 35 days. *Planet Space Sci* 198:182–190.
- Dodd, M.S., Papineau, D., Grenne, T., Slack, J.F., Rittner, M., Pirajno, F., O'Neil, J., and Crispin, T.S.L. (2017) Evidence for early life on Earth's oldest hydrothermal vent precipitates. *Nature* 543:60–64.
- Elkins-Tanton, L.T. (2008) Linked magma ocean solidification and atmosphere growth for Earth and Mars. *Earth Planet Sci Lett* 271:181–191.
- Feldman, W.C., Maurice, S., Binder, A.B., Barraclough, B.I., Elphic, R.C., and Lawrence, D.J. (1998) Fluxes of fast and epithermal neutrons from Lunar Prospector: evidence for water ice at the lunar poles. *Science* 281:1496–1500.
- Galletta, G., Ferri, F., Fanti, G., D'Alessandro, M., Bertoloni, G., Pavarin, D., Bettanini, C., Cozza, P., Pretto, P., and Bianchini, G. (2006) S.A.M., the Italian martian simulation chamber. *Orig Life Evol Biosph* 36:625–627.
- Gutiérrez, J.L. (2002) Terrene meteorites in the Moon: their relevance for the study of the origin of life in the Earth. In *Proceedings of the First European Workshop on Exo-Astrobiology*, ESA SP-518, edited by H. Lacoste, ESA Publications Division, Noordwijk, the Netherlands, pp 187–191.
- Hassenkam, T., Anderson, M.P., Dalby, K.N., Mackenzie, D.M.A., and Rosing, M.T. (2017) Elements of Eoarchean life trapped in mineral inclusions. *Nature* 548:78–81.
- Hauri, E.H., Saal, A.E., Nakajima, M., Anand, M., Rutherford, M.J., Van Orman, J.A., and LeVoyer, M. (2017) Origin and evolution of water in the Moon's interior. *Annu Rev Earth Planet Sci* 45:89–111.
- Hood, L.L. (2011) Central magnetic anomalies of Nectarian-aged lunar impact basins: probable evidence for an early core dynamo. *Icarus* 211:1109–1128.
- Jensen, L.L., Merrison, J., Hansen, A.A., Mikkelsen, K.A., Kristoffersen, T., Nørnberg, P., Lomstein, B.A., and Finster, K. (2008) A facility for long-term Mars simulation experiments: the Mars Environmental Simulation Chamber (MESCH). *Astrobiology* 8:537–548.
- Joy, K.H., Visscher, C., Zolensky, M.E., Mikouchi, T., Hagiya, K., Ohsumi, K., and Kring, D.A. (2015) Identification of magnetite in lunar regolith breccia 60016: evidence for oxidized conditions at the lunar surface. *Meteorit Planet Sci* 50: 1157–1172.
- Kasting, J.F., Whitmire, D.P., and Reynolds, R.T. (1993) Habitable zones around main sequence stars. *Icarus* 101:108–128.
- Lawrence, D.J. (2017) A tale of two poles: toward understanding the presence, distribution, and origin of volatiles at the polar regions of the Moon and Mercury. *J Geophys Res Planets* 122:21–52.
- Lazcano, A. and Miller, S.L. (1994) How long did it take for life to begin and evolve to cyanobacteria. *J Mol Evol* 39:549–554.
- Li, S. and Milliken, R.E. (2017) Water on the surface of the Moon as seen by the Moon Mineralogy Mapper: distribution, abundance, and origins. *Sci Adv* 3, doi:10.1126/sciadv.1701471.
- Lin, Y., Tronche, E.J., Streenstra, E.S., and van Westrenen, W. (2017) Evidence for an early wet Moon from experimental crystallization of the lunar magma ocean. *Nat Geosci* 10:14–19.
- Liu, Y., Schulze-Makuch, D., de Vera, J.-P., Cockell, C., Leya, T., Baqué, M., and Walther-Antonio, M. (2018) Development of effective bacterial single cell lysis method suitable for whole genome amplification in microfluidic platforms. *Micro-machines*, in press.
- Matthewman, R., Court, R.W., Crawford, I.A., Jones, A.P., Joy, K.H., and Sephton, M.A. (2015) The Moon as a recorder of organic evolution in the early Solar System: a lunar regolith analog study. *Astrobiology* 15:154–168.
- Melosh, H.J. and Vickery, M. (1989) Impact erosion of the primordial atmosphere of Mars. *Nature* 338:487–489.
- Mileikowsky, C., Cucinotta, F.A., Wilson, J.W., Gladman, B., Horneck, G., Lindegren, L., Melosh, J., Rickman, H., Valtonen, M., and Zheng, J.Q. (2000) Natural transfer of viable microbes in space. 1. From Mars to Earth and Earth to Mars. *Icarus* 145:391–427.
- Nakajima, M. and Stevenson, D.J. (2014) Investigation of the initial state of the Moon-forming disk: bridging SPH simulations and hydrostatic models. *Icarus* 233:259–267.

- Needham, D.H. and Kring, D.A. (2017) Lunar volcanism produced a transient atmosphere around the ancient Moon. *Earth Planet Sci Lett* 478:175–178.
- Nutman, A.P., Bennett, V.C., Friend, C.R.L., Van Kranendonk, M.J., and Chivas, A.R. (2016) Rapid emergence of life shown by discovery of 3700-million-year-old microbial structures. *Nature* 537:535–538.
- Orgel, L.E. (1998) The origin of life—how long did it take? *Orig Life Evol Biosph* 28:91–96.
- Pierazzo, E. and Chyba, C.F. (1999) Amino acid survival in large cometary impacts. *Meteorit Planet Sci* 34:909–918.
- Pieters, C.M., Goswami, J.N., Clark, R.N., Annadurai, M., and Boardman, J. (2009) Character and spatial distribution of OH/H₂O on the surface of the Moon seen by M3 on Chandrayaan-1. *Science* 326:568–572.
- Schidlowski, M. (2001) Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of Earth history: evolution of a concept. *Precambrian Res* 106:117–134.
- Schopf, J.W. (1993) Microfossils of the early Archean Apex Chert: new evidence of the antiquity of life. *Science* 260:640–646.
- Schopf, J.W., Kitajima, K., Spicuzza, M.J., Kudryavtsev, A.B., and Valley, J.W. (2018) SIMS analysis of the oldest known assemblage of microfossils document their taxon-correlated carbon isotope compositions. *Proc Natl Acad Sci USA* 115:53–58.
- Schulze-Makuch, D. (2013) Organic molecules in lunar ice: a window to the early evolution of life on Earth. In *Habitability on other Planets and Satellites*, edited by J. Seckbach and J.-P. de Vera, Springer, Dordrecht, the Netherlands.
- Schulze-Makuch, D. and Irwin, L.N. (2008) *Life in the Universe: Expectations and Constraints*, 2nd ed., Springer, Berlin.
- Schulze-Makuch, D., Méndez, A., Fairén, A.G., von Paris, P., Turse, C., Boyer, G., Davila, A.F., António, M.R.S., Catling, D., and Irwin, L.N. (2011) A two-tiered approach to assessing the habitability of exoplanets. *Astrobiology* 11:1041–1052.
- Shearer, C.K., Sharp, Z.D., Burger, P.V., McCubbin, F.M., Provencio, P.P., Brearley, A.J., and Steele, A. (2014) Chlorine distribution and its isotopic composition in “rusty rock” 66095: implications for volatile element enrichments of “rusty rock” and lunar soils, origin of “rusty” alteration, and volatile element behavior on the Moon. *Geochim Cosmochim Acta* 139:411–433.
- Stern, S.A. (1999) The lunar atmosphere: history, status, current problems, and context. *Rev Geophys* 37:453–491.
- Stevenson, D.J. and Halliday, A. (2014) The origin of the Moon. *Philos Trans A Math Phys Eng Sci* 372, doi:10.1098/rsta.2014.0289.
- Sunshine, J.M., Farnham, T.L., Feaga, L.M., Groussin, O., Merlin, F., Milliken, R.E., and A’Hearn, M.F. (2009) Temporal and spatial variability of lunar hydration as observed by the Deep Impact spacecraft. *Science* 326:565–568.
- Svetsov, V.V. and Shuvalov, V.V. (2015) Water delivery to the Moon by asteroidal and cometary impacts. *Planet Space Sci* 117:444–452.
- Vaniman, D., Reedy, R., Heiken, G., Olhoeft, G., and Mendell, W. (1991) The lunar environment. In *The Lunar Sourcebook: A User’s Guide to the Moon*, edited by G.H. Heiken, D. Vaniman, and B.M. French, Cambridge University Press, Cambridge, UK, pp 27–60.
- Vondrak, R.R. (1974) Creation of an artificial lunar atmosphere. *Nature* 248:657–659.

Address correspondence to:

Dirk Schulze-Makuch
Zentrum für Astronomie und Astrophysik
Technische Universität Berlin
Hardenbergstr 36
Berlin 10623
Germany

E-mail: schulze-makuch@tu-berlin.de

Submitted 21 February 2018

Accepted 20 June 2018

Associate Editor: Christopher McKay

Abbreviations Used

ISS = International Space Station
LMO = lunar magma ocean