(~3rd hospitalization day onwards) in-hospital food-intake stabilizes. Multivariate logistic regression models showed that eating half or less of the served meals at the first 48 hours was associated with higher odds of cognitive decline (OR=1.70, 95%CI=1.13–2.55), whereas low intake after 48 hours of admission was associated with higher odds of ADL decline (OR=1.48, 95%CI=1.01–2.19), controlling for baseline functional, clinical, and nutritional status. These associations between low food-intake at various time points and worse outcomes shows the importance of routine food-intake assessment along the hospitalization sequelae, with emphasis on both the first 48 hours of admission and at least one consecutive day. Future studies should investigate whether the performance of consecutive food-intake assessments can improve food consumption and outcomes.

FRAILITY INDICES AND THE CLINICAL FRAILITY SCALE EACH PREDICT 30-DAY MORTALITY IN ED PATIENTS

K. Rockwood¹, A. van der Valk², R. Eendebak¹, K. Lines¹, S. Howlett³, O. Theou¹, 1. *Dalhousie University*, 2. *Nova Scotia Health Authority*, 3. *Department of Pharmacology, Dalhousie University*

We examined the ability of frailty indices and the Clinical Frailty Scale to predict mortality in acutely ill, older (aged 65+ years) Emergency Department (ED) patients referred to internal medicine. Here we report on the first 415 patients (Mage = 80.6 ± 8.3, 58.1% women) with complete data. Acuity was assessed using the Canadian Triage and Acuity Scale (CTAS). A geriatrician (KR) conducted a Comprehensive Geriatric Assessment from which an FI-score was calculated based on the patient's current state (FI-CGA). The Clinical Frailty Scale (CFS) was also completed. An FI-Lab was constructed using 30 common laboratory tests collected within 2 days of CGA administration. Of the 415 patients, 321 (77.3%) were admitted, staying an average of 26.9 ± 45.8 days. Mortality within 30 days was 2.6% (CFS), 8% (FI-CGA) and 7.9% (FI-Lab) for the least frail groups and 41.8% (CFS), 28.2% (FI-CGA), and 29.7% (FI-Lab) for the frailest groups. After adjusting for age, sex, and CTAS, all three frailty tools independently predicted 6-month mortality. The hazard ratio was 1.67 (95% CI: 1.48–1.89) per 1-grade increase in CFS, 1.04 (1.03–1.06) per 0.01-point increase in the FI-CGA and 1.05 (1.02–1.07) per 0.01-point increase in the FI-Lab. ED patients requiring CTAS-defined urgent assistance were at higher risk for mortality than less urgent cases; 2.65 (1.71–4.12) CFS model, 2.84 (1.80–4.48) FI-CGA model, 1.81 (1.08–3.02) FI-Lab model. An FI from common laboratory tests is a promising tool for grading frailty and predicting mortality for ED patients. It can be automated readily and adds prognostic information to the CFS.

HOME HEALTH SERVICES: PREVENTING ADVERSE OUTCOMES AFTER TOTAL HIP REPLACEMENT

M. Meucci¹, P. Gozalo², D. Dosa², S. Allen³, 1. *University of Rhode Island*, 2. *Brown University*, 3. *Department of Health Services, Policy and Practice, Brown University*

Background and Significance: The number of total hip replacement (THR) surgeries performed has increased dramatically in recent years from 138,700 in 2000 to

310,800 in 2010 with nearly 50% of patients being discharged directly to home. However, the rate of 30 day hospital readmission after THR has recently surpassed 10%. Although it is standard practice for home health services (HHS) to be prescribed after THR, no previous study has examined the impact of home-based therapy and nursing services on reducing the likelihood of rehospitalization. **Objective:** To determine whether home health services reduce the likelihood of 30 day rehospitalization post-THA among Medicare beneficiaries. **Methods:** Using Medicare claims data from 2007–2013 we identified all beneficiaries who underwent THR and were discharged directly home from the hospital with and without HHS. We used a logistic mixed effects model to estimate the impact of HHS on rehospitalization rates and 30 days after discharge for THA while adjusting for potential confounders and accounting for geographic variation in practice. **Findings:** Of the 410,162 individuals discharged directly home post-THA, 68.2% had a home health visit in the next 30 days. Among those with and without HHS, 4.2% and 6.9% were rehospitalized, respectively (χ^2 P-value <0.0001). After adjusting for confounding and regional variations, the likelihood of being rehospitalized among those without HHS was twice that (OR 2.03; 95% CI 1.97–2.10) of those with HHS. **Conclusions:** Among Medicare beneficiaries discharged directly home after THR home health services help to prevent costly rehospitalizations during the first 30 days.

HOW HOSPITALIZED OLDER ADULTS SHARE CONCERNS DURING DAILY ROUNDS

K. Pecanac, E. LeSage, E. Stephens, *University of Wisconsin-Madison*

In order to deliver patient-centered care, providers need to elicit and address patient concerns during a hospital stay. Patients have the opportunity to share concerns with providers daily during morning rounds. The purpose of this analysis was to determine how patients raise their concerns during daily rounds with providers. Eleven conversations between providers and older adult patients (age 65 or older) during daily rounds were audio recorded. Data collection occurred in the ICU and on a medical floor of one hospital. Data were analyzed using conversation analysis, a qualitative method, which focuses on the sequence of actions of utterances (not just what is said but how and when it is said). Preliminary analysis shows that, as expected, patients often communicated concerns after providers invited patients to ask questions or share how they were feeling. However, at times patients also took the opportunity to reorient the conversation from the physician's agenda or concerns to the patient's own concerns. Patient concerns included physiological issues such as distressing symptoms not yet discussed, but they often involved concerns about when they would go home or if it was expected that they would not return home, where they would go. These findings show that patients are able to communicate their concerns to providers, even when not elicited. Nevertheless, providers may want to address the developing plan for the patient's discharge daily to demonstrate that they recognize it is a common concern and to help patients understand why uncertainty regarding discharge may exist in their situation.