were the most important issues identified in caring for these patients, with ambivalence about possible dangers of strong pain medication and utility of feeding tubes at EOL. Top priorities identified for improving PC/EOL care included: greater family involvement (50%); education/training on pain control (48%) and management of other symptoms (33%); and use of a PC team (34%). Findings provide guidance on how best to target PC education/support.

ASSESSING TECHNICAL FEASIBILITY AND ACCEPTABILITY OF PROVIDING TELEHEALTH PALLIATIVE CARE IN NURSING HOMES

V. Wertz¹, N. Bui, MS², E. Uy, MS³, P. Barrientos, MS⁴, D. David, PhD⁵, S.J. Lee, MD⁶, C. Ritchie, MD⁷, C. Stephens, PhD⁸, 1. UCSF School of Nursing, Dept. of Community Health Systems, San Francisco, California, United States, 2. Asian Health Services, Oakland, CA, USA; UCSF School of Nursing, Dept. of Community Health Systems, San Francisco, CA, USA, 3. UCSF School of Nursing, San Francisco, CA, USA, 4. UCSF School of Nursing, Dept. of Community Health Systems, San Francisco, CA, USA, 5. San Francisco VA Medical Center, San Francisco, CA, USA; UCSF School of Nursing, Dept. of Community Health Systems, San Francisco, CA, USA, 6. San Francisco VA Medical Center, San Francisco, CA, USA; UCSF School of Medicine, Division of Geriatrics, San Francisco, CA, USA, 7. UCSF School of Medicine, Division of Geriatrics, San Francisco, CA, USA, 8. UCSF School of Nursing, Dept. of Community Health Systems, San Francisco, CA, USA

Use of telehealth to provide NH palliative care(PC) has been extremely limited, only focusing on end-of-life care. No studies have examined how it can be used to proactively deliver PC expertise/support to NH residents, families, and staff. We conducted 6 feasibility tests of telehealth PC in 3 NHs with 19 residents, families, providers and staff (53% female, 47% white, 79% age 50+). None had prior experience with telehealth. 100% were comfortable with the video visit and could see themselves using it in the near future (74% for own care; 90% for family care). 84% strongly believed technology could improve communication between patients/families/providers; 74% strongly believed technology could improve care coordination between NHs and hospitals. 37% rated audio quality fair, 63% good/very good. 47% rated visual quality fair, 47% good/very good. NH telehealth PC is technically feasible and acceptable, providing an innovative opportunity to improve access to needed NH PC services/supports.

SESSION 2440 (SYMPOSIUM)

SUBJECTIVE COGNITION, AFFECT, & PERSONALITY: IMPLICATIONS FOR COGNITIVE SELF-REPORT

Chair: N. Hill, *The Pennsylvania State University, University Park, Pennsylvania*

Evidence suggests that cognitive self-report is a potentially valuable indicator of preclinical Alzheimer's disease. However, subjective cognition is also associated with affective symptoms and certain personality traits, which can confound its utility as a predictor of cognitive decline. Furthermore,

current measures of subjective cognition are highly variable across studies and tap into a variety of different constructs. The purpose of this symposium is to present findings from multiple studies that examined aspects of subjective cognition, its association with psychological well-being, the influence of individual characteristics (i.e., personality and sex) on cognitive self-report, and sources of response bias in current approaches to assessment. First, we present the results of a daily diary study that examined the influence of memory lapses on positive and negative affect. Second, we describe the temporal associations between memory self-report and depressive symptoms over an 18 year period. Our third presentation examines the influence of neuroticism on memory self-report in women specifically. Fourth, we present the results of a study that investigated the influence of temperament and personality on subjective executive function among older adults. And finally, we discuss sources of response bias in subjective cognition measures with a report on interviews conducted with older adults to elucidate their interpretations of, and responses to, common cognitive self-report items.

SOURCES OF RESPONSE BIAS IN COGNITIVE SELF-REPORT ITEMS: "WHICH MEMORY ARE YOU TALKING ABOUT?"

N. Hill¹, J. Mogle, PhD², E. Whitaker, MS³, A. Gilmore-Bykovskyi, PhD⁴, S. Bhargava, PhD⁵, I. Bhang, MS⁶, L. Sweeder, BSN⁷, K. Van Haitsma, PhD⁸, 1. *The Pennsylvania State University, University Park, Pennsylvania, United States*, 2. *The Pennsylvania State University, University Park, PA, USA*, 3. *The Pennsylvania State University, University Park, PA, USA*, 4. *University of Wisconsin-Madison, Madison, Wisconsin, USA*, 5. *The Pennsylvania State University, University Park, PA, USA*, 6. *The Pennsylvania State University, University Park, PA, USA*, 7. *The Pennsylvania State University, University Park, PA, USA*, 8. *The Pennsylvania State University, University Park, PA, USA*

Understanding how older adults interpret and respond to questions about cognition is critical to identifying response biases that may limit the accuracy of cognitive self-reports in identifying risk for cognitive decline. In this study, we used cognitive interviewing to identify features of cognitive self-report item content and structure that contribute to: 1) differing interpretations among respondents, and 2) older adults' decisional processes when responding. Older adults without dementia (n=49; Mage = 74.5 years; 36.7% male) completed an audio-recorded cognitive interview during which common cognitive self-report items were evaluated; the Question Appraisal System was used to analyze the resultant transcripts. The most common sources of response bias were vague item wording leading to multiple interpretations of meaning, incorrect assumptions regarding consistency of problems across situations, and provocation of an emotional reaction that influenced responses. Findings from this study identify modifiable sources of response bias to inform refinement of cognitive self-report measures.

DAILY MEMORY LAPSES AND PSYCHOLOGICAL WELL-BEING

J. Mogle¹, N. Hill, PhD², D. Almeida, PhD³, 1. *Penn State University, University Park, Pennsylvania, United States*,