AWARENESS AND ANXIETY IN HELP-SEEKERS FOR DEMENTIA ASSESSMENT SERVICE

D. Leung, G. Wong, *The University of Hong Kong*

Anxiety is common in people with dementia, and is associated with awareness of cognitive deficits. Awareness is a multicomponent construct but little is known about how other components, such as functional ability, influences anxiety. Few studies looked at awareness and anxiety in people seeking help from dementia assessment service. This study explores the potential relationship between awareness and anxiety in help-seekers with suspected dementia. The study included 110 dyads of caregivers and people with suspected dementia from a dementia early detection service in Hong Kong. Awareness of cognitive performance was estimated by comparing participants' Mini-Mental State Examination (MMSE) and self-rated memory on the Quality of Life-Alzheimer's Disease (QoL-AD). Awareness of daily activities functioning was estimated by comparing caregivers-rated Alzheimer's Disease Cooperative Study-Activities of Daily Living (ADCS-ADL) and participants' self-rated ability to do chores on QoL-AD. Anxiety symptoms were assessed using the relevant item on Neuropsychiatric Inventory-Questionnaire. Anxiety was reported in 31.8% of the participants. Self-reported memory and ability to do chores were correlated with MMSE ($r=0.21$, $p=.033$) and ADCS-ADL ($r=0.76$, $p<.01$) respectively. Regression analysis showed that age, gender, education, and awareness of cognitive performance and daily activity functioning explained a significant proportion of the variance in anxiety ($R^2=0.13$, $F(5, 103)=1.73$, $p=.016$). Greater discrepancy between self-reported and caregiver-reported ability to do chores predicted anxiety ($\beta=0.45$, $p=.045$). Awareness of cognitive and functional deficits appears intact in this sample of people with suspected dementia. Anxiety in this population may be predicted by a lower awareness of daily activities functioning ability.

CHARACTERISTICS OF RETENTION IN BEHAVIORAL HEALTH SERVICES IN PRIMARY CARE

L. Schneider¹, J. Dieker², K. Johanson¹, C. Yeager¹, L. Feliciano², 1. *University of Colorado at Colorado Springs*, 2. *University of Colorado Colorado Springs*

One role of behavioral health clinicians in primary care is to engage patients in brief solution-focused therapy for behavioral health problems identified during medical visits. While some patients choose not to return to sessions following these visits, other patients return to behavioral health services for multiple sessions. The purpose of the present research was to determine the demographic and psychosocial traits of patients over the age of 50 who choose to follow up with behavioral health services following initial contact with a clinician in a primary care setting. Archival data was analyzed from a system of federally qualified health centers in Colorado. A sample of patients above the age of 50 who returned for additional behavioral health services beyond one visit was identified ($N = 1751$). On average, patients attended 5 ($SD = 5.20$) behavioral health sessions. The mean age of patients was 60.58 years ($SD = 7.62$). The majority were Caucasian (66.3%), followed by Hispanic/Latino (13.4%), Black/African American (8.9%), Native American/American

Indian (1.3%), and other/unknown (10.1%). More than half of patients who returned for services were opioid users (54.9%). A relatively small number were homeless (6.4%). Patients were single (38.4%), followed by divorced (23.9%), married (26.8%), and widowed (8.3%). Findings suggest opioid use is related to retention in behavioral health services. This information can be used to identify patients less likely to attend follow up sessions and work to increase engagement in behavioral health services.

COMPREHENSIVE GERIATRIC ASSESSMENTS IN INTEGRATED CARE PROGRAMS: A SYSTEMATIC LITERATURE REVIEW

A. Stoop¹, M. Lette², P. van Gils¹, G. Nijpels², C. Baan¹, S. de Bruin¹, 1. *National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands*, 2. *Amsterdam Public Health research institute, VU University Medical Center, Amsterdam, the Netherlands*

Introduction: Integrated care for older adults living at home aims to provide holistic, multidisciplinary and person-centered care. One of the services provided in several integrated care programs is the comprehensive geriatric assessment (CGA), conducted to identify older adults' care needs. There are many different ways to conduct a CGA. However, it is still unclear what distinguishes these CGAs from each other. Also, it is yet unknown to what extent the principles of integrated care (holism, multidisciplinary and person-centredness) are reflected in CGAs. The objectives of this study were to: 1) compare different CGAs conducted within integrated care programs for older adults living at home, and 2) describe how the principles of integrated care were applied in these CGAs. Methods: A systematic literature search was conducted to identify integrated care programs including a CGA. Data were extracted on main characteristics of identified CGA tools and procedures, and how principles of integrated care were applied in these CGAs. Results: Twenty-six programs were included in this study, of which most were implemented in the Netherlands and the U.S. Twenty different CGA tools and procedures were identified. The majority of CGAs aimed to have a holistic, multidisciplinary and person-centred approach, although the way and extent to which CGAs incorporated these principles differed. Discussion: This systematic review highlights the variability of CGAs used in integrated care for older people living at home. The overview of CGAs may promote further exchange of CGA tools and procedures, and help researchers and professionals to not reinvent the wheel.

DO REPORT CARDS PREDICT FUTURE QUALITY? THE CASE OF SKILLED NURSING FACILITIES

P. Cornell¹, D. Grabowski², E. Norton³, M. Rahman⁴, 1. *Brown University*, 2. *Harvard Medical School*, 3. *Department of Health Policy and Management, University of Michigan*, 4. *Department of Health Services, Policy and Practice, Brown University*

Report cards are intended to improve consumer decision-making and foster a market for quality. However, inadequate risk adjustment of report card measures often biases comparisons across firms. We test whether skilled nursing facility (SNF) star ratings causally predict the

quality-related outcomes that are important to patients and providers. We examine to effect of an increase in the star rating of a chosen SNF after an inpatient hospital stay on the number of days spent in SNF, hospital, in the community with and without home health care, and deceased. To identify the causal effect of star rating on health care outcomes, we exploit variation over time in the distance from a patient's residential ZIP code to SNFs. We found that patients who go to higher-rated SNFs achieved better outcomes, supporting the validity of the SNF report card ratings. A one-star increase is associated with a 5% decrease in mortality over 180 days after SNF admission, as well as 3.3% and 2.4% decrease in number of days spent in inpatient hospital and SNF care, respectively; and 4.0% and 2.5% increase in number of days spent at home in the community with and without home health care.

DOES RETIREMENT AFFECT MAMMOGRAPHY USE?

P. Eibich¹, L. Goldzahl², 1. *Max Planck Institute for Demographic Research*, 2. *University of Manchester*

Breast cancer is one of the major causes of death among women aged 50 and above. Early detection considerably improves survival, and therefore many countries have introduced breast cancer screening programs. This paper examines the causal impact of retirement on the uptake of mammography screening. We use data from a series of Eurobarometer surveys conducted between 1996 and 2006, covering 25 different European countries. We address the endogeneity of retirement by using age thresholds for pension eligibility as instruments in a fuzzy regression discontinuity design. The results show that retirement reduces the probability of participating in breast cancer screening. This effect is not driven by changes in health, income or physician behavior. Instead, we find evidence that retired women invest less in preventive care in general. Specification curve analysis indicates that these results are robust to changes in model specification.

EXPLORING THE BALANCE: A PATH ANALYSIS EXAMINATION OF THE MAINTAINING THE BALANCE MODEL

S. Pennell, C. Jacelon, J. Choi, *U. Massachusetts Amherst*

Home is the desired site of care for most adults in the US (Eckert, Morgan & Swamy, 2004). The number of older adults who potentially require supportive care at home is rapidly growing. Our current healthcare infrastructure must evolve to accommodate the preference for homecare. This investigation provides quantitative testing of a qualitatively-derived model called Maintaining the Balance (MTB) developed by Jacelon (2010). MTB describes the factors that drive a community-dwelling older adult's ability to achieve optimal wellness (OW) in the home environment. Secondary analysis of data collected for the National Social Life, Health, and Aging Project (NSHAP) was utilized. Path analysis technique was employed to explore the effect of the 5 balance variables: health, activity, autonomy, attitude, and relationships on the outcome variable of OW. Attitude was the variable with the largest effect on OW. Activity, health, and relationships also had significant effects on OW. Autonomy and age were non-significant factors. The hypothesized model was partially supported by the NSHAP dataset.

HEALTH COMMUNICATION EFFECTIVENESS: INTERACTIONS BETWEEN AGE, HEALTH RISK, AND MESSAGE FRAMING

E. Pinyan, T. Hess, *North Carolina State University*

Health communications convey information about the effects of behavior modifications on health outcomes, and thus have the potential of influencing health-enhancing decisions. Message framing is important in such communications, in that focusing on the benefits of adopting a behavior or the costs of failing to adopt a behavior may interact with person or situational factors in determining effectiveness. We hypothesized that age differences in goal orientation (growth vs. loss prevention) and risk status for a health outcome will influence the effectiveness of framed messages. Middle-aged ($n=60$, $M=50.0$ years) and older adults ($n=60$, $M=75.4$ years) completed a questionnaire to estimate 20-year risk of developing cardiovascular disease (CVD) and were informed if they were low or high risk. Participants were then presented with either loss- or gain-framed messages discussing eight behavioral factors associated with CVD (e.g., diet, smoking). Participants indicated their intentions to engage in behavioral modifications of the eight factors. ANOVAs examining behavioral modifications revealed a significant three-way interaction between age, risk, and message. Consistent with hypothesized age changes in goal orientation, loss-framed messages were more effective in influencing intentions in older than in middle-aged adults, whereas the opposite was true for gain-framed messages. Risk status further moderated these effects, with effectiveness of loss-based messages in older adults being especially strong in those identified as high-risk. These findings have important implications for optimizing health messages to motivate behavior change, suggesting that both age and risk status need to be considered in constructing effective communications.

IMPACT OF SOCIAL FACTORS ON RISK ADJUSTMENT FOR THE QUALITY MEASURE FOR SELF-REPORTED PAIN IN NURSING HOME RESIDENTS

L. Campbell¹, A. Vadnais², Q. Li¹, D. Barch¹, T. McMullen³, L. Smith¹, 1. *RTI International*, 2. *RTI International*, Waltham, MA, USA, 3. *Centers for Medicare & Medicaid Services*

When healthcare outcomes vary by social factors, including these characteristics in risk-adjustment models may improve quality measurement accuracy. Therefore, we examined the impact of potential social risk factors available in the Minimum Data Set 3.0 on the nursing home quality measure, Percent of Residents Who Self-Report Moderate to Severe Pain (Long-Stay). This measure is risk-adjusted for cognitive independence (CI), as evidence suggests this population is better able to self-report pain. Social factors tested included age (>85), sex, marital status, and Medicaid status. In national analyses, we examined the distribution of social factors across facilities to determine variability, and used Chi-squared tests to determine whether self-reported pain prevalence was significantly different in residents with and without each social factor. Bivariate results indicated age (>85) and male sex were associated with lower pain prevalence ($p<0.001$); thus, we tested risk-adjustment models with CI and dummy variables for age (>85 ; Model 1), sex (Model