B-973, a Novel $\alpha 7$ nAChR Ago-PAM: Racemic and Asymmetric Synthesis, Electrophysiological Studies, and *in Vivo* Evaluation

Sumanta Garai,[†] Krishnamohan S. Raja,[†] Roger L. Papke,[§] Jeffrey R. Deschamps,^{||} M. Imad Damaj,[‡] and Ganesh A. Thakur^{*,†,§}

[†]Department of Pharmaceutical Sciences, Bouvé College of Health Science, Northeastern University, 140 The Fenway, 360 Huntington Avenue, Boston, Massachusetts 02115, United States

[§]Department of Pharmacology and Therapeutics, University of Florida, Gainesville, Florida 32611, United States

^{||}Naval Research Laboratory, Code 6930, 4555 Overlook Avenue, Washington, D.C. 20375, United States

[‡]Department of Pharmacology and Toxicology, Virginia Commonwealth University, Richmond, Virginia 23284, United States

S Supporting Information

ABSTRACT: We report here the total synthesis of B-973 (five steps), a recently identified $\alpha 7$ nAChR ago-PAM, its enantiomeric resolution, and its electrophysiological characterization in *Xenopus* oocytes to identify (–)-B-973B as the bioactive enantiomer. The asymmetric synthesis of B-973B was accomplished in 99% ee, and X-ray crystallography studies revealed its absolute “S” stereochemistry. B-973B was effective in attenuating pain behavior and decreasing paw edema (formalin test), and its analgesic effects were mediated through $\alpha 7$ nAChR.

KEYWORDS: Alpha7 nicotinic acetylcholine receptors, ligand-gated ion channels, positive allosteric modulators, Ago-PAMs, neuropathic pain, asymmetric synthesis, electrophysiological characterization

Nicotinic acetylcholine receptors (nAChRs) are members of the Cys-loop superfamily of cationic ligand-gated ion channels and are involved in the physiological responses to the neurotransmitter, acetylcholine (ACh).^{1–3} These receptors are distributed throughout the central and peripheral nervous systems and play important roles in several (patho)-physiological processes. Compounds that selectively target the homopentameric $\alpha 7$ nAChR subtype are being pursued as potential pharmacotherapy agents for treating cognitive dysfunctions in schizophrenia and Alzheimer’s disease, as well as for treating neuropathic and inflammatory pain. This type of targeting is safe and is not associated with opioids-like side effects such as addiction liability.^{4–7} Attempts to target $\alpha 7$ nAChR has mainly focused on two approaches: partial agonism via the ACh binding site and positive allosteric modulation (PAM) at sites topographically distinct from ACh binding. Although $\alpha 7$ nAChR-selective and potent (partial) agonists have been developed and studied (pre)clinically, the rapid desensitization of $\alpha 7$ nAChR agents in response to high agonist concentration, together with the possibility of endogenous tone disruption, has led to some concerns regarding their utility as clinical candidates.^{2,4,5} PAMs of the $\alpha 7$ nAChR provide an alternative targeting approach that

facilitates natural signaling by endogenous ACh and thus is differentiated from partial agonists.⁸

In addition to PAMs, allosteric agonists as well as silent agonists of $\alpha 7$ nAChR have been discovered.⁶ However, despite all of these efforts, only limited success has been achieved clinically with galantamine (Figure 1). JNJ-39393406 is an $\alpha 7$ PAM that is currently undergoing phase II clinical trials for depressive disorders and smoking cessation.^{9,10} Most of the drug candidates that were pursued as $\alpha 7$ nAChR (partial) agonists or PAMs, however, failed in Phase II clinical trials for cognitive disorders due to insufficient efficacy. Recently, a unique class of compounds with dual modes of action—allosteric agonism and positive allosteric modulation—have been discovered and are referred to as “ago-PAMs” (e.g., GAT107, Figure 1).^{11–14}

Ago-PAMs of $\alpha 7$ nAChR activate receptors through site distinct from orthosteric sites and positively modulate orthosteric cholinergic signaling while preserving the spatio-temporal features of synaptic transmission.^{13,14} Such modu-

Received: September 5, 2018

Accepted: October 11, 2018

Published: October 11, 2018

Figure 1. Representative selective positive allosteric modulators of $\alpha 7$ nAChR: an approved drug (1), phase II clinical candidate (2), and preclinical candidates (3–6).

lators that act through dual modes of action are expected to be more efficacious and potentially address the limitations observed in partial agonists and PAMs in clinical trials. Indeed, we recently demonstrated that one such dual modulator, GAT107, from the tetrahydroquinoline (THQ) class of compounds is an orally bioavailable and efficacious analgesic for treating inflammatory and neuropathic pain in animal models.¹⁵

Recently, Bristol Myers Squibb reported HTS screening of a diverse small molecule library ($>10^6$ molecules) using a fluorescence-based assay configured to detect $\alpha 7$ nAChR PAMs.¹⁶ Active compounds identified from the screen included a series of piperazine-containing molecules that displayed robust $\alpha 7$ nAChR PAM activity. One compound from this series, B-973 (Figure 1), was unique in that it was shown to be a potent ago-PAM. Although this important work led to the identification of a novel scaffold with $\alpha 7$ nAChR ago-PAM activity, it left several questions unanswered, thus limiting further drug optimization and developmental efforts on this scaffold. Based on the information provided, it is not clear if B-973 is a racemate or a single enantiomer. Neither its absolute stereochemistry nor the activity of the other enantiomer are known. Information on the chemical synthesis of this compound to facilitate *in vivo* studies and library syntheses is also not reported.

Given the therapeutic potential of $\alpha 7$ nAChR ago-PAMs and the excitement surrounding this structurally novel scaffold, an effort was made to pursue its synthesis and enantiomeric separation and identify the stereochemical requirements for $\alpha 7$ nAChR ago-PAM activity through electrophysiological studies followed by *in vivo* evaluation of the active enantiomer. We report an efficient and scalable synthesis of the racemate as well as of the active enantiomer in 99% ee. Access to copious amounts of the active enantiomer enabled its preclinical evaluation for inflammatory pain (formalin test) in a mouse model, where it was shown to be efficacious. The five-step synthetic route reported here was optimized for producing gram-scale quantities of B-973 and is depicted in Scheme 1.

The synthesis begins with the formation of tricyclic bromo pyrazine derivative **9** by monosubstitution of dibromo pyrazine **7** with piperazine **8** in the presence of K_2CO_3 in DMF in 92% yield. To install a keto functionality on **9**, we examined several reaction conditions involving $BuLi$ -mediated lithiation followed by quenching with appropriate electrophiles. Our initial

Scheme 1. Racemic Synthesis of B-973

efforts were plagued by the formation of a range of undesired products, which was presumably due to the sensitivity of the pyrazine ring to these conditions. We further attempted installing the ketone moiety under mild Pd-catalyzed cross-coupling conditions.¹⁷ Stille cross coupling of compound **9** with tributyl (1-ethoxyvinyl)tin and $Pd(PPh_3)_2Cl_2$ in the presence of CuI yielded the enol ether **10**. The crude product was hydrolyzed with aq. HCl to give the desired ketone compound **11** in 82% yield over two steps. Thus, after obtaining the key keto intermediate **11** in multigram quantities, we carried out its conversion to an imine with NH_4OAc , followed by *in situ* reduction with $NaCNBH_4$ to yield racemic amine **12** in 96% yield. The conversion of acid **13** to the corresponding acid chloride **14**, followed by its coupling with the amine fragment **12** in the presence of pyridine and catalytic DMAP yielded the racemic B-973, **6** in 54% yield (Scheme 1).

B-973 has one chiral center with two possible configuration. To test whether the allosteric agonist and PAM activity was attributed to a single enantiomer or was present in both compounds, we first developed conditions for the enantiomeric separation of B-973 (Scheme 2; see SI). Using chiral super fluid chromatography (SFC), baseline separation of the B-973 enantiomers was accomplished in 99% ee.

Both enantiomers were then tested in *Xenopus* oocytes expressing human $\alpha 7$ nAChR with a two-electrode voltage clamp method (Figure 2). Following the measurement of 60

Scheme 2. Enantiomeric Resolution of Racemic B-973

Figure 2. Electrophysiological characterization of the B-973 isomers on human $\alpha 7$ nAChR expressed in *Xenopus* oocytes.

μ M ACh control responses, the B-973 isomers were coapplied with a control concentration at 10 or 100 μ M. The average net-charge responses ($\pm$ SEM) of the oocytes ($n \geq 5$) to the A and B isomers, normalized to the initial ACh control responses from the same cells, are shown in Figure 2A. While there were negligible effects from the B-973A isomer, the B-973B isomer greatly increased the responses compared to those evoked by ACh alone.

Following measurement of the 60 μ M ACh control responses, the B-973 isomers were applied alone at a concentration of 100 μ M, followed by another application of 60 μ M ACh. The average net-charge responses ($\pm$ SEM) of the oocytes ($n = 6$) to the A and B isomers, normalized to the initial ACh control responses from the same cells, are shown in Figure 2B. The B-973A isomer failed to evoke a response, and the subsequent ACh controls were not significantly different from the initial controls. In contrast, B-973B evoked large responses, and the ACh responses greatly increased after washout compared to initial controls. The normalized net-charge responses of $\alpha 7$ -expressing cells to 10 μ M B-973B were compared to 10 μ M of the previously characterized ago-PAM GAT107 (Figure 2C).^{12,13} While the initial allosteric activation produced by the two agents was similar, the primed potentiation of the subsequent 60 μ M ACh response was greater ($p < 0.05$) for B-973B than for GAT107. Additionally, the responses to racemic B-973 were smaller ($p < 0.05$) than the response to the B-973B enantiomer. These studies thus concluded that the ago-PAM activity resided exclusively in (–)-B-973B, while (+)-B-973A was inactive at $\alpha 7$ nAChR.

To understand the absolute stereochemical requirements for $\alpha 7$ nAChR ago-PAM activity, we attempted to crystallize both enantiomers to obtain crystals suitable for X-ray studies. However, even after several crystallization attempts, neither of the B-973 enantiomers could be crystallized in an acceptable form. Furthermore, to facilitate structure–activity relationship (SAR) studies, there was a need for easy access to the homochiral version of amine 12 due to the observed high enantiospecificity for $\alpha 7$ nAChR activity. We therefore decided to develop an enantioselective approach that would yield the active enantiomer in high ee and allow derivatization of the

chiral amine to expand the possibility of obtaining crystals suitable for X-ray studies.

Among the various methods for the preparation of optically active amines,¹⁸ we planned to synthesize chiral 12 via an approach involving imine formation with chiral *tert*-butanesulfinamide, followed by diastereoselective reduction with the appropriate reducing agent as a key step.¹⁹ Advantageously, both the enantiomers of *tert*-butanesulfinamide are inexpensive and commercially available in high optical purity. Moreover, condensation of *tert*-butanesulfinamide and ketone gives stable ketimine under mild reaction conditions, yields good diastereoselectivity during reduction, and allows easy removal of the *N*-*tert*-butanesulfinyl in high yield.

Accordingly, the chiral *N*-*tert*-butanesulfinyl ketimine 17 was synthesized from ketone 11 with *R*-(+)-2-methyl-2-propane sulfinamide in the presence of $\text{Ti}(\text{OEt})_4$ in dichloromethane under reflux conditions as a yellow solid in 89% yield. The chiral *N*-*tert*-butanesulfinyl ketimine 17 was stereospecifically reduced by 9-BBN^{20–22} in THF at -78°C to yield sulfinamide 18 as a single diastereomer ($\text{dr} \geq 99\%$; determined by ^1H and ^{13}C NMR spectroscopy) in 96% yield. The hydrolysis of the sulfinamide group of 17 was accomplished using 6 M HCl in methanol and gave chiral amine 19 in 86% yield. Finally, the coupling of chiral amine 19 with acid chloride 14 yielded optically pure 15 in 94% ee (Scheme 3). By comparing the optical rotation of 15 with the enantiomers obtained from chiral SFC, 15 was identified as the inactive enantiomer.

Scheme 3. Asymmetric Synthesis of (+)-B-973A

To obtain the other enantiomer, *N*-*tert*-butanesulfinyl ketimine 17 was stereospecifically reduced with 1-Selectride^{20–22} in anhydrous THF at -78°C to yield 20 as a single diastereomer ($\text{dr} \geq 99\%$; determined by ^1H and ^{13}C NMR spectroscopy) in 92% yield. An X-ray crystallographic study revealed the absolute stereochemistry of 20 at C4 to be “S”. The hydrolysis of the sulfinamide group of 20 using 6 M HCl gave the chiral amine 21 in 86% yield. The absolute “S” stereochemistry of the chiral center C4 was further confirmed by a single-crystal X-ray study of *p*-nitrobenzoyl derivative 22 of chiral amine 21 (see SI). Finally, the coupling of 21 with acid chloride 14 yielded the optically pure bioactive isomer 16, (S)-(–)-B-973B, in 56% yield and with 99% optical purity (Scheme 4).

We previously demonstrated the antinociceptive and anti-inflammatory efficacy of GAT107 toward pain in a series of

Scheme 4. Asymmetric Synthesis of (–)-B-973B

mouse models.¹⁵ Here, we tested whether the systemic administration of B-973B would produce antinociception in mouse formalin testing. As seen in Figure 3A, B-973B did significantly reduce nociceptive behavior during Phase I [$F_{(3,28)} = 3.941$, $P = 0.018$] at the highest dose of 10 mg/kg. Moreover, B-973B (1–10 mg/kg, i.p.) dose-dependently attenuated formalin-induced paw licking behavior in Phase II [$F_{(3,28)} = 13.07$, $P < 0.001$] with 3 and 10 mg/kg doses, differing from the vehicle group.

Figure 3. *In vivo* evaluation of the active enantiomer B-973B in a mouse model of inflammatory pain and establishment of target specificity.

In addition, B-973B significantly reduced paw edema [$F_{(3,28)} = 21.05$, $P < 0.001$; Figure 3B] in mice treated with 3 and 10 mg/kg doses, differing from the vehicle group. We next explored the possible role of $\alpha 7$ nAChRs as related to B-973B's effect in the formalin test. To study this, we tested whether the $\alpha 7$ nAChR antagonist, MLA, blocks the antinociceptive effects of B-973B in the formalin test. As shown in Figure 3C, a significant effect of treatment was found [$F_{(3,15)} = 33.05$; $P < 0.0001$]. MLA given alone significantly affected the nociceptive behavior ($P = 0.77$). MLA (10 mg/kg, s.c.) totally blocked the antinociceptive effect of B-973B in both phases I and II of the formalin test ($P < 0.001$).

Acute B-973B mainly attenuated pain behavior in the second phase of the formalin test, which is associated with the development of inflammation and spinal dorsal horn sensitization but was only effective in phase I at the highest dose (immediately after formalin injection), which is mediated

by C-fiber activity.^{23,24} Moreover, B-973B decreased formalin-induced paw edema, which is consistent with the idea that it acts on the inflammatory phase of the formalin test. Furthermore, using a complementary pharmacological agent (i.e., the $\alpha 7$ nAChR antagonist, MLA), we confirmed that the antinociceptive effects of B-973B in the formalin test were mediated through $\alpha 7$ nAChRs.

Thus, in this work, we have developed the first total, gram-scale synthesis of B-973, a structurally novel $\alpha 7$ nAChR ago-PAM. We also developed conditions for its enantiomeric separation using chiral SFC to obtain both enantiomers in high enantiomeric excess. The (–)-B-973B enantiomer retained all $\alpha 7$ nAChR ago-PAM activity, while the (+)-B-973B was inactive in electrophysiological assays. Asymmetric synthesis of (–)-B-973B was accomplished in high enantiomeric excess (99% ee). When tested *in vivo*, B-973B was effective in attenuating pain behavior and decreased formalin-induced paw edema (formalin test). Target specificity of B-973B for analgesic activity was established using MLA. B-973B was as effective as structurally distinct GAT107 in *in vitro* assays and *in vivo* experiments, thus providing a suitable lead for future SAR optimization.

■ ASSOCIATED CONTENT

S Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsmmedchemlett.8b00407.

Experimental details, structural characterization data, chiral SFC separation details, NMR spectra, X-ray study details, and biological studies (PDF)

Accession Codes

CCDC 1842480 and 1842483 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: g.thakur@northeastern.edu.

ORCID

Ganesh A. Thakur: 0000-0002-7468-8819

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the National Institute of Health (NIH) RO1-EY024717 to G.A.T. and RO1-GM057481 to R.L.P., and a grant from a Pilot Project from the VCU Massey Cancer Center and RO1-CA206028 to M.I.D. Oocyte experiments were conducted by Clare Stokes and Alexander den Boef.

■ REFERENCES

- (1) Bertrand, D.; Terry, A. V. The wonderland of neuronal nicotinic acetylcholine receptors. *Biochem. Pharmacol.* **2018**, *151*, 214.
- (2) Papke, R. L. Merging old and new perspectives on nicotinic acetylcholine receptors. *Biochem. Pharmacol.* **2014**, *89*, 1.

- (3) Stokes, C.; Treinin, M.; Papke, R. L. Looking below the surface of nicotinic acetylcholine receptors. *Trends Pharmacol. Sci.* **2015**, *36*, 514.
- (4) Bagdas, D.; Gurun, M. S.; Flood, P.; Papke, R. L.; Damaj, M. I. New Insights on Neuronal Nicotinic Acetylcholine Receptors as Targets for Pain and Inflammation: A Focus on $\alpha 7$ nAChRs. *Curr. Neuropharmacol* **2018**, *16*, 415.
- (5) Hone, A. J.; McIntosh, J. M. Nicotinic acetylcholine receptors in neuropathic and inflammatory pain. *FEBS Lett.* **2018**, *592*, 1045.
- (6) Manetti, D.; Bellucci, C.; Chiaramonte, N.; Dei, S.; Teodori, E.; Ro manelli, M. N. Designing selective modulators for the nicotinic receptor subtypes: challenges and opportunities. *Future Med. Chem.* **2018**, *10*, 433.
- (7) Hudzik, T. J.; Basso, A. M.; Lynch, J. J.; Bracken, W. M.; Mohler, E. G.; Kohlhaas, K. L.; Xu, H.; Haig, G.; Gault, L. Preclinical abuse liability assessment of ABT-126, an agonist at the $\alpha 7$ nicotinic acetylcholine receptor (nAChR). *Pharmacol., Biochem. Behav.* **2017**, *158*, 22.
- (8) Williams, D. K.; Wang, J.; Papke, R. L. Positive allosteric modulators as an approach to nicotinic acetylcholine receptor-targeted therapeutics: advantages and limitations. *Biochem. Pharmacol.* **2011**, *82*, 915.
- (9) Perkins, K. A.; Roy Chengappa, K. N.; Karelitz, J. L.; Boldry, M. C.; Michael, V.; Herb, T.; Gannon, J.; Brar, J.; Ford, L.; Rassnick, S.; Brunzell, D. H. Initial Cross-Over Test of A Positive Allosteric Modulator of Alpha-7 Nicotinic Receptors to Aid Cessation in Smokers With Or Without Schizophrenia. *Neuropsychopharmacology* **2018**, *43*, 1334.
- (10) Winterer, G.; Gallinat, J.; Brinkmeyer, J.; Musso, F.; Kornhuber, J.; Thuerauf, N.; Rujescu, D.; Favis, R.; Sun, Y.; Franc, M. A.; Ouwerkerk-Mahadevan, S.; Janssens, L.; Timmers, M.; Streffer, J. R. Allosteric $\alpha 7$ nicotinic receptor modulation and P50 sensory gating in schizophrenia: a proof-of-mechanism study. *Neuropharmacology* **2013**, *64*, 197.
- (11) Gill, J. K.; Savolainen, M.; Young, G. T.; Zwart, R.; Sher, E.; Millar, N. S. Agonist activation of $\alpha 7$ nicotinic acetylcholine receptors via an allosteric transmembrane site. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 5867.
- (12) Thakur, G. A.; Kulkarni, A. R.; Deschamps, J. R.; Papke, R. L. Expedient synthesis, enantiomeric resolution, and enantiomer functional characterization of (4-(4-bromophenyl)-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]quinoline-8-sulfonamide (4BP-TQS): an allosteric agonist-positive allosteric modulator of $\alpha 7$ nicotinic acetylcholine receptors. *J. Med. Chem.* **2013**, *56*, 8943.
- (13) Papke, R. L.; Horenstein, N. A.; Kulkarni, A. R.; Stokes, C.; Corrie, L. W.; Maeng, C. Y.; Thakur, G. A. The activity of GAT107, an allosteric activator and positive modulator of $\alpha 7$ nicotinic acetylcholine receptors (nAChR), is regulated by aromatic amino acids that span the subunit interface. *J. Biol. Chem.* **2014**, *289*, 4515.
- (14) Horenstein, N. A.; Papke, R. L.; Kulkarni, A. R.; Chaturbhuj, G. U.; Stokes, C.; Manther, K.; Thakur, G. A. Critical Molecular Determinants of $\alpha 7$ Nicotinic Acetylcholine Receptor Allosteric Activation: Separation of Direct Allosteric Activation and Positive Allosteric Modulation. *J. Biol. Chem.* **2016**, *291*, 5049.
- (15) Bagdas, D.; Wilkerson, J. L.; Kulkarni, A.; Toma, W.; AlSharari, S.; Gul, Z.; Lichtman, A. H.; Papke, R. L.; Thakur, G. A.; Damaj, M. I. The $\alpha 7$ nicotinic receptor dual allosteric agonist and positive allosteric modulator GAT107 reverses nociception in mouse models of inflammatory and neuropathic pain. *Br. J. Pharmacol.* **2016**, *173*, 2506.
- (16) Post-Munson, D. J.; Pieschl, R. L.; Molski, T. F.; Graef, J. D.; Hendricson, A. W.; Knox, R. J.; McDonald, I. M.; Olson, R. E.; Macor, J. E.; Weed, M. R.; Bristow, L. J.; Kiss, L.; Ahljanian, M. K.; Herrington, J. B-973, a novel piperazine positive allosteric modulator of the $\alpha 7$ nicotinic acetylcholine receptor. *Eur. J. Pharmacol.* **2017**, *799*, 16.
- (17) Sato, N.; Narita, N. Studies on Pyrazines; 38: Acylation of Bromopyrazines and 2-Bromopyridine via Copper-Cocatalytic Stille Reaction. *Synthesis* **2001**, *2001*, 1551.
- (18) Nugent, T. C.; El-Shazly, M. Chiral Amine Synthesis – Recent Developments and Trends for Enamide Reduction, Reductive Amination, and Imine Reduction. *Adv. Synth. Catal.* **2010**, *352*, 753.
- (19) Robak, M. T.; Herbage, M. A.; Ellman, J. A. Synthesis and Applications of tert-Butanesulfinamide. *Chem. Rev.* **2010**, *110*, 3600.
- (20) Chelucci, G.; Baldino, S.; Chessa, S. Diastereoselective reduction of enantiopure N-p-toluenesulfinyl ketimines derived from pyridyl ketones. *Tetrahedron* **2006**, *62*, 619.
- (21) Chelucci, G.; Baldino, S.; Chessa, S.; Pinna, G. A.; Soccolini, F. An easy route to optically active 1-substituted-1-pyridyl-methylamines by diastereoselective reduction of enantiopure N-tert-butanesulfinyl ketimines. *Tetrahedron: Asymmetry* **2006**, *17*, 3163.
- (22) Chelucci, G.; Baldino, S.; Solinas, R.; Baratta, W. Asymmetric synthesis of 1-substituted-1-(pyridin-2-yl)methylamines by diastereoselective reduction of enantiopure N-p-toluenesulfinyl ketimines. *Tetrahedron Lett.* **2005**, *46*, 5555.
- (23) Abbott, F. V.; Franklin, K. B.; Westbrook, R. F. The formalin test: scoring properties of the first and second phases of the pain response in rats. *Pain* **1995**, *60*, 91.
- (24) Davidson, E. M.; Carlton, S. M. Intraplantar injection of dextrorphan, ketamine or memantine attenuates formalin-induced behaviors. *Brain Res.* **1998**, *785*, 136.