Published in final edited form as:

Biochemistry. 2018 March 13; 57(10): 1577–1581. doi:10.1021/acs.biochem.8b00053.

Improved Analysis of RNA Localization by Spatially-Restricted Oxidation of RNA-Protein Complexes

Ying Li¹, Mahima B. Aggarwal¹, Ke Ke¹, Kim Nguyen¹, and Robert C. Spitale^{1,2}

¹Department of Pharmaceutical Sciences University of California, Irvine. Irvine, California. 92697

²Department of Chemistry. University of California, Irvine. Irvine, California. 92697

Abstract

Recent analysis of transcriptomes has revealed that RNAs perform a myriad of functions beyond coding for proteins. Critical to RNA function is its transport to unique subcellular locations. Despite the importance of RNA localization it is still very challenging to study in an unbiased manner. We recently described the ability to tag RNA molecules within subcellular locations through spatially-restricted nucleobase oxidation. Herein we describe a dramatic improvement of this protocol through the localized oxidation and tagging of proteins. Isolation of RNA-protein complexes enabled the enrichment of challenging RNA targets on chromatin and presented a considerably optimized protocol for the analysis of RNA subcellular localization within living cells.

Graphical Abstract

Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.

RNA molecules perform key functions at the heart of many normal biological pathways and are key drivers in the onset of many diseases.^(1–3) The localization or sequestration of RNA molecules to specific subcellular compartments is critical to the regulation of their functions.

Corresponding Author: rspitale@uci.edu.

Author Contributions

All authors have given approval to the final version of the manuscript.

Funding Sources

No competing financial interests have been declared.

Supporting Information

Biochemical methods, plasmids, primers, antibodies, protein gels, western blot, dot blot, scheme of enrichment.

(4–6) Despite the importance of RNA localization, there is a real lack of methods to systematically analyze RNA location inside cells in an unbiased manner^(7–10).

Our laboratory recently developed a novel method to assay RNA localization within intact living cells through spatially-restricted nucleobase oxidation.⁽¹¹⁾ Using Halo-tag fusion proteins, we localized the oxygen photosensitizer dibromofluorescein (DBF) within subcellular compartments. Blue light exposure generated short-lived singlet oxygen, resulting in guanosine oxidation in RNAs nearby in space. Oxidized guanosines can react with nucleophiles in solution and as such we utilized propargyl amine (PA) to append alkyne handles onto these RNAs for downstream study. RT-qPCR data demonstrated that this oxidative approach provides subcellular, and even sub-organellar resolution for RNA localization studies (Scheme 1A).

Despite our exciting results, two observations convinced us to seek an improved RNA tagging approach. First, the overall yield of guanosine labeling is far from satisfactory (~10% on a free oligonucleotide in solution⁽¹¹⁾). Considering the various singlet oxygen quenchers and competitive nucleophiles in cells, we anticipated even lower reactivity. Second, several positive-control chromatin-associated RNAs were not tagged and enriched, even when localizing DBF to chromatin, through Histone 2B (H2B)-Halo tethering. This result was especially surprising as confocal fluorescence imaging data indicated robust and localized PA adduct formation in the nucleus. These observations challenged us to interrogate the molecular details of our approach. Within this Communication we report a dramatic improvement in our approach, through the isolation of RNA-protein complexes, which have been tagged through singlet oxygen-mediated tagging (Scheme 1B). We anticipate that our improved method provides a streamlined and experimentally superior approach to increase the yield of subcellular RNA tagging in an unbiased manner.

To systematically identify the biomolecules which form adducts with PA, cells were processed as previously described, lightly fixed, permeabilized, and incubated with RNase, DNase, or proteinase, respectively (Supporting Information, SI). PA adducts were visualized in fluorescence confocal microscopy through Cy5 signals after copper-catalyzed azide-alkyne cycloaddition (CuAAC) treatment. Imaging results (Figure 1A) revealed that RNase and DNase treatment didn't cause significant Cy5 signal reduction. In contrast, Cy5 signal was undetectable after proteinase treatment, while Hoechst stain remained. These results suggested that the large majority of PA adducts in cells, and the corresponding signal in imaging experiments, stem from PA-protein adducts.

Many examples in the literature have demonstrated that electron-rich side chains (phenylalanine, tryptophan, histidine, cysteine) can be oxidized in presence of singlet oxygen.^(12–16) Oxidized amino acids can then be attacked by nucleophiles to form adducts.^(16, 17) These observations, coupled with our results above, prompted us to explore whether our approach was capable of forming PA adducts with proteins *in vitro*.

We utilized bovine serum albumin (BSA) as a model protein. We incubated BSA with tetrabromofluorescein (TBF; Figure 1B) and PA in the presence of blue light. After incubation, PA adducts were labeled by Cu-AAC with Cy5 azide and imaged using

denaturing gel electrophoresis. As shown in Figure 1C, Cy5 signal was only observed in the presence of TBF, PA, and light. We also performed a time-course analysis of adduct formation and observed increasing Cy5 signals over time (Figure 1D). In contrast, increasing concentrations of sodium azide (singlet oxygen quencher⁽¹⁸⁾, NaN₃) resulted in decreasing adduct formation (Figure 1E). The same concentrations of sodium chloride (NaCl) didn't affect the Cy5 signals, suggesting that the protein modification was indeed driven by singlet oxygen. These results confirmed that proteins are also subjected to singlet-oxygen induced oxidation and nucleophilic addition under the conditions previously used for imaging, RNA tagging and enrichment.

Our observations suggest that proteins are robustly tagged with PA *in vitro* and they contribute most to the PA adduct formation in cells. These results may explain the inability to tag certain RNAs that are tightly associated with protein complexes and have limited solvent accessibility. Recent data demonstrated that the large majority of RNA sequences are covered or interacting with protein molecules.^(19–21) RNAs might have even more hampered solvent accessibility within densely packed heterochromatin regions^(22–24), thus were not detected by our previous labeling method. We were enticed to explore if protein tagging could also take place in a spatially-restricted manner, providing a roadmap for the possibility of utilizing such characteristics for assaying RNA localization.

To ensure that protein labeling still affords tight spatial resolution, we performed protein analysis in H2B-Halo expressing cells (Figure 2A). We first confirmed the adduct formation with adjacent histone protein H3 via histone protein extraction after light treatment (SI, Figure S1). To systematically test the spatial resolution of proteins in detail, we picked six proteins to compare: H3, HDAC1, and hnRNPH (chromatin associated)⁽²⁵⁾; Nucleolin (nucleolus)⁽²⁶⁾; Gapdh and Alpha Tubulin (cytoplasm)^(27, 28). After treatment, the cells were lysed and the lysate were submitted to biotinylation through CuAAC with biotin-azide, enrichment via streptavidin-biotin interaction, gel electrophoresis and western blot (SI). As shown in Figure 2B, the chromatin associated proteins, H3, HDAC1 and hnRNPH, were highly enriched (~10% of the input). Nucleolin, Gapdh, and alpha Tubulin, despite being highly abundant in the cells were not enriched.

To further test our resolution, we compared these results to a nuclear localization sequence (NLS)-Halo fusion, which has diffuse nuclear and nucleolus localization (Figure S2).⁽²⁹⁾ As shown in Figure S3, the nuclear and particularly nucleolus proteins were enriched, albeit to a much lower percent, compared to the 10% input loading. These results further suggest that tagging is tightly controlled by the spatial proximity of DBF to the protein of interest, which could lay the foundation for us to assay RNA localization through RNA-protein complexes (Scheme 1B). We foresee our protein tagging approach may complement other existing technologies such as APEX⁽³⁰⁾ and Bio-ID⁽³¹⁾ in a light controllable manner. Very recently, APEX has been applied to assay cellular RNA spatial organization with short-lived phenoxyl radicals (< 1 ms⁽³²⁾).⁽³³⁾ We anticipate that reactive singlet oxygen (< 3.5 μ s^(34, 35)) will provide a complementary approach.

To take advantage of spatially-restricted protein labeling for the analysis of RNA localization we utilized mild crosslinking (0.1% PFA), which has been shown to trap *in*

cellulo RNA-protein complexes.^(25, 33) We developed an optimized protocol for enrichment of protein-RNA complexes that had been modified with PA (SI, Figure S4). We profiled the RNA isolated from these ribonucleoprotein complexes by dot blot (Figure S5) and RT-qPCR. We utilized the positive-control chromatin-associated RNA 7SK as our benchmark and also focused on other chromatin-associated RNAs (XIST, KCNQ1OT1, HOTAIR)⁽²⁵⁾, which were barely detectable in our previous study. RNAs from other sites in the cell (U13 and U3 – nucleolus⁽³⁶⁾, GAPDH and ACTIN – cytoplasm^(37, 38)) were chosen as negative controls. As shown in Figure 3A, our new protocol based on RNA isolation after enrichment (IAE, shown in green) effectively enriched chromatin-associated RNAs. The fold enrichment was dramatically increased compared to our previous method (RNA isolation before enrichment (IBE); shown in black). Meanwhile, RNAs that localized primarily in the nucleolus and cytoplasm (shown in red) were not enriched by the RNP approach. These data jointly demonstrated that the enrichment of protein-RNA crosslinks improved the yield of RNAs without sacrificing the spatial resolution.

Our RT-qPCR result is consistent with literature reports, which rely on more complicated and laborious fractionation protocols.⁽³⁹⁾ Ruthenburg and coworkers assayed the distribution of RNAs in the nucleus by isolating RNAs separately from the soluble nuclear fraction and the insoluble chromatin fraction through multiple rounds of laborious cellular fractionation and centrifugation. For their calculation of enrichment they compared the Chromatin Pellet Extract (CPE) with the Soluble-Nuclear Extract (SNE) ratios. We compared our fold-enrichment analyses with the chromatin/nuclear ratios of the RNAs we interrogated. High enrichment of XIST and KCNQ1OT1 is consistent with their tight association with chromatin (high CPE/SNE); HOTAIR is more diffuse in the soluble nuclear fraction (low CPE/SNE ratio) and were less enriched.⁽³⁹⁾ Lastly, consistent between our approaches cytoplasmic RNAs GAPDH and ACTB had very low CPE/SNE ratios and extremely low enrichment by our approach. These results strongly suggest our experimental method can enrich RNAs within tightly associated subsections of cells with high resolution.

Finally, RNA enrichment with the NLS-Halo construct (diffuse nuclear localization) showed that chromatin-associated RNAs have much lower fold-enrichment and RNAs localized in the nucleolus are now enriched at a similar level if not higher to those on chromatin (Figure 3C). These results further demonstrated that the proximity of singlet oxygen determines the efficiency of PA-adduct formation (Figure 3D) and the new RNP protocol affords tight spatial resolution.

The results described here extend the possibilities of using spatially restricted singlet-oxygen formation to assay cellular localization of various biomolecules. We have confirmed the tight spatial resolution for protein labeling. In addition, we have reported an improved protocol to assay RNA localization by enriching RNA-protein complexes. We envision our optimized protocol will be readily applied to many other locations in cells as Halo-fusion proteins are now commercially available for many human proteins. The critical investigation herein lays an experimentally optimized roadmap for wide-scale and systematic analysis of cellular RNA localization in parallel to protein localization. Such studies are currently underway in our lab.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We thank members of the Spitale lab for their careful reading and critique of the manuscript. Spitale lab is supported by start up funds from the University of California, Irvine, and the NIH (1DP2GM119164 RCS). RCS is a Pew Biomedical Scholar.

ABBREVIATIONS

PA,	propargyl amine
DBF,	dibromofluorescein
TBF,	tetrabromofluorescein
ORF,	open reading frame
CuAAC,	coppercatalyzed azide-alkyne cycloaddition

REFERENCES

1. Mehler MF, and Mattick JS (2007) Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease, *Physiol. Rev* 87, 799–823. [PubMed: 17615389]
2. Pek JW, and Okamura K (2015) Regulatory RNAs discovered in unexpected places, *Wiley Interdiscip Rev RNA* 6, 671–686. [PubMed: 26424536]
3. Liu EY, Cali CP, and Lee EB (2017) RNA metabolism in neurodegenerative disease, *Disease Models & Mechanisms* 10, 509–518.
4. Geisler S, and Collier J (2013) RNA in unexpected places: long non-coding RNA functions in diverse cellular contexts, *Nat. Rev. Mol. Cell Biol* 14, 699–712. [PubMed: 24105322]
5. Rinn J, and Guttman M (2014) RNA Function. RNA and dynamic nuclear organization, *Science* 345, 1240–1241. [PubMed: 25214588]
6. Lesnik C, Golani-Armon A, and Arava Y (2015) Localized translation near the mitochondrial outer membrane: An update, *RNA Biol* 12, 801–809. [PubMed: 26151724]
7. Mili S, and Steitz JA (2004) Evidence for reassociation of RNA-binding proteins after cell lysis: implications for the interpretation of immunoprecipitation analyses, *RNA* 10, 1692–1694. [PubMed: 15388877]
8. Riley KJ, Yario TA, and Steitz JA (2012) Association of Argonaute proteins and microRNAs can occur after cell lysis, *RNA* 18, 1581–1585. [PubMed: 22836356]
9. Shaffer SM, Wu MT, Levesque MJ, and Raj A (2013) Turbo FISH: a method for rapid single molecule RNA FISH, *PLoS One* 8, e75120. [PubMed: 24066168]
10. Crosetto N, Bienko M, and van Oudenaarden A (2015) Spatially resolved transcriptomics and beyond, *Nat. Rev. Genet* 16, 57–66. [PubMed: 25446315]
11. Li Y, Aggarwal MB, Nguyen K, Ke K, and Spitale RC (2017) Assaying RNA Localization in Situ with Spatially Restricted Nucleobase Oxidation, *ACS Chem. Bio* 12, 2709–2714. [PubMed: 28952711]
12. Buettner GR, and Hall RD (1987) Superoxide, hydrogen peroxide and singlet oxygen in hematoporphyrin derivative-cysteine, - NADH and -light systems, *Biochimica et Biophysica Acta (BBA) - General Subjects* 923, 501–507. [PubMed: 3030441]

13. Rougee M, Bensasson RV, Land EJ, and Pariente R (1988) DEACTIVATION OF SINGLET MOLECULAR OXYGEN BY THIOLS AND RELATED COMPOUNDS, POSSIBLE PROTECTORS AGAINST SKIN PHOTOSENSITIVITY, *J. Photochem. Photobiol* 47, 485–489.
14. Maskos Z, Rush JD, and Koppenol WH (1992) The hydroxylation of tryptophan, *Arch. Biochem. Biophys* 296, 514–520. [PubMed: 1321587]
15. Michaeli A, and Feitelson J (1994) REACTIVITY OF SINGLET OXYGEN TOWARD AMINO ACIDS AND PEPTIDES, *J. Photochem. Photobio* 59, 284–289.
16. Jin F, Leitich J, and von Sonntag C (1995) The photolysis ($\lambda = 254$ nm) of tyrosine in aqueous solutions in the absence and presence of oxygen. The reaction of tyrosine with singlet oxygen, *J. Photochem. Photobiol. A: Chem* 92, 147–153.
17. Ali FE, Barnham KJ, Barrow CJ, and Separovic F (2004) Metal catalyzed oxidation of tyrosine residues by different oxidation systems of copper/hydrogen peroxide, *J. Inorg. Biochem* 98, 173–184. [PubMed: 14659647]
18. Bancirova M (2011) Sodium azide as a specific quencher of singlet oxygen during chemiluminescent detection by luminol and Cypridina luciferin analogues, *Luminescence* 26, 685–688. [PubMed: 21491580]
19. Spitale RC, Crisalli P, Flynn RA, Torre EA, Kool ET, and Chang HY (2012) RNA SHAPE analysis in living cells, *Nat. Chem. Bio* 9, 18. [PubMed: 23178934]
20. Spitale RC, Flynn RA, Zhang QC, Crisalli P, Lee B, Jung J-W, Kuchelmeister HY, Batista PJ, Torre EA, Kool ET, and Chang HY (2015) Structural imprints in vivo decode RNA regulatory mechanisms, *Nature* 519, 486. [PubMed: 25799993]
21. Van Nostrand EL, Nguyen TB, Gelboin-Burkhart C, Wang R, Blue SM, Pratt GA, Louie AL, and Yeo GW (2017) Robust, Cost-Effective Profiling of RNA Binding Protein Targets with Single-end Enhanced Crosslinking and Immunoprecipitation (seCLIP), In *mRNA Processing: Methods and Protocols* (Shi Y, Ed.), pp 177–200, Springer New York, New York, NY.
22. Balasubramanian B, Pogozelski WK, and Tullius TD (1998) DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone, *Proc. Natl. Acad. Sci. U. S. A* 95, 9738–9743. [PubMed: 9707545]
23. Lan L, Nakajima S, Wei L, Sun L, Hsieh C-L, Sobol RW, Bruchez M, Van Houten B, Yasui A, and Levine AS (2014) Novel method for site-specific induction of oxidative DNA damage reveals differences in recruitment of repair proteins to heterochromatin and euchromatin, *Nucleic. Acids Res* 42, 2330–2345. [PubMed: 24293652]
24. Ou HD, Phan S, Deerinck TJ, Thor A, Ellisman MH, and O'Shea CC (2017) ChromEMT: Visualizing 3D chromatin structure and compaction in interphase and mitotic cells, *Science* 357, eaag0025. [PubMed: 28751582]
25. G Hendrickson D, Kelley DR, Tenen D, Bernstein B, and Rinn JL (2016) Widespread RNA binding by chromatin-associated proteins, *Genome Biol* 17, 28. [PubMed: 26883116]
26. Taha MS, Nouri K, Milroy LG, Moll JM, Herrmann C, Brunsveld L, Piekorz RP, and Ahmadian MR (2014) Subcellular Fractionation and Localization Studies Reveal a Direct Interaction of the Fragile X Mental Retardation Protein (FMRP) with Nucleolin, *PLOS ONE* 9, e91465. [PubMed: 24658146]
27. Hammond JW, Cai D, and Verhey KJ (2008) Tubulin modifications and their cellular functions, *Curr. Opin. Cell Biol* 20, 71–76. [PubMed: 18226514]
28. Tristan C, Shahani N, Sedlak TW, and Sawa A (2011) The diverse functions of GAPDH: Views from different subcellular compartments, *Cell. Signal* 23, 317–323. [PubMed: 20727968]
29. Wang J, Zhao Y, Wang C, Zhu Q, Du Z, Hu A, and Yang Y (2015) Organelle-Specific Nitric Oxide Detection in Living Cells via HaloTag Protein Labeling, *PLOS ONE* 10, e0123986. [PubMed: 25923693]
30. Rhee H-W, Zou P, Udeshi ND, Martell JD, Mootha VK, Carr SA, and Ting AY (2013) Proteomic Mapping of Mitochondria in Living Cells via Spatially Restricted Enzymatic Tagging, *Science* 339, 1328. [PubMed: 23371551]
31. Roux KJ, Kim DI, and Burke B (2001) BioID: A Screen for Protein-Protein Interactions, In *Curr. Protoc. Protein Sci*, John Wiley & Sons, Inc.

32. Mortensen A, and Skibsted LH (1997) Importance of Carotenoid Structure in Radical-Scavenging Reactions, *J. Agri.Food Chem* 45, 2970–2977.
33. Kaewsapsak P, Shechner DM, Mallard W, Rinn JL, and Ting AY (2017) Live-cell mapping of organelle-associated RNAs via proximity biotinylation combined with protein-RNA crosslinking, *eLife* 6, e29224. [PubMed: 29239719]
34. Egorov SY, Kamalov VF, Koroteev NI, Krasnovsky AA, Toleutaev BN, and Zinukov SV (1989) Rise and decay kinetics of photosensitized singlet oxygen luminescence in water. Measurements with nanosecond time-correlated single photon counting technique, *Chem. Phys. Lett* 163, 421–424.
35. Kuimova MK, Yahioglu G, and Ogilby PR (2009) Singlet Oxygen in a Cell: Spatially Dependent Lifetimes and Quenching Rate Constants, *J. Am. Chem. Soc* 131, 332–340. [PubMed: 19128181]
36. Tyc K, and Steitz JA (1989) U3, U8 and U13 comprise a new class of mammalian snRNPs localized in the cell nucleolus, *The EMBO Journal* 8, 3113–3119. [PubMed: 2531075]
37. Zhou L, Nguyen AN, Sohal D, Ying Ma J, Pahanish P, Gundabolu K, Hayman J, Chubak A, Mo Y, Bhagat TD, Das B, Kapoun AM, Navas TA, Parmar S, Kambhampati S, Pellagatti A, Braunchweig I, Zhang Y, Wickrema A, Medicherla S, Boultonwood J, Platanius LC, Higgins LS, List AF, Bitzer M, and Verma A (2008) Inhibition of the TGF- β receptor I kinase promotes hematopoiesis in MDS, *Blood* 112, 3434. [PubMed: 18474728]
38. Chang T-C, Zeitels LR, Hwang H-W, Chivukula RR, Wentzel EA, Dews M, Jung J, Gao P, Dang CV, Beer MA, Thomas-Tikhonenko A, and Mendell JT (2009) Lin-28B transactivation is necessary for Myc-mediated let-7 repression and proliferation, *Proc. Natl. Acad. Sci. U. S. A* 106, 3384–3389. [PubMed: 19211792]
39. Werner Michael S., and Ruthenburg Alexander J. (2015) Nuclear Fractionation Reveals Thousands of Chromatin-Tethered Noncoding RNAs Adjacent to Active Genes, *Cell Reports* 12, 1089–1098. [PubMed: 26257179]

Figure 1. Singlet oxygen-based protein tagging.

A. Imaging results after cells being treated with nothing, DNase, RNase and proteinase, respectively. DBF denoted in green, PA adducts denoted in red, and nuclear stained with Hoechst and shown in blue. B. Schematic of protein tagging with singlet oxygen. C. Denaturing gel-imaging with Cy5; E represents "Experimental", which has TBF, PA and hv, while three other headers represent different negative control conditions. D. Time course of adduct formation (red: over-saturated signal). E. Quenching of protein tagging with NaN_3 . CB: Coomassie blue.

Figure 2. Spatially-restricted singlet oxygen-based protein tagging.

A. Schematic of spatially-restricted chromatin tagging with singlet oxygen. B. Biotin-Streptavidin enrichment-western blot analysis of tagged proteins with the H2B-Halo fusion construct.

Figure 3. RT-qPCR analysis of RNAs tagged by enrichment of RNA-protein complexes.
A. RT-qPCR comparison of the previous method IBE and the current method IAE. B. Reported ratio of Chromatin Pellet Extract (CPE) to Soluble-Nuclear Extract (SNE)⁽³⁹⁾. C. RT-qPCR comparison of H2B-Halo and NLS-Halo constructs by IAE approach. D. Fluorescence imaging signals of Cy5 azide (PA adducts denoted in red) for H2B-Halo and NLS-Halo with yellow arrows pointing at enhanced nucleolus signal in NLS but lack of signal in H2B. Enrichment was calculated against negative control with Ot method (SI), biological duplicates.

Scheme 1. Assaying RNA localization with spatially-restricted oxidation.

A. Schematic of RNA oxidation with fluorophores (DBF, shown in green) and subcellular RNA tagging with PA. B. Schematic of comparison between previous work and this work. In previous work, free tagged RNAs were isolated and profiled. In this work, pulldown of RNA-protein complexes increased the yield of RNA enrichment. POI: protein of interest; $h\nu$: light irradiation.