Published in final edited form as:

Toxicol Appl Pharmacol. 2018 April 01; 344: 74. doi:10.1016/j.taap.2018.03.001.

Corrigendum to “Convergence of hepcidin deficiency, systemic iron overloading, heme accumulation, and REV-ERB α/β activation in aryl hydrocarbon receptor-elicited hepatotoxicity” [Toxicol. Appl. Pharmacol. 321 (2017) 1–17]

K.A. Fader^{a,b}, R. Nault^{a,b}, M.P. Kirby^a, G. Markous^a, J. Matthews^c, and T.R. Zacharewski^{a,b,*}

^aDepartment of Biochemistry & Molecular Biology, Michigan State University, East Lansing, MI 48824, United States

^bInstitute for Integrative Toxicology, Michigan State University, East Lansing, MI 48824, United States

^cDepartment of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Oslo 0316, Norway

After publication of this manuscript, the authors were notified by Abcam that the antibody used to quantify hepatic protein levels of iron responsive element (IRE)-binding protein 2 (IREB2; ab181153) was determined to detect not only IREB2 as marketed, but also aconitase 1 (ACO1; aka IREB1). As a result, Fig. 4B depicts total hepatic levels of both IRE-binding proteins (ACO1 and IREB2) rather than IREB2 specifically. However, both ACO1 and IREB2 regulate transcription/translation of IRE-containing genes through the same mechanism. Therefore, this does not significantly impact our conclusions regarding the effect of TCDD on IRE-binding proteins or overall dysregulation of intracellular iron homeostasis.

The authors would like to apologise for any inconvenience caused.

*Corresponding author at: Michigan State University, 603 Wilson Road, Room 309, East Lansing, MI 48824, United States., tzachare@msu.edu (T.R. Zacharewski).