CARE COORDINATION AND SOCIAL NETWORK DYNAMICS IN GERIATRIC CARE USING WEARABLE PROXIMITY SENSORS

S. Wei¹, K. Corazzini², E. McConnell¹, 1. *Duke University School of Nursing*, 2. *Duke University*

Background: Care coordination involves multiple participants (healthcare providers and patients) working together to ensure the appropriate delivery of care. Face-to-face interactions between individuals is a critical component of care coordination in clinical settings. An accurate and detailed description and quantification of contacts provide key information about how providers interact with patients and for intervention development to improve care coordination. Using proximity data collected from Radio-Frequency Identification (RFID) wearable sensors, this study aimed to understand network structure and interaction patterns in relation to different positional roles in care coordination.

Methods and Findings: Social network analysis was used to analyze a secondary dataset collected from RFID sensors that detect close-range face-to-face interactions between individuals in a geriatric unit of a university hospital in France. Two questions were explored: (1) how do interaction patterns differ related to different roles (nursing staff, doctors, patients and administrative staff)? And (2) how do the interaction patterns differ within the same roles on the geriatric unit? Over 4 days and nights, interaction patterns and network measures, such as centrality and cohesion, among providers ($n = 46$) and patients ($n = 29$) varied greatly depending on the shifts of day and different clinical roles.

Conclusion: Wearable sensors provide a novel way to objectively capture network data to understand ego-centric and system-level interaction patterns in care coordination. Real-time longitudinal measurement of face-to-face interactions can provide important information on structural mechanisms in relation to different role responsibilities and functions in the context of care coordination in geriatric clinical care settings.

CAREGIVER CHALLENGES AFTER A DRIVING ASSESSMENT CLINIC RECOMMENDATION TO CEASE DRIVING: IMPLICATIONS FOR SUPPORT

C. Pierpaoli Parker¹, M. Cox², C. Owsley³, J. Elgin², A. McBrayer², L. Dreer², 1. *University of Alabama*, 2. *Department of Ophthalmology and Visual Sciences, University of Alabama at Birmingham*, 3. *University of Alabama at Birmingham*

Purpose: To examine the challenges family caregivers are confronted with following a recommendation for their relative to cease driving.

Methods: The sample consisted of 63 caregivers, ages 38 to 93 years old ($M = 61.40$, $SD = 11.20$; 69.8% female; 90.5% Caucasian), who provided assistance to an adult recommended to cease driving following an evaluation at a Driving Assessment Clinic (DAC) for adults with medical conditions. Participants were contacted at least 2 months post-evaluation in order to have had time to experience potential caregiver challenges following the recommendation. Caregivers completed a semi-structured needs assessment survey via telephone. For the current study, qualitative responses regarding the unique challenges caregivers found most distressing following their relative's driving evaluation

were coded into themes separately by two researchers, with a third researcher resolving any discrepancy.

Results: 70% of participants reported 1 or more distressing challenges related to caregiving. Fourteen unique themes emerged with the top 5 being: (1) caregiver emotional distress, (2) time constraints, (3) difficulty managing care-recipient emotions and behaviors (e.g., arguments around continuing to drive or transportation alternatives), and (4) problems with coordinating and distributing caregiving tasks. Providing assistance with instrumental activities of daily living (IADLs) and family relationship strain both tied as the fifth theme.

Conclusions: Caregivers experience a variety of challenges following a relative's recommendation to cease driving. The findings will be discussed in terms of enhancing provider communication about such challenges, available resources, and ongoing support programs to minimize family stress.

CAREGIVER IDENTITY THEORY AND PREDICTORS OF BURDEN AND DEPRESSION: FINDINGS FROM THE REACH II STUDY

V. Miller¹, M. Killian², N. Fields¹, 1. *The University of Texas at Arlington*, 2. *Florida State University*

Due to the degenerative nature of Alzheimer's disease (AD), caregivers (CGs) of persons with dementia face many challenges. CGs are often subject to stressors, placing them at a greater risk for depression and other mental health problems. Guided by the caregiver identity theory, we examine the relationship between care recipient's ability to perform daily tasks, ADLs and IADLs, and CG depression as mediated by CGs' perceived burden while caregiving for one with AD. Using baseline data from the REACH II study, relationships between burden, reported depression, and each ADLs and IADLs were tested. To explore CG burden mediating the relationship between and depression each ADLs and IADLs, structural equation modeling (SEM) was conducted. Reported CG total scores were significantly correlated with total ADLs (Spearman's $\rho = .524$, $p < .001$) and IADLs ($\rho = .495$, $p < .001$) scores, indicating increased perceived CG burden with greater number of assisted daily activities. CG depression scores were significantly predicted by burden scores ($\beta = 0.386$, $t = 18.27$, $p < .001$, $r^2 = .346$) and CG's ability to pay for basic needs ($\beta = 1.10$, $t = 4.50$, $p < .001$, $r^2 = .031$). SEM modeling indicated that CG burden fully mediated the relationship between ADLs and IADLs and CG depression after controlling for CG's ability to pay for basic care needs. Our results shed light on the relationship between a CG's perceived burden and depression. Findings suggest the need to closely examine the relationships between AD caregiving, CG burden, and mental health. Results suggest bolstering policies and programs that offer tangible supports and services to ameliorate CG burden of informal AD caregiving.

CHALLENGES IN ASSESSMENT AND MANAGEMENT OF ALCOHOL USE DISORDER IN OLDER ADULTS

M. DiBartolo, J. Jarosinski, *Salisbury University*

Alcohol use disorder (AUD) is a significant, under-recognized, under-reported diagnosis and a hidden epidemic in the U.S. The prevalence is expected to rise dramatically as the number of adults aged 65 and older increases to one in five by 2030 and in this group, the health consequences