CORRESPONDENCE

Bowlegs and Intensive Football Training in Children and Adolescents

A Systematic Review and Meta-Analysis

by Dr. med. Peter Helmut Thaller, Dr. med. Julian Fürmetz, Fuhuan Chen, Dr. med. Nikolaus Degen, Kirs Marjaana Manz, Dipl.-Phys., and M. Sc., Dr. med. Florian Wolf in issue 24/2018

No Obvious Association

The article (1) spells doom for football, and perhaps other sports during adolescence: football causes genu varum (bowlegs), and bowlegs in adolescents turns into knee osteoarthritis in adults. Only: this is not supported by the data, and neither is there any evidence for the consequences. The meta-analysis included case-control studies, which without longitudinal studies are not suitable for establishing causalities (in this case: between football and bowlegs).

The association between bowlegs and later knee osteoarthritis is not obvious. In the cited review by Tanamas et al. (2), only 1 of 14 included studies found an increased risk for the development of knee osteoarthritis as a result of leg malalignment, and only in overweight persons. Other longitudinal studies—for example, that reported by Hunter et al. (3)—did not find any increase in risk.

The orthopedic myth, that bowlegs and genu valgum deformity (“knock-knees”) lead to knee osteoarthritis, results from the ex post view—namely, the pronounced malalignment of the leg axis in knee osteoarthritis. The conclusion: bowlegs and knock-knees don’t cause knee osteoarthritis, but knee osteoarthritis causes bowlegs and knock-knees.

An intercondylar distance of more than 1.5 cm in young people who play football should not be construed into a drama. Cruciate ligament injury and especially cartilage loss—whether as a result of traumatic injury or iatrogenic—advance knee osteoarthritis substantially. And: in addition to age, overweight and obesity also have a confirmed effect in terms of advancing knee osteoarthritis (4).

DOI: 10.3238/arztebl.2018.0683a

References

1. Thaller PH, Fürmetz J, Chen F, Degen N, Manz KM, Wolf F: Bowlegs and intensive football training in children and adolescents—a systematic review and meta-analysis. *Dtsch Arztebl Int* 2018; 115: 401–8.
2. Tanamas S, Hanna FS, Cicuttini FM, Wluka AE, Berry P, Urquhart DM: Does knee malalignment increase the risk of development and progression of knee osteoarthritis? A systematic review. *Arthritis Rheum* 2009; 61: 459–67.
3. Hunter DJ, Niu J, Felson DT, et al.: Knee alignment does not predict incident osteoarthritis: the Framingham osteoarthritis study. *Arthritis Rheum* 2007; 56: 1212–8.
4. Schiltenswolf M, Liebers F, Bolm-Audorff U, et al.: Welche konkurrierenden Faktoren sind bei der Begutachtung der Gonarthrose zu berücksichtigen? In: Schiltenswolf M, Grosser V, Thomann KD (eds.): *Berufskrankheit Gonarthrose (BK 2112)*. Frankfurt am Main: Referenz Verlag 2012.

Prof. Dr. med. Marcus Schiltenswolf

Zentrum für Orthopädie, Unfallchirurgie und Paraplegiologie, Universitätsklinikum Heidelberg
marcus.schiltenswolf@urz.uni-heidelberg.de

The Mikulicz Line as a Parameter

The article on the association of genu varum (bowlegs) and intensive football training during childhood and adolescence mentions that epidemiological studies in children and adolescents require clinical investigative methods. However, much space is then given to radiological criteria in terms of text and figures (1). In addition to some ambiguities in the naming of the radiological parameters mentioned in the figures, any mention of the Mikulicz line (the mechanical longitudinal axis of the leg) is conspicuous by its absence—in text, tables, and figures—although the Mikulicz line is one of the crucial radiological parameters that provides immediate insights into the load distribution in the knee joint, in a scenario of deviations of the leg axis.

Curiously, this mechanical axis is then depicted in *Figure 5* as the only measurement line in the whole-leg photographs, but it is not explained in the legend—namely, as the mechanical load bearing line of the leg at the frontal level, as the link between the hip and ankle joints, with the resultant intersection in the knee joint, which describes the clinical relevance. For planning operative correction of the mechanical axis, the Mikulicz line is also the crucial parameter (2). This is obviously also the case in the clinical case shown in *Figure 5*, because the clinical picture shows for the worse affected left side a preoperatively shaved leg, evidently preceding valgization osteotomy. In addition to operative aspects, the Mikulicz line is generally an important measure for assessing leg axis deviations in children and adolescents in very different circumstances (3) and is mentioned as the mechanical axis of the leg in the cited references of the article too. The association between the clinically and biologically relevant changes in the load distribution in the compartments of the knee joint and the shifting of the Mikulicz line in axis deviations, as well as before and after corrective surgery, has been described elsewhere (4).

DOI: 10.3238/arztebl.2018.0683b

References

1. Thaller PH, Fürmetz J, Chen F, Degen N, Manz KM, Wolf F: Bowlegs and intensive football training in children and adolescents—a systematic review and meta-analysis. *Dtsch Arztebl Int* 2018; 115: 401–8.

2. Lobenhoffer P, Van Heerwaarden R, Agneskirchner JD: Kniegelenknahe Osteotomien: Indikation- Planung- Operationstechniken mit Plattenfixateuren. New York: Georg Thieme Verlag, 2nd revised edition 2014.
3. Landauer F, Huber G, Paulmichl K, O'Malley G, Mangge H, Weghuber D: Timely diagnosis of malalignment of the distal extremities is crucial in morbidly obese juveniles. *Obes Facts* 2013; 6: 542–51.
4. Agneskirchner JD, Hurschler C, Wrann CD, Lobenhoffer P: The effects of valgus medial opening wedge high tibial osteotomy on articular cartilage pressure of the knee: a biomechanical study. *Arthroscopy* 2007; 23: 852–61.

Prof. Dr. med. Andreas Lahm MPH

Kliniken Maria Hilf GmbH Mönchengladbach
Akademisches Lehrkrankenhaus der Uniklinik RWTH Aachen
Andreas.Lahm@mariahilf.de

In Reply:

In our article, we investigated indications that football as a competitive sport in adolescents increases the risk of developing genu varum (bowlegs). Leisure time sport is explicitly excluded, and the great importance of sport for the development of adolescents was emphasized (1). In accordance with the article's topic and the timing of its publication, immediate reporting in the media was extensive, as expected. Lively (we hope) scientific discussion will take its time to run its course. We therefore wish to express our particular thanks for the feedback from specialists and would respond to our correspondents as follows.

We thank Prof. Lahm for his additions to our article. It is correct that the longitudinal axis of the leg constitutes the central topic of our article. It is the most important parameter in the context of deformities close to the knee joint in the clinical/anatomical and radiological contexts. The terms (mechanical) leg axis or Mikulicz line are used synonymously, alongside other terms. A long standing radiograph is the gold standard for determining the leg's mechanical axis (2). But this is not an option for studies of healthy adolescents. We therefore evaluated only clinical studies in our article. We were not able to give "much space" to the "radiological criteria" for reasons of restricting ourselves to include the essentials: in only one sentence on page 403 and in a paragraph on page 406. The mechanical axis of the leg was not shown/labeled in *Figure 2* for reasons of clarity. It could have been mentioned in the described clinical case on page 406. But here, too, the positive late result after bilateral realignment (with a clinically

and radiologically centered longitudinal axis in both legs) fell victim to space restrictions.

We also thank Prof. Schiltewolf for his comments. Our article does indeed include a cautious warning. Cautious, because the literature on the topic is scarce, warning because the meta-analysis indicates a clear direction. When discussing the relevance of genu varum, we cautiously explained the hypothesis that, on the basis of the existing evidence, deviations of the leg axis encourage knee osteoarthritis. A reference not cited in the article for reasons of space supports this hypothesis also on the basis of long standing radiographs (3). If a clear contradiction to this hypothesis is supported merely by the cited 2007 article "Knee alignment does not predict incident osteoarthritis," then methodological weaknesses of this article should be mentioned. A limitation of the so-called Framingham osteoarthritis study lies in the determination of the leg axis by using focused views of the knee joint only. We wish to reiterate that in contrast to the long standing radiograph (2), focused views of unilateral knee joints are less appropriate for determining the leg's mechanical axis (4).

In conclusion, further, high quality studies are needed.

DOI: 10.3238/arztebl.2018.0684

References

1. Thaller PH, Fürmetz J, Chen F, Degen N, Manz KM, Wolf F: Bowlegs and intensive football training in children and adolescents—a systematic review and meta-analysis. *Dtsch Arztebl Int* 2018; 115: 401–8.
2. Thaller PH, Baumgart R, Burghardt R, Knuellig S, Buerklein D, Mutschler W: Digital imaging in lower limb bone deformities—standards and new perspectives. *ELSEVIER International Congress Series* 2005; 1281: 154–8.
3. Sharma L, Song J, Felson DT, Cahue S, Shamiyeh E, Dunlop DD: The role of knee alignment in disease progression and functional decline in knee osteoarthritis. *JAMA* 2001; 286: 188–95.
4. Zampogna B, Vasta S, Amendola A, et al.: Assessing lower limb alignment: comparison of standard knee xray vs long leg view. *Iowa Orthop J* 2015; 35: 49–54.

On behalf of the authors

Dr. med. Peter Helmut Thaller

Arbeitsgruppe 3D-Chirurgie
Klinik für Allgemeine, Unfall- und Wiederherstellungschirurgie
Klinikum der Ludwig-Maximilians-Universität München
Campus Innenstadt, pthaller@med.lmu.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.