CORRESPONDENCE

Oxygen Treatment in Intensive Care and Emergency Medicine

by Dr. med. Jörn Grensemann, PD Dr. med. Valentin Fuhrmann, and Prof. Dr. med. Stefan Kluge in issue 27–28/2018

O₂ saturation target of 96–100% should be abandoned

We thank the authors for raising this important topic (1). The meta-analysis by Chu et al. showed that oxygen confers no overall benefit in normotoxic patients and is associated with increased mortality in patients at risk. The risk of hypercapnic respiratory failure induced by hyperoxia has been described not only in patients with chronic obstructive pulmonary disease (COPD), but, among others, also in patients with community acquired pneumonia, severe asthma, and obesity hypoventilation syndrome. Although oxygen is a drug by definition, 42% of UK hospital patients receiving oxygen had no prescription for it (2).

An O₂ saturation measured by pulse oximetry (SpO₂) of below 92% corresponds to an arterial saturation (SaO₂) lower than 90% with a high sensitivity and specificity. In patients with SpO₂ < 92%, those at risk for hypercapnia, and critically ill patients, arterial blood gases are required. In more than 97% of the population at sea level, SaO₂ at rest is above 92%. Even patients with comorbidities tolerate SpO₂ saturations between 80% and 90% for long periods. Only if SaO₂ is continuously < 85%, indications are found of anaerobic metabolism in the myocardium.

In order to define a hyperoxic range, the Australian guideline suggests a target SpO₂ range of 92–96% with oxygen therapy. Patients at risk for hypercapnia should be given oxygen only if their SpO₂ is < 88% (SpO₂ target range then 88–92%), all other patients if their SpO₂ is < 92% (3).

In contrast, many patients in German emergency departments and intensive care units have SpO₂ values above 96% during oxygen therapy. In electronic charting systems at Hannover Medical School (MHH), an SpO₂ “normal range” of 96–100% has recently been completely abandoned.

A target value for SpO₂ or SaO₂ should be defined in all inpatients receiving oxygen supplementation. Oxygen administration by automated titration systems is promising (4). Careful administration of oxygen may also save costs.

DOI: 10.3238/arztebl.2018.0685a

References

1. Grensemann J, Fuhrmann V, Kluge S: Oxygen treatment in intensive care and emergency medicine. *Dtsch Arztebl Int* 2018; 115: 455–62.
2. O'Driscoll R. British Thoracic Society, Emergency Oxygen Audit Report 2015. www.brit-thoracic.org.uk/document-library/audit-and-quality-improvement/audit-reports/bts-emergency-oxygen-audit-report-2015/ (last accessed on 15 July 2018).
3. Beasley R, Chien J, Douglas J, et al.: Thoracic Society of Australia and New Zealand oxygen guidelines for acute oxygen use in adults: ‘Swimming between the flags’. *Respirology* 2015; 20: 1182–91.

4. L'Her E, Dias P, Gouillou M, et al.: Automatic versus manual oxygen administration in the emergency department. *Eur Respir J* 2017; 50: 1602552.

Prof. Dr. med. Jens Gottlieb

Klinik für Pneumologie, Deutsches Zentrum für Lungenforschung (DZL)
gottlieb.jens@mh-hannover.de

Dr. med. Anna Bertram

Klinik für Nieren- und Hochdruckerkrankungen, Zentrale Notfallaufnahme

Christoph Duesberg

Klinik für Pneumologie, Zentrale Notfallaufnahme

Dr. med. Gernot Beutel

Klinik für Hämatologie, Hämostaseologie, Onkologie und Stammzelltransplantation, Zentrale Notfallaufnahme
Medizinische Hochschule Hannover (MHH)

Conflict of interest statement

The authors declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

In Reply:

We thank our correspondents for their valuable additional comments on our article (1). The meta-analysis by Chu et al., which was published just before our article went into print, underlines clearly that hyperoxia can be associated with increased mortality (2). On this background, the administration of oxygen without prescription has to be regarded as a particular worry and should prompt intensive training measures.

As we described in our article, specialty societies have issued clear recommendations for the target ranges for oxygen supplementation for different conditions. Future studies should investigate these target ranges and optimize them.

For the best possible treatment of our patients it is crucial to put into practice what is currently known about the negative effects of hyperoxia. For this reason we welcome and support explicitly that staff in all hospital areas are alerted to the problem of iatrogenic hyperoxia, so as to avoid unnecessary administration of oxygen leading to increases in morbidity and mortality.

Alerts can, for example, be triggered by electronic devices, such as pre-set gas monitoring alarm ranges, which go off as soon as pulse oximetry identifies an oxygen saturation (SpO₂) above a certain limit, and which can be facilitated by SpO₂ targets that are patient-specific and individual. Automated titration systems are promising in this setting, but no results from studies exist to date. Structured implementation is a further important building block when setting physiological SpO₂ targets. This could consist, for example, of specifically training the staff immediately involved in treatment and using electronic monitoring systems (3).

DOI: 10.3238/arztebl.2018.0685b

References

1. Grensemann J, Fuhrmann V, Kluge S: Oxygen treatment in intensive care and emergency medicine. *Dtsch Arztebl Int* 2018; 115: 455–62.
2. Chu DK, Kim LH, Young PJ, et al.: Mortality and morbidity in acutely ill adults treated with liberal versus conservative oxygen therapy (IOTA): a systematic review and meta-analysis. *Lancet* 2018; 391: 1693–705.
3. Helmerhorst HJ, Schultz MJ, van der Voort PH, et al.: Effectiveness and clinical outcomes of a two-step implementation of conservative oxygenation targets in critically ill patients: a before and after trial. *Crit Care Med* 2016; 44: 554–63.

Dr. med. Jörn Grensemann

PD Dr. med. Valentin Fuhrmann

Prof. Dr. med. Stefan Kluge

Klinik für Intensivmedizin, Universitätsklinikum Hamburg-Eppendorf
skluge@uke.de

Conflict of interest statement

Prof. Kluge received consultancy fees from Baxter, Fresenius, and Xenios. He received reimbursement of travel and accommodation expenses as well as lecture fees from Baxter, Fresenius, Sorin, and Xenios. He received consumables for the conduct of clinical and preclinical studies from ETVision Ltd and Fisher & Paykel as well as funds from Xenios.

Dr. Grensemann and Dr. Fuhrmann declare that no conflict of interests exists.

CLINICAL SNAPSHOT

STEMI Equivalent in Ventricular Pacemaker Rhythm

A 90-year-old female patient was brought to our emergency department complaining of chest pain. Ventricular paced rhythm was detected on 12-lead ECG. The widespread opinion that this constellation does not permit a diagnosis of ischemia on 12-lead ECG is not accurate, as this example shows. In lead V2 the patient showed concordant (on the same side of the QRS main vector) ST elevations >1 mm. Discordant (on the opposite side of the QRS main vector) ST elevations and depressions are not pathological in ventricular paced rhythm. However, the discordant ST elevations in lead V3 were too pronounced (ratio of ST elevation at the J-point to the S-wave approximately 1:1 and thus clearly $\geq 25\%$). Thus, according to Sgarbossa and Smith-modified Sgarbossa criteria, the ECG findings corresponded to STEMI equivalent of the anterior wall. As expected, an occlusion of the middle segment of the left anterior descending artery was diagnosed and successfully recanalized on emergent coronary angiography. Both in ventricular paced rhythm and in left bundle branch block, chest pain should prompt 12-lead ECG to scrutinize for signs of ischemia according to the above-mentioned criteria or, if applicable, other criteria.

Dr. med. Steffen Grautoff, EBCEM, Zentrale Notaufnahme, Klinikum Herford, steffen.grautoff@klinikum-herford.de

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare.

Cite this as: Grautoff S: STEMI equivalent in ventricular pacemaker rhythm. *Dtsch Arztebl Int* 2018; 115: 686. DOI: 10.3238/arztebl.2018.0686